



**STATE OF CONNECTICUT
DEPARTMENT OF ENERGY AND ENVIRONMENTAL
PROTECTION**

Robert Klee
Commissioner

Bureau of Natural Resources
Marine Fisheries Division
www.ct.gov/deep/fishing

**A STUDY OF MARINE RECREATIONAL
FISHERIES IN CONNECTICUT**



Federal Aid in Sport Fish Restoration
F14AF00296 (F-54-R-34)
Annual Performance Report
March 1, 2014 – February 28, 2015



Job 5 (LIS Trawl Survey)

Cover photo: *Fisheries Biologist (retired) Roderick MacLeod with a trophy striped bass he caught in Long Island Sound.*

Roderick (Rod) MacLeod retired on April 30, 2015, after more than 34 years of service with the Marine Fisheries Division. During that time Rod helped initiate the marine angler survey program including the Volunteer Angler Program – one of the first of its kind on the east coast. Rod headed up the marine angler survey for nearly 30 years including managing the transition from Connecticut's independent creel surveys to participation in the Marine Recreational Fishery Statistics Survey in 1987. Rod also contributed significantly to the design and implementation of our return to independent creel surveys in 2013 intended to complement the federal MRIP survey.

In addition to oversight of state marine angler surveys, Rod served as Connecticut's representative on ASMFC Interstate Tagging and Artificial Reef Committees and on the ACCSP Operations and Recreational Technical Committees.

Rod's day-to-day duties also included heading up fish kill investigations and looking out for anglers' interests in the agency's review of structures and dredging permits, including providing critical advice on the development of fishing access sites whenever such opportunities arose. Rod also played a central role in our outreach efforts targeting the angling community, including producing the weekly fishing reports and helping hundreds of callers over the years looking for a good place to get out fishing or crabbing. His detailed knowledge of fishing and fishing access along the entire coast, gleaned from years of creel survey work and his own extensive fishing experience, has been invaluable to this office and the fishing public.

Rod will be missed not only for the contributions he has made to this agency over more than a third of a century, but also as a longtime friend and colleague. We wish Rod and his family the best as he enters this new and exciting phase in his life in retirement. Try to leave a few fish on the shoal for us weekend warriors, Rod!

State of Connecticut
Department of Energy and Environmental Protection
79 Elm Street
Hartford, CT 06106-5127
www.ct.gov/deep

Federal Aid in Sport Fish Restoration
F14AF00296 (F-54-R-34)
Annual Performance Report

Project Title: *A Study of Marine Recreational Fisheries in Connecticut*

Period Covered: March 1, 2014 - February 28, 2015

Job Title

Job 1: Marine Angler Survey
Job 2: Volunteer Angler Survey
Job 3: Enhanced Shore Fishing
Job 4: Tackle Shop Co-op Survey
Job 5: Marine Finfish Survey

Job 6: Studies in Conservation Engineering
Job 7: Alosine Survey
Job 8: Estuarine Seine Survey

Job 9: Volunteer Estuarine Fisheries Database
Job 10: Cooperative Interagency Resource Monitoring

Job 11: Public Outreach
Job 12: Marine Fisheries GIS

Prepared by:

Roderick E. MacLeod
Roderick E. MacLeod
Gregory Wojcik
Gregory Wojcik
Kurt F. Gottschall
Deborah J. Pacileo
Inactive
Jacqueline M. Benway
David R. Molnar
Penelope T. Howell
Penelope T. Howell
Matthew J. Lyman
Katie O'Brien-Clayton
David R. Molnar
Deborah J. Pacileo
Jacqueline M. Benway



Approved by:

David G. Simpson, Director
Marine Fisheries Division

Date: May 1, 2015

JOB 5: MARINE FINFISH SURVEY

Long Island Sound Trawl Survey

LONG ISLAND SOUND TRAWL SURVEY

TABLE OF CONTENTS

LIST OF TABLES	iv
LIST OF FIGURES	vi
Cruise results from the 2014 Spring & fall surveys	1
STUDY PERIOD AND AREA.....	1
GOAL.....	1
OBJECTIVES.....	1
INTRODUCTION.....	2
METHODS.....	2
Sampling Design	2
Sampling Procedures	3
Data Analysis.....	5
<i>Indices of Abundance: Annual Mean Count and Weight per Tow</i>	5
<i>Indices of Abundance: Indices-at-Age and Age Group</i>	5
<i>Species Richness by Group</i>	8
<i>Open Water Forage Abundance</i>	8
RESULTS AND DISCUSSION.....	9
Overview of LISTS 2014 Spring and Fall Surveys.....	9
Cooperative Sample and Data Collection.....	9
Number of Species Identified.....	10
Total Catch	10
Length Frequencies	10
Seasonal Indices of Abundance.....	11
Indices of Abundance: Important Recreational Species.....	11
Winter Flounder Average Size at Maturity	12
Species Richness by Group	12
MODIFICATIONS.....	12
LITERATURE CITED.....	13
TABLES 5.1 - 5.29	15
TABLES 5.30 - 5.65 (Length Frequencies)	43
FIGURES 5.1 - 5.19	81
APPENDICES.....	101
Appendix 5.1. List of finfish species identified by <i>A Study of Marine Recreational Fisheries in Connecticut (F54R)</i> and other CT DEEP Marine Fisheries Division programs.....	102

Appendix 5.2. Annual total count of finfish, lobster and squid taken in the LISTS, 1984-2014..... 105

Appendix 5.3. Annual total weight (kg) of finfish, lobster and squid taken in LISTS, 1992-2014..... 108

Appendix 5.4. Total number and weight (kg) of finfish and invertebrates caught in LISTS, 1984-2014. 110

Appendix 5.5. Endangered Species Interactions 140

Appendix 5.6. Cold and warm temperate species captured in LISTS 141

LIST OF TABLES

Table 5.1.	Specifications for the Wilcox 14 m high-rise trawl net and associated gear.....	16
Table 5.2.	The number of sites scheduled for sampling each month within the 12 depth-bottom type strata.....	16
Table 5.3.	Length and age data collected in 2014.	17
Table 5.4.	Number of Long Island Sound Trawl Survey (LISTS) samples taken by year and cruise.....	18
Table 5.5.	Station information for LISTS April 2014.	19
Table 5.6.	Station information for LISTS May 2014.	20
Table 5.7.	Station information for LISTS June 2014.	21
Table 5.8.	Station information for LISTS September 2014.....	22
Table 5.9.	Station information for LISTS October 2014.....	23
Table 5.10.	Samples with non-standard tow durations and reason for incomplete tow, spring and fall 2014.....	24
Table 5.11.	Data requests by month, 2014.	25
Table 5.12.	Sample requests by month, 2014.....	26
Table 5.13.	List of finfish species observed in 2014.	27
Table 5.14.	List of invertebrates observed in 2014.	28
Table 5.15.	Total number and weight (kg) of finfish and invertebrates caught in 2014.	29
Table 5.16.	Total counts and weight (kg) of finfish taken in the spring and fall sampling periods, 2014.	30
Table 5.17.	Total catch of invertebrates taken in the spring and fall sampling periods, 2014.	31
Table 5.18.	Spring indices of abundance for selected species, 1984-2014.	32
Table 5.19.	Fall indices of abundance for selected species, 1984-2014.....	33
Table 5.20.	Finfish and invertebrate biomass indices for the spring sampling period, 1992-2014.	34
Table 5.21.	Finfish and invertebrate biomass indices for the fall sampling period, 1992-2014.....	35
Table 5.22.	Bluefish indices of abundance, 1984-2014.....	36
Table 5.23.	Scup indices at-age, 1984-2014.....	37
Table 5.24.	Age frequency of striped bass taken in spring, 1984-2014.	38
Table 5.25.	Striped bass indices-at-age, 1984-2014.	38
Table 5.26.	Summer flounder indices-at-age, 1984-2014.	39
Table 5.27.	Tautog indices-at-age, 1984-2014.	40
Table 5.28.	Weakfish age 0 and age 1+ indices of abundance, 1984-2014.....	41
Table 5.29.	Winter flounder indices-at-age, 1984-2014.....	42
Table 5.30.	Alewife length frequencies, spring and fall, 1 cm intervals, 1989-2014.....	44
Table 5.31.	American shad length frequencies, spring and fall, 2 cm intervals (midpoint given), 1989-2014.....	45
Table 5.32.	American lobster length frequencies - spring, female, 1 mm intervals, 1984-2014.....	46
Table 5.33.	American lobster length frequencies - fall, female, 1 mm intervals, 1984-2014.	47
Table 5.34.	American lobster length frequencies - spring, male, 1mm intervals, 1984-2014.....	48
Table 5.35.	American lobster length frequencies - fall, male, 1 mm intervals, 1984-2014.....	49

Table 5.36. Atlantic herring length frequencies, spring and fall, 1 cm intervals, 1989-2014.	50
Table 5.37. Atlantic menhaden length frequency, spring and fall, 1 cm intervals, 1996-2014.	51
Table 5.38. Black sea bass length frequency, spring, 1 cm intervals, 1987-2014....	52
Table 5.39. Black sea bass length frequency, fall, 1 cm intervals, 1987-2014.....	53
Table 5.40. Blueback herring length frequencies, spring and fall, 1 cm intervals, 1989-2014.	54
Table 5.41. Bluefish length frequencies, spring, 1 cm intervals (midpoint given), 1984-2014.....	55
Table 5.42. Bluefish length frequencies, fall, 1 cm intervals (midpoint given), 1984-2014.	56
Table 5.43. Butterfish length frequencies, 1 cm intervals, spring and fall, 1986-1990, 1992-2014.....	57
Table 5.44. Clearnose skate length frequencies, spring, 1 cm intervals, 1993-2014.....	58
Table 5.45. Clearnose skate length frequencies, fall, 1 cm intervals, 1993-2014	59
Table 5.46. Fourspot flounder length frequencies, spring and fall, 2 cm intervals (midpoint given), 1989, 1990, 1996-2014.....	60
Table 5.47. Hickory shad length frequencies, spring and fall, 1 cm intervals, 1991-2014.	61
Table 5.48. Horseshoe crab length frequencies by sex, spring, 1 cm intervals, 1998-2014....	62
Table 5.49. Horseshoe crab length frequencies by sex, fall, 1 cm intervals, 1998-2014.....	63
Table 5.50. Long-finned squid length frequencies, spring, 1 cm intervals (midpoint given), 1986- 1990, 1992-2014.....	64
Table 5.51. Long-finned squid length frequencies, fall, 1 cm intervals (midpoint given), 1986-1990, 1992-2014.....	65
Table 5.52. Scup spring length frequencies, 1 cm intervals, 1984-2014.....	66
Table 5.53. Scup fall length frequencies, 1 cm intervals, 1984-2014.....	67
Table 5.54. Striped bass spring length frequencies, 2 cm intervals (midpoint given), 1984-2014.....	68
Table 5.55. Striped bass fall length frequencies, 2 cm intervals (midpoint given), 1984-2014.	69
Table 5.56. Summer flounder length frequencies, spring, 2 cm intervals (midpoint given), 1984-2014.....	70
Table 5.57. Summer flounder length frequencies, fall, 2 cm intervals (midpoint given), 1984-2014.....	71
Table 5.58. Tautog length frequencies, spring, 1 cm intervals (midpoint given), 1984-2014.....	72
Table 5.59. Weakfish length frequencies, spring, 2 cm intervals (midpoint given), 1984-2014.....	73
Table 5.60. Weakfish length frequencies, fall, 2 cm intervals (midpoint given), 1984-2014.	74
Table 5.61. Windowpane flounder length frequencies, spring, 1 cm intervals, 1989, 1990, 1994-2014.....	75
Table 5.62. Windowpane flounder length frequencies, fall, 1 cm intervals, 1989, 1990, 1994-2014.....	76
Table 5.63. Winter flounder length frequencies, April-May, 1 cm intervals, 1984-2014.....	77
Table 5.64. Winter flounder length frequencies, fall, 1 cm intervals, 1984-2014	78
Table 5.65. Winter skate length frequencies, spring and fall, 2 cm intervals (midpoint given), 1995-2014.....	79

LIST OF FIGURES

Figure 5.1. Trawl Survey site grid	82
Figure 5.2. April 2014 sites selected and sampled.....	83
Figure 5.3. May 2014 sites selected and sampled.....	84
Figure 5.4. June 2014 sites selected and sampled.....	85
Figure 5.5. September 2014 sites selected and sampled.....	86
Figure 5.6. October 2014 sites selected and sampled.....	87
Figure 5.7. The number of finfish species observed annually, 1984-2014.....	88
Figure 5.8. Plots of abundance indices for: black sea bass, bluefish (total, age 0 and ages 1+), butterfish, cunner, and dogfish (smooth and spiny).....	89
Figure 5.9. Plots of abundance indices for: flounders (fourspot, summer, windowpane, winter and winter ages 4+) and hakes (red, silver and spotted).....	90
Figure 5.10. Plots of abundance indices for: herrings (alewife, Atlantic, blueback), hogchoker, Northern kingfish, Atlantic menhaden, moonfish, and ocean pout.....	91
Figure 5.11. Plots of abundance indices for: fourbeard rockling, rough scad, longhorn sculpin, sea raven, and scup (all ages, age 0, and ages 2+).....	92
Figure 5.12. Plots of abundance indices for: searobins (striped and northern), shad (American and hickory), skates (clearnose, little, and winter), and spot.....	93
Figure 5.13. Plots of abundance indices for: striped bass, Atlantic sturgeon, tautog, and weakfish (all ages, age 0 and ages 1+).....	94
Figure 5.14. Plots of abundance and biomass indices for: crabs (lady, rock and spider), horseshoe crab, American lobster, and long-finned squid.....	95
Figure 5.15. Mean number of finfish species per sample, spring and fall, 1984-2014.....	96
Figure 5.16. Open water forage abundance, 1992-2014.....	96
Figure 5.17. Geometric mean biomass of finfish and invertebrates per sample, spring and fall, 1992- 2014.....	97
Figure 5.18. Percent of sampled winter flounder that were sexually mature by length group for female and male flounder captured in LISTS over five time periods, 1990-2013.....	98
Figure 5.19. Trends in the number of cold temperate versus warm temperate species per sample captured in spring and fall LIS Trawl Surveys.....	99

JOB 5: LONG ISLAND SOUND TRAWL SURVEY (LISTS)

CRUISE RESULTS FROM THE 2014 SPRING AND FALL SURVEYS

STUDY PERIOD AND AREA

The Connecticut DEEP Marine Fisheries Division completed the thirty-first year of the Long Island Sound Trawl Survey in 2014. The Long Island Sound Trawl Survey encompasses an area from New London to Greenwich, Connecticut and includes waters from 5 to 46 meters in depth in both Connecticut and New York state waters. Typically, Long Island Sound is surveyed in the spring, from April through June, and during the fall, from September through October. This report includes results from the 2014 spring and fall sampling periods and provides time series information since the commencement of the survey in 1984.

GOAL

To provide long term monitoring of abundance, biomass and size composition of marine fishery resources along with environmental parameters, in order to evaluate the effects of fishing and environmental conditions on the distribution and abundance of living resources in Long Island Sound.

OBJECTIVES

Provide:

- 1) Annual indices of counts and biomass per standard tow for 40 common species and age-specific indices of abundance for winter flounder, tautog, scup, summer flounder, bluefish (Age 0, 1+) and weakfish (Age 0, 1+).
- 2) Length-frequency distributions of bluefish, scup, summer flounder, winter flounder, tautog, striped bass, weakfish, black sea bass, and other ecologically important species.
- 3) Annual total counts and biomass for all finfish species taken and annual total biomass for all common macro-invertebrate species taken.
- 4) Species list for Long Island Sound based on LIS Trawl Survey sampling, noting the presence of additional species from other sampling conducted by the Marine Fisheries Division.
- 5) Fishery independent survey data to cooperative state researchers or agencies, such as the National Marine Fisheries Service, Atlantic States Marine Fisheries Commission, New England and Mid-Atlantic Councils, and researchers associated with state or local universities

INTRODUCTION

The Long Island Sound Trawl Survey (LISTS) was initiated in 1984 to provide fishery independent monitoring of important recreational species in Long Island Sound. A stratified-random design based on bottom type and depth interval was chosen and forty sites were sampled monthly from April through November to establish seasonal patterns of abundance and distribution. Seven finfish species were initially of primary interest: bluefish, scup, striped bass, summer flounder, tautog, weakfish, and winter flounder. Length data for these species were collected from every tow; scup, tautog, and winter flounder were sampled for aging. Lobster were also enumerated and measured from every tow. All fish species were identified and counted.

Since 1984, several changes have been incorporated into the Survey. In 1991, the sampling schedule was changed to a spring/fall format, although sampling is still conducted on a monthly basis (April - June, September, and October). Beginning in 1992, species were weighed in aggregate with an onboard scale to provide indices of biomass. Furthermore, more species have been sampled for lengths, such as windowpane and fourspot flounders, and important forage species such as butterfish, long-finned squid, and several herring species. By 2003, the list of species measured expanded to 20 finfish species and two invertebrate species (lobster and long-finned squid), plus rarely occurring species. In 2014, lengths were collected from all finfish species on each tow. In addition, age structures were collected from bluefish, menhaden, tautog, scup, winter flounder, weakfish and large summer flounder (>59 cm). All of these changes serve to improve the quality and quantity of information made available to fishery managers for local and regional assessment of stock condition, and to provide a more complete annual inventory of LIS (Long Island Sound) fishery resources.

METHODS

Sampling Design

LISTS is conducted from longitude 72° 03' (New London, Connecticut) to longitude 73° 39' (Greenwich, Connecticut). The sampling area includes Connecticut and New York waters from 5 to 46 m in depth and is conducted over mud, sand and transitional (mud/sand) sediment types. Sampling is divided into spring (April-June) and fall (Sept-Oct) periods, with 40 sites sampled monthly for a total of 200 sites annually. The sampling gear employed is a 14 m otter trawl with a 51 mm codend (Table 5.1). To reduce the bias associated with day-night changes in catchability of some species, sampling is conducted during daylight hours only (Sissenwine and Bowman 1978).

LISTS employs a stratified-random sampling design. The sampling area is divided into 1.85 x 3.7 km (1 x 2 nautical miles) sites (Figure 5.1), with each site assigned to one of 12 strata defined by depth interval (0 - 9.0 m, 9.1 - 18.2 m, 18.3 - 27.3 m or, 27.4+ m) and bottom type (mud, sand, or transitional as defined by Reid et al. 1979). For each monthly sampling cruise, sites are selected randomly from within each stratum. The number of sites sampled in each stratum was determined by dividing the total stratum area by 68 km² (20 square nautical miles), with a minimum of two sites sampled per stratum (Table 5.2). Discrete stratum areas smaller than a sample site are not sampled.

Sampling Procedures

Prior to each tow, temperature (°C) and salinity (ppt) are measured at 1 m below the surface and 0.5 m above the bottom using a YSI model 30 S-C-T meter. Water is collected at depth with a five-liter Niskin bottle, and temperature and salinity are measured within the bottle immediately upon retrieval.

The survey's otter trawl is towed from the 15.2 m aluminum R/V John Dempsey for 30 minutes at approximately 3.5 knots, depending on the tide. At completion of the tow, the catch is placed onto a sorting table and sorted by species. Finfish, lobsters and squid are counted and weighed in aggregate (to the nearest 0.1 kg) by species with a precision marine-grade scale (30 kg, +/- 10 gm capacity). Catches weighing less than 0.1 kg are recorded as 0.1 kg. During the initial two years of the survey (1984 & 1985), lobsters were the only invertebrates recorded. Squid abundance has been recorded since 1986. Since 1992, additional invertebrate species have been weighed in aggregate, and some have been counted. The complete time series of species counted and weighed in the survey is documented in Appendix 5.4.

For finfish species, lengths are recorded to the centimeter as either total length or fork length (e.g. measurements from 100 mm to 109 mm are recorded as 10 cm) and entered in the database as 105 mm (Table 5.3). Lobsters are measured to 0.1 mm carapace length. Squid are measured using the mantle length (cm), horseshoe crab measurements are taken using prosomal width (cm) and whelk (knobbed and channeled) shell widths are measured in millimeters.

The number of individuals measured from each tow varies by species, and also depends on the size of the catch and range of lengths (Table 5.3). If a species is subsampled, the length frequency of the catch is determined by multiplying the proportion of measured individuals in each centimeter interval by the total number of individuals caught. Some species are sorted and subsampled by length group so that, for example, all large individuals are measured and a subsample of small (often young-of-year) specimens is measured. All individuals not measured in a length group are counted. The length frequency of each group is estimated as described above, i.e. the proportion of individuals in each centimeter interval of the subsample is expanded to determine the total number of individuals caught in the length group. The estimated length frequencies of each size group are then appended to complete the length frequency for that species. This procedure is often used with catches of bluefish, scup, and weakfish, which are usually dominated by young-of-year or discrete age/length classes.

Bluefish, menhaden, scup, summer flounder, tautog, weakfish (ageing was discontinued in 2013) and winter flounder are sampled for age determination (Table 5.3). The target number of age samples (otolith) for bluefish were 50 from the spring period (defined by ASMFC Bluefish Technical Committee as Jan-July) and 50 from the fall period (August-December). Subsamples of scup, stratified by length group, are measured to the nearest mm (fork length) and scales from each individual are taken for ageing. Scup scales are removed posterior to the pectoral fin and ventral to the lateral line. The scales are pressed onto plastic laminate with an Ann Arbor roller press to obtain an impression of the scale, which is then viewed with a microfiche reader at 21x. Scales are also taken from all summer flounder greater than 59 cm. At least 15 scales are removed from the caudal peduncle area. These scales are pressed and aged to supplement the National Marine Fisheries Service age key and are also included in the formulation of LISTS summer flounder catch-at-age matrix (see below).

Menhaden scales are collected from roughly 50 fish each year as required by Amendment 2 of the ASMFC Atlantic menhaden management plan. Amendment 2 introduced a requirement for biological sampling of the commercial bait harvest to support improved stock assessments. However since Connecticut has such a small menhaden commercial fishery, sampling it would be difficult. The same size/age component of the menhaden population taken in the commercial fishery is available to LISTS so collections are taken as part of each survey cruise. Menhaden fork length (mm), and sex are recorded and scales are taken about mid-body (lateral line) and below the insertion of the dorsal fin. Most tautog taken in LISTS are aged due to the low numbers caught in recent years (under 250 fish). Tautog are iced and taken to the lab, where their total length (mm), sex, and total weight (gm) are recorded and their age is determined from opercular bones (Cooper 1967). At the request of the ASMFC Tautog Technical Committee, LISTS began collecting tautog otoliths in addition to opercles in 2012. Results from a recent ASMFC Tautog Ageing Workshop (May 2012) indicated there was no clear benefit to switching from opercles to otoliths for CT, so tautog otoliths will be collected (minimum of 50 per/ASMFC) and archived for potential use in the future. Subsamples of winter flounder, stratified by length group and area (as listed in bottom of Table 5.3), are iced and taken to the lab where they are measured to the millimeter (total length), weighed (gm) and sexed. Their maturity stage is determined (NMFS 1989), and they are aged with whole and/or sectioned otoliths (Simpson et al. 1988). Weakfish scales were obtained and processed as described above for scup, and, prior to 2013, otoliths were sectioned and read using procedures described in Simpson et al. 1988. However, since the compliance criteria for a number of other species managed through ASMFC Fishery Management Plans have increased to include ageing over the years, coincidence with stagnant (or declining) levels of funding and staff, age structures for weakfish will no longer be collected because it is not required.

In reports prior to 2001, three species were not included in annual and seasonal totals: American sand lance, bay anchovy, and striped anchovy. These species, with the possible exception of striped anchovy, can be very abundant in Long Island Sound, but are not retained well in the otter trawl. Additionally, many of these fish are young-of-year and often drop out of the net as it is retrieved and wound on the net reel. For this reason they were not included in the list of species to be counted when LISTS was started in 1984. However, to document the occurrence of these species in LISTS catches, American sand lance was added in 1994, striped anchovy was added in 1996, and bay anchovy was added in 1998. Since 2001, adults of these three species have been included in the annual and seasonal totals and the young-of-year are listed if present in the year's catch but are not quantified (Table 5.15, Appendix 5.4). Young-of-year for these three species are included in the database but are cataloged with a separate species identifier and quantities are considered estimates (Appendix 5.2).

In 2014, the only endangered species encountered by LIS Trawl Survey was Atlantic sturgeon, a species that was listed as Endangered by NOAA in 2012. Sampling procedures have been modified in recent years to minimize the likelihood of injury to Atlantic sturgeon. When sampling in a season and area where the chance of catching a sturgeon is high (based on historic LISTS catch) and water depth is greater than 27m, gear retrieval speed is reduced to decrease the stress induced by rapid changes in pressure. When a sturgeon is detected in the net, it is removed as quickly and carefully as possible. Subsequent handling and processing follow protocols described in A Protocol for Use of Shortnose, Atlantic, Gulf, and Green Sturgeons (Kahn and Mohead. 2010. U.S. Dep. Commerce, NOAA Tech Memo, NMFS-OPR-45, 62p.,

http://www.nmfs.noaa.gov/pr/pdfs/species/kahn_mohead_2010.pdf) and adhere to the Reasonable and Prudent Measures, as well as, the Terms and Conditions spelled out in the ESA Section 7 Biological Opinion's Incidental Take Statement issued by NOAA for CT in June 2012 (http://www.greateratlantic.fisheries.noaa.gov/protected/section7/bo/actbiops/usfws_state_fisheries_surveys_2013.pdf). Future LISTS interactions with sturgeon will follow requirements of the subsequent biological opinion issued by NOAA for the 11 Northeast States and District of Columbia. All interactions with endangered species are detailed in Appendix 5.5.

Data Analysis

Indices of Abundance: Annual Mean Count and Weight per Tow

To evaluate the relative abundance of common species, an annual spring (April - June) and fall (September - October) geometric mean number per tow and weight per tow (biomass, kg) is calculated for the common finfish and invertebrate species. To calculate the geometric mean, the numbers and weight per tow are logged (\log_e) to normalize the highly skewed catch frequencies typical of trawl surveys:

$$\text{Transformed variable} = \ln(\text{variable}+1).$$

Means are computed on the log scale and then retransformed to the geometric mean:

$$\text{geometric mean} = \exp(\text{mean})-1.$$

The geometric mean count per tow was calculated from 1984 - 2014 for 38 finfish species, lobster, and long-finned squid (1986 - 2014). The geometric mean weight per tow was calculated using weight data collected since 1992 for the same species, plus an additional 13 invertebrates.

For the seven finfish species that were measured on every tow (bluefish, scup, striped bass, summer flounder, tautog, weakfish, and winter flounder) biomass indices were calculated for the years 1984 - 1991 by using length/weight equations to convert length frequencies to weight per tow. Bluefish, scup, weakfish and winter flounder lengths were converted using equations from Wilk et al. 1978; striped bass conversions were accomplished using an equation from Young et al. 1994; summer flounder and tautog conversions were accomplished using equations developed from LISTS data from 1984 -1987 and 1984 -1996 respectively.

Indices of Abundance: Indices-at-Age and Age Group

Annual age specific indices (indices-at-age matrices) were calculated for scup, striped bass, summer flounder, winter flounder and tautog. The age data used to calculate the indices came from three sources: striped bass ages were derived using the von Bertalanffy (1938) equation; summer flounder age-length keys were obtained from the National Marine Fisheries Service (NMFS) Northeast Fisheries Science Center spring and fall trawl surveys combined with LISTS ages (>59 cm); scup, winter flounder and tautog age-length keys (in 1 cm intervals) were obtained directly from LISTS. Since fish growth can fluctuate annually as a function of population size or other environmental factors, a year and season specific age-length key was used wherever possible. Once lengths have been converted to age, the proportion at age is multiplied by the abundance index of the appropriate season to produce an index of abundance at age.

Recruitment (young-of-year) and age 1+ (all fish age one and older) indices were calculated for bluefish and weakfish. Observed modes in the length frequencies were used to separate the two groups.

The specific methods used to calculate indices-at-age for each species were as follows:

- ◆ **Bluefish.** Otoliths were taken from 190 bluefish, 22 from the spring period and 168 from the fall period. Of the 22 samples taken in the spring, only 6 were obtained from LISTS; the bulk of the samples came from recreational anglers. All of the fall samples were obtained from LISTS (168 fish). No samples were obtained through donations from a fishing tournaments in 2014. In 2012 a coast wide biological sampling program was initiated through ASMFC addendum 1 of the bluefish management plan. Since there is only three years of data from the northeast, there are still limited results available at this time. Therefore, the method of using modes observed in the fall length frequencies to separate bluefish into age 0 and age 1+ groups, and calculate a geometric mean catch per tow for each group (Table 5.22) was continued through 2014. Comparison of the mean length-at-ages reported for young-of-year and age 1 bluefish in the New York Bight (Chiarella and Conover 1990) and Long Island Sound (Richards 1976) with LISTS length frequencies suggests that bluefish can easily be identified as either age 0 (snapper bluefish) or adults (age 1+). Richards (1976) and Chiarella and Conover (1990) determined that most bluefish less than 30 cm are age 0. A discontinuity in the LISTS fall length frequencies occurs most years between 26 cm and 39 cm (Table 5.42). Therefore 30 cm was determined to be a suitable length for partitioning age 0 and age one fish. With the addition the biological sampling programs along the coast, a regional northeast key is being finalized through ASMFC and from 2013 forward, a LISTS key will be utilized for calculating a full index-at-age for Long Island Sound.

Prior to 2012, there was limited bluefish ageing in the northeast. Although North Carolina state biologists have aged bluefish for some time, their age keys were not used to age Long Island Sound bluefish because North Carolina mean lengths-at-age are not consistent with modes observed in Long Island Sound bluefish length frequencies. This difference suggests that growth may vary by region, or that early and late spawned bluefish may be differentially distributed along the coast (Kendall and Walford 1979).

- ◆ **Scup.** An index-at-age matrix was developed for 1984-2014 using spring (May-June only) and fall (September-October) LISTS data (Table 5.23). April data was omitted since very few scup are taken at this time. A total of 12,910 scup aged between 1984 and 2014 were used to make year and season specific age-length keys (1 cm intervals). In the relatively few instances when the season/year specific key failed at a given 1 cm length interval, a three-year pooled key was used to determine the age. Three-year pooled keys were calculated using the years preceding and following the “run” year. For the terminal year, only two years were used for the pooled key. The final index-at-age was computed for both spring and fall indices-at-age. Since very few scup older than age 9 are taken (less than 4% in any given year), an age 10+ group is calculated by summing indices for ages 10 and up. To represent the full adult portion of the population an age 2+ index is calculated by summing the indices for ages 2 through 10+.

- ◆ **Striped bass.** To approximate the ages of striped bass taken in the spring survey (Table 5.24), the average of the Chesapeake Bay and Hudson River striped bass von Bertalanffy parameters ($L_{\max} = 49.9$ in, $K = 0.13$, $t_0 = 0.16$, Vic Crecco, pers. comm.) were used in the rearranged von Bertalanffy equation:

$$t = (1/K) * (-\log_e ((L_{\max} - L_t) / L_{\max})) + t_0$$

Since this equation estimates age t as a fraction of a year, the estimates were rounded to the nearest year (e.g. age 3 = ages 2.5 to 3.4). A spring catch-at-age matrix was developed for 1984 through 2014 by apportioning the spring index by the percentage of fish at each age (Table 5.25).

- ◆ **Summer flounder.** The year and season specific age-length keys (1 cm intervals) used to age LISTS catches were provided by NMFS from their spring and fall trawl surveys. These keys were supplemented with fish caught and aged by LISTS (typically 60 cm and over). In 2014 LISTS had sample requests for summer flounder and scale samples from these fish (< 60cm) were collected. In 2014, 19 summer flounder, were aged; 14 from the spring and 5 (all > 60cm) from the fall. Since 2001, whenever the season/year specific key failed at a given 1 cm length interval a pooled year key using only adjacent years was used (Gottschall and Pacileo 2002). Since it is thought that growth rates for summer flounder have changed over time, a pooled key using only adjacent years would more accurately represent fish that could not be aged by the season/year specific key. Using this methodology, the catch-at-age matrix (Table 5.26) will remain unchanged for all but the terminal year, which will be updated as the following years' data becomes available.
- ◆ **Tautog.** An index-at-age matrix was developed for 1984-2014 using all survey months (Gottschall and Pacileo 2007) (Table 5.27). During 2014, 167 tautog were captured and opercles were collected from all; 131 collected in the spring and 36 were collected in the fall. Ageing for 2006-2012 has been completed. Ageing for 2013-2014 samples has not yet been completed. A 2012 age key was used for the 2013-2014 un-aged fish and a pooled key was used where the 2012 key failed. Therefore, the 2013-2014 indices-at age are preliminary; the 2013-2014 tautog samples will be aged during the summer of 2015 and an updated index-at-age matrix will be constructed.
- ◆ **Weakfish.** Age 0 and age 1+ indices were calculated for both spring (1984 – 2013) and fall surveys (1984 – 2009, 2013) (Table 5.28). Since few weakfish are taken in April, the spring geometric mean was calculated using only May and June. All weakfish taken in spring are assumed to be age 1+. Similar to bluefish, the fall age 0 and 1+ indices were calculated by using length frequencies to separate the catch. Since a break in the fall length frequencies generally occurs between 24 and 32 cm each year (Table 5.57), weakfish less than 30 cm are considered to be age 0 while those greater than or equal to 30 cm are ages 1+. Ageing for weakfish was discontinued in 2013.
- ◆ **Winter flounder.** An index-at-age matrix was developed for 1984-2014 using April and May LISTS data (Table 5.29). June data were not used since length frequency data suggest that many adult winter flounder have left the Sound by this time (an exception

was made for 1984, the first year of LISTS, because very few samples were taken in the spring months). A total of 23,272 winter flounder aged between 1984 and 2014 were used to make year and region (east of Stratford Shoal, west of Stratford Shoal) specific age-length keys in 1 cm intervals. Similar to scup and summer flounder, three year pooled keys using only the adjacent years (two years for the terminal year runs) were used to assign ages if year specific keys were not available.

Each flounder aged as described above was also assessed for maturity stage (following Burnett 1989) by sex. CT DEEP staging of winter flounder was verified in a cooperative study with NMFS in 2009-2010 (Gottschall and Pacileo 2011). The percentage of male and female fish in each centimeter length group that was sexually mature (ripe, resting, or spent) was calculated in order to determine the length group at which 50% was mature each year.

Species Richness by Group

The Long Island Sound Trawl Survey monitors species richness using groups of species classified as either cold temperate or warm temperate. For the purposes of tracking species richness, American sand lance, bay anchovy, and striped anchovy were omitted (see *Sampling Procedures* section). All other finfish species captured in LISTS were divided into groups based on their temperature preferences and seasonal spawning habits as documented in the literature (Collette and Klein-MacPhee 2002, Murdy et al. 1997). Species in the cold temperate group prefer water temperatures below 15⁰C (60⁰F), tend to spawn at the lower end of their temperature tolerance range, and are more abundance north of Long Island Sound than south of New York. Species in the warm temperate group prefer warmer temperatures (11-22⁰C or 50-77⁰F), tend to spawn in the upper range of their temperature tolerance, and are more abundant south of the Sound than north of Cape Cod (Appendix 5.6). Species that are not tolerant of cold temperatures, are abundant only south of Chesapeake Bay but stray into northern waters mostly as juveniles, and spawn only in the mid-Atlantic Bight and south were placed into a separate group (subtropical) and were not included in the analysis because they are typically only present in the fall LISTS.

Open Water Forage Abundance

A Long Island Sound open water forage index of abundance was compiled to measure the available food base which supports resident and migratory species within the Sound. This index is formulated as a biomass index that is assembled from 11 of the forage species that are most common in LISTS catches along with three other species that are considered forage at an early life stage (young-of-year, YOY). The species used to generate the index are; Atlantic herring, long-finned squid, butterfish, alewife, blueback herring, American shad, hickory shad, menhaden, whiting, spotted hake, and red hake along with young-of-year stage of scup, bluefish, and weakfish. The geometric mean biomass is calculated using the aggregate of these 14 species on a per tow basis and calculated using the same methodology as described above for individual species biomass indices.

RESULTS AND DISCUSSION

Overview of LISTS 2014 Spring and Fall Surveys

Each month of the survey, sampling aboard the R/V John Dempsey generally began in the east end of Long Island Sound and progressed westward. The April survey commenced on April 9, 2014, and continued until May 2 for a total of six (6) days underway and 40 tows completed. May sampling started on May 12 and continued till May 22 with eight (8) sampling days underway and 40 sites completed. June sampling began on June 9 and ended on June 24, taking ten (10) days underway to complete the 40 sites. The Fall Survey commenced on September 9 and needed 8 days underway to complete 40 tows. The October survey fell short by 1 tow (39 completed instead of 40 tows) because of weather and took 11 sampling days. The October cruise continued until November fourth. A total of 199 LISTS tows were completed in 43 days underway during the spring and fall 2014 surveys (Table 5.4); not including transit days or weather days.

Maps showing the sites selected versus the sites sampled during each month of sampling are provided in Figure 5.2 (April), Figure 5.3 (May), Figure 5.4 (June), Figure 5.5 (September) and Figure 5.6 (October). Within each figure the red bordered sites are the sites selected for the month and the solid blue dots indicate the actual sites sampled. If a site had to be relocated during sampling, an explanation of why it was moved is provided under the figure. Additional site/station information is provided in Table 5.5 (April), Table 5.6 (May), Table 5.7 (June), Table 5.8 (September) and Table 5.9 (October). These tables provide date of sample, time, tow duration, latitude/longitude, surface and bottom temperature and salinity, average tow speed, distance towed and approximate area swept for each tow.

Sometimes, a full 30-minute tow cannot be completed. Typical reasons for short tows include lack of room because of observed pot gear set in the immediate area, a drop in speed due to entanglement with some object on the bottom (frequently derelict pot gear), or a complete stop in forward motion (submerged wreck or rock pile). Survey crew will often attempt to finish an interrupted tow by clearing the net (if needed) and resetting beyond the obstruction or observed gear. If this is not possible, a site may have to be moved to another site nearby with the same stratum (bottom type and depth). If the site was moved, the data from the initial site will not be used. Typically, a minimum of 15-20 minutes is required for a LISTS tow to be recorded. However, there are occasions when a tow with less than 15 minutes will be accepted, usually because there is no alternate site in the designated strata in the vicinity. Short tow information for each month in the 2014 survey is summarized in Table 5.10.

Cooperative Sample and Data Collection

Throughout the time series, LISTS staff have been participating in cooperative efforts for sample collections, data requests, and special projects using survey personnel, equipment, and other resources. Most of these cooperative efforts are with state researchers or agencies, the National Marine Fisheries Service, Atlantic States Marine Fisheries Commission, New England and Mid-Atlantic Councils, and researchers or graduate students associated with state or local universities. Table 5.11 illustrates many of the organizations that requested data in 2014, while Table 5.12 shows sample request received and fulfilled. In recent years many requests for samples have come from high schools, aquariums, or other educational organizations needing

finfish and invertebrates for teaching purposes. Additionally, our own staff often have sample or data requests for media or other public outreach events (see Job 11 of this report).

Number of Species Identified

Sixty finfish species were observed in the 2014 Long Island Sound Trawl Survey (Table 5.13). This includes one new species for the survey; blue spotted cornet fish (*Fistularia tabacaria*). From 1984 to 2014, LIS Trawl Survey has identified one hundred six (106) finfish species (Appendix 5.1), averaging 58 species per year with a range of 43 to 70 species (Figure 5.7). In addition, a total of 42 types of invertebrates were collected in 2014 (Table 5.14). Most invertebrates are identified to species. However, in some cases, invertebrates were identified to genus or a higher level taxon.

Total Catch

Appendix 5.4 presents a time series (1984-2014) of the finfish species collected each year and their respective rank by numbers. Annual total biomass of invertebrates is also included in this appendix (1992-2014), ranked by weight (kg). A total of 153,100 finfish weighing 16,174 kg were sampled in 2014 (Table 5.15). A total of 33,919 finfish weighing 8,816 kg were sampled in spring of 2014 (Table 5.16). A total of 119,182 finfish weighing 7,358 kg were sampled in fall of 2014 (Table 5.16). A total of 1,529 kg of invertebrates were taken in 2014 (Table 5.15). The total biomass of invertebrate catch taken in the spring of 2014 was 866 kg (Table 5.17). A total of 663 kg of invertebrates were taken in fall of 2014 (Table 5.17).

Length Frequencies

Length frequency tables are provided primarily to give the reader an understanding of the size range of various species taken in LISTS. Lengths are converted to age frequencies for analysis of principal species such as scup, bluefish, striped bass, summer flounder, tautog, winter flounder, and weakfish. Changes such as an expansion in the size (age) range for some important recreational species are apparent in recent years including more large scup (Table 5.52-5.53), striped bass (Table 5.54-5.55), and summer flounder (Table 5.56-5.57).

Length frequencies were prepared for 22 species:

alewife	spring and fall	1989 - 2014	Table 5.30;
American shad	spring and fall	1989 - 2043	Table 5.31;
American lobster	spring and fall (M&F)	1984 - 2014	Table 5.32-Table 5.35;
Atlantic herring	spring and fall	1989 - 2014	Table 5.36;
Atlantic menhaden	spring and fall	1996 - 2014	Table 5.37;
black sea bass	spring and fall	1987 - 2014	Table 5.38, Table 5.39
blueback herring	spring and fall	1989 - 2014	Table 5.40;
bluefish	spring and fall	1984 - 2014	Table 5.41, Table 5.42;
butterfish	spring and fall	1986 - 1990, 1992 - 2014	Table 5.43;
clearnose skate	spring and fall	1993 - 2014	Table 5.44, Table 5.45;
fourspot flounder	spring and fall	1989 - 1990, 1996 - 2014	Table 5.46;
hickory shad	spring and fall	1991 - 2014	Table 5.47;

horseshoe crab	spring and fall (M&F)	1998 - 2014	Table 5.48, Table 5.49;
long-finned squid	spring and fall	1986 - 1990, 1992 - 2014	Table 5.50, Table 5.51;
scup	spring and fall	1984 - 2014	Table 5.52, Table 5.53;
striped bass	spring and fall	1984 - 2014	Table 5.54, Table 5.55;
summer flounder	spring and fall	1984 - 2014	Table 5.56, Table 5.57;
tautog	spring	1984 - 2014	Table 5.58;
weakfish	spring and fall	1984 - 2014	Table 5.59, Table 5.60;
windowpane flounder	spring and fall	1989, 1990, 1994 - 2014	Table 5.61, Table 5.62;
winter flounder	April-May and fall	1984 - 2014	Table 5.63, Table 5.64;
winter skate	spring and fall	1995 - 2014	Table 5.65.

For the years where length data are available, length frequencies were prepared for the seasons or months for which the preferred indices of abundance and catch-at-age matrices are calculated; for some species length frequencies are provided for both seasons.

Seasonal Indices of Abundance

The geometric mean count per tow was calculated from 1984-2014 for 38 finfish species plus lobster and long-finned squid (squid since 1986). All spring (April-June) and fall (September-October) data are used to compute the abundance indices presented in Tables 5.18 (spring) and 5.19 (fall), with the preferred seasonal index (for counts) denoted by an asterisk. Geometric mean biomass-per-tow indices have been calculated for 38 finfish and 15 invertebrate species (or species groups) since 1992, for both spring and fall (Table 5.20 and 5.21, respectively). Age specific indices of abundance were calculated for selected important recreational species, including scup, striped bass, summer flounder, and winter flounder (see below). For two other species, bluefish and weakfish recruitment indices were calculated using modal analysis of the length frequencies. For each of the thirty-eight finfish species, plots including catch per tow in numbers and biomass in kilograms are illustrated in Figures 2.8 through 5.13. These figures also include plots of each of the age specific indices and recruitment indices mentioned above. Figure 5.14 provides plots of abundance (biomass) indices for crabs (lady, rock, spider; 1992-2014), American lobster (1984-2014), horseshoe crab (1992-2014), and long-finned squid (1986-2014).

Indices of Abundance: Important Recreational Species

Spring and fall abundance indices are presented in Tables 5.18-5.19. Indices of abundance at age were also calculated for seven important recreational species: bluefish (Table 5.22), scup (Table 5.23), striped bass (Table 5.24 age frequency, Table 5.25 indices at age), summer flounder (Table 5.26), tautog (Table 5.27), weakfish (Table 5.28) and winter flounder (Table 5.29). Bluefish and striped bass indices-at-age are based on the fall and spring surveys, respectively, whereas winter flounder indices-at-age are based on only the April and May cruises of the spring survey. In 2014, LISTS collected otoliths from 602 winter flounder, 601 of which were used in the development of age keys and the final catch-at-age matrix. Both scup and weakfish indices-at-age are calculated and presented separately for each season. Scales from 702 scup were collected and aged in 2014, 689 of which were used in the keys and calculations of the age matrix. Weakfish and bluefish use modal distributions for calculating their respective recruitment index although a small number of weakfish are taken each year for ageing purposes (see methods).

Winter Flounder Average Size at Maturity

Average size at maturity for winter flounder captured in April and May cruises has increased since maturation data recording began in 1990. The number mature by cm-interval and sex was calculated for the subset of fish examined in the laboratory each year, and a five-year average computed to maximize sample size. The resulting maturation curves (Figure 5.18) skew right for both sexes from 1990-94 to 2010-2013. The 50%-midpoint for females has increased from 24-26cm in the 1990s to 27cm after 2000. The 50%-midpoint for males has increased from 16-19cm in the 1990s to 20-22cm after 2000. These results indicate not only a larger average size at maturation but also a greater synchronization of the maturation process over a smaller size range.

Species Richness by Group

The number of cold temperate and warm temperate species captured in each tow was averaged by seasonal cruise (April-June and September-October) for each year from 1984-2014 as an indicator of annual biological diversity or species richness. Trends in these indicators were tested for statistical significance by regression analysis. Results (Figure 5.19) show that the average number of warm temperate species captured/tow in spring and fall cruises has increased ($F=26.8$ and 76.0 respectively, $p<0.001$); while the average number of cold temperate species has decreased, especially in spring ($F=36.8$, $p<0.001$) but also in fall cruises ($F=11.7$, $p=0.002$).

MODIFICATIONS

No modifications.

LITERATURE CITED

- American Fisheries Society. 2004. Common and Scientific Names of Fishes from the United States, Canada, and Mexico Sixth ed. American Fisheries Society Special Publication 29, Bethesda, MD. 386 pp.
- von Bertalanffy, L. 1938. A quantitative theory of organic growth (Inquiries on growth laws. II). Hum. Biol. 10 (2): 181-213.
- Burnett, J., L. O'Brien, R.K. Mayo, J.A. Darde and M. Bohan. 1989. Finfish maturity sampling and classification schemes used during Northeast Fisheries Center bottom trawl surveys, 1963 – 89. NOAA Technical Memorandum NMFS-F/NEC-76: 14 pp. (<http://www.nefsc.noaa.gov/nefsc/publications/tm/tm76.pdf>).
- Chiarella, L.A. and D.O. Conover. 1990. Spawning season and first-year growth of adult bluefish from the New York Bight. Transactions of the American Fisheries Society 119:455-462.
- Collette, B. and G. Klein-MacPhee, 2002, editors. Bigelow and Schroeder's Fishes of the Gulf of Maine, 3rd edition. Smithsonian Institution Press, Washington DC.
- Cooper, R.A. 1967. Age and growth of the tautog, *Tautog onitis* (Linnaeus), from Rhode Island. Trans. Amer. Fish. Soc. 96: 132-134.
- Fahay, M.P., P.L. Berrien, D.L. Johnson and W.W. Morse. 1999. Essential Fish Habitat Source document: Atlantic Cod, *Gadus morhua*, Life History and habitat characteristics. NOAA Technical Memorandum NMFS-NE-124: 41 pp. (<http://www.nefsc.noaa.gov/publications/tm/tm124/tm124.pdf>).
- Flescher, D.D. 1980. Guide to some trawl-caught marine fishes from Maine to Cape Hatteras, North Carolina. NOAA Tech. Rpt. NMFS Circular 431, 34 pp.
- Gosner, K.L. 1978. A Field Guide to the Atlantic Seashore. Peterson Field Guide Series. Houghton Mifflin Company, Boston, MA. 329 pp.
- Gottschall, K.F, M.W. Johnson and D.G. Simpson. 2000. The distribution and size composition of finfish, American lobster, and long-finned squid in Long Island Sound based on the Connecticut Fisheries Division Bottom Trawl Survey, 1984-1994. U.S. Dep. Commer., NOAA Tech Rep. NMFS 148, 195p.
- Gottschall, K and D. Pacileo. 2011. Marine Finfish Survey, Job 2. In: A Study of Marine Recreational Fisheries in Connecticut. Annual Progress Report, CT DEP/Marine Fisheries Division, Old Lyme, CT. 203 pp.
- Gottschall, K and D. Pacileo. 2008. Expansion of the DEP Long Island Sound Trawl Survey, Job 2 (100 pp). In: Assessment and Monitoring of the American Lobster Resource and Fishery in Long Island Sound. State of CT, Final Project Report to NOAA NMFS Northeast Region for Grant # NA16FW1238, 474 pp.
- Gottschall, K and D. Pacileo. 2007. Marine Finfish Survey, Job 2. In: A Study of Marine Recreational Fisheries in Connecticut. Annual Progress Report, CT DEP/Fisheries Division, Old Lyme, CT. 203 pp.
- Gottschall, K and D. Pacileo. 2002. Marine Finfish Survey, Job 2. In: A Study of Marine

- Recreational Fisheries in Connecticut. Annual Progress Report, CT DEP/Fisheries Division, Old Lyme, CT. 176 pp.
- Johnson, M and D. Shake. 2000. Marine Finfish Survey, Job 2. In: A Study of Marine Recreational Fisheries in Connecticut. Annual Progress Report, CT DEP/Fisheries Division, Old Lyme, CT. 160 pp.
- Kahn, Jason, and Malcolm Mohead. 2010. A Protocol for Use of Shortnose, Atlantic, Gulf, and Green Sturgeons. U.S. Dep. Commerce, NOAA Tech Memo, NMFS-OPR-45, 62p.
- Kendall, A.W., Jr., and L.A. Walford. 1979. Sources and distribution of bluefish, *Pomatomus saltatrix*, larvae and juveniles off the east coast of the United States. U.S. Fish and Wildlife Service Fishery Bulletin 77:213-227.
- Murdy, E., R. Birdsong and J. Musick, 1997, editors. Fishes of Chesapeake Bay. Smithsonian Institution Press, Washington DC.
- O'Brien, L., J. Burnett and R. Mayo. 1993. Maturation of Nineteen Species of Finfish off Northeast Coast of the United States, 1985-1990. NOAA Technical Report NMFS 113. 66 pp.
- Reid, R.N., A.B. Frame and A.F. Draxler. 1979. Environmental baselines in Long Island Sound, 1972-73. NOAA Tech. Rpt. NMFS SSRF-738, 31 pp.
- Richards, S. W. 1976. Age, growth and food of the bluefish (*Pomatomus saltatrix*) from east-central Long Island Sound from July through November 1975. Transactions of the American Fisheries Society 105:523-525.
- Simpson, D.G., P.H. Howell and M. Johnson. 1988. Marine Finfish Survey, Job 2. In: A Study of Marine Recreational Fisheries in Connecticut. Final report, Ct DEP/Fisheries Division, Old Lyme, Ct. 265 pp.
- Simpson, D.G., K Gottschall and M Johnson. 1991. Marine Finfish Survey, Job 2. In: A Study of Marine Recreational Fisheries in Connecticut. Annual performance report, Ct DEP/Fisheries Division, Old Lyme, Ct. 80 pp.
- Sissenwine, M.P. and L. Bowman. 1978. Factors affecting the catchability of fish by bottom trawls. ICNAF Research Bulletin No.13: 81-87.
- Wilk, S.J., W.W. Morse and D.E.Ralph. 1978. Length-weight relationships of fishes collected in the New York Bight. Bull. New Jersey Acad. Sci. Vol 23, No 2, pp58-64.
- Young, B.H., K.A. McKnown and P.S. Savona. 1994. A study of the striped bass in the marine district for New York, VII. Completion Rept., N.Y. DEC. 133pp.

TABLES 5.1 - 5.29
LISTS

Table 5.1. Specifications for the Wilcox 14 m high-rise trawl net and associated gear.

Component	Description
Headrope	9.1 m long, 13 mm combination wire rope
Footrope	14.0 m long, 13 mm combination wire rope
Sweep	Combination type, 9.5 mm chain in belly, 7.9 mm chain in wing
Floats	7 floats, plastic, 203 mm diameter
Wings	102 mm mesh, #21 twisted nylon
Belly	102 mm mesh, #21 twisted nylon
Tail Piece	76 mm mesh, #21 twisted nylon
Codend	51 mm mesh, #54 braided nylon
Ground Wires	18.2 m long, 6x7 wire, 9.5 mm diameter
Bridle Wires:	top legs 27.4 m long, 6x7 wire, 6.4 mm diameter
Bottom Legs	27.4 m long, 6x7 wire, 11.1 mm, rubber disc type, 40 mm diameter
Doors	Steel "V" type, 1.2 m long x 0.8 m high, 91 kg
Tow Warp	6x7 wire, 9.5 mm diameter

Table 5.2. The number of sites scheduled for sampling each month within the 12 depth-bottom type strata.

Bottom type	Depth Interval (m)				Totals
	0 - 9.0	9.1 - 18.2	18.3 - 27.3	27.4+	
Mud	2	3	5	5	15
Sand	2	2	2	2	8
Transitional	3	5	5	4	17
Totals	7	10	12	11	40

Table 5.3. Length and age data collected in 2014.

In addition to the species listed below, other rarely occurring species (totaling less than 30 fish/year each) were measured. During 2014, thirty-one other species were measured during LISTS sampling as either rarely occurring species or for other research related projects

Species measured	Measurement	# tows/day	# fish measured
Alewife	FL (cm)	All	min of 15 / tow
American lobster	CL (0.1 mm)	All	min of 50 / tow
American shad	FL (cm)	All	min of 15 / tow
Atlantic herring	FL (cm)	All	min of 15 YOY and min of 30 adults / tow
Atlantic menhaden	FL (cm)	All	min of 15 / tow
Atlantic sturgeon	FL (cm)	All	All
Blueback herring	FL (cm)	All	min of 15 / tow
Bluefish	FL (cm)	All	min of 30 YOY / tow, all adults
black sea bass	TL (cm)	All	All
butterfish	FL (cm)	All	min of 15 YOY and 15 adults / tow
cunner	TL (cm)	All	All
dogfish, smooth	FL (cm)	All	All
dogfish, spiny	FL (cm)	All	All
fourspot flounder	TL (cm)	All	min of 30/tow
hake, red	TL (cm)	All	min of 30/tow
hake, silver (whiting)	TL (cm)	All	min of 30/tow
hake, spotted	TL (cm)	All	min of 30/tow
hickory shad	FL (cm)	All	All
horseshoe crab	PW (cm)	All	All
northern searobin	FL (cm)	All	min of 30/tow
moonfish	FL (cm)	All	min of 10/tow
smallmouth flounder	TL (cm)	All	min of 10/tow
striped bass	FL (cm)	All	All
striped searobin	FL (cm)	All	min of 30/tow
scup	FL (cm)	All	min of 15 YOY and 30 / mode for age 1+
long-finned squid	ML (cm)	All	min of 30 / tow
summer flounder	FL (cm)	All	All
tautog	TL (cm)	All	All
weakfish	FL (cm)	All	min of 15 YOY / tow, all adults
whelk , channeled	PW (mm)	All	All
whelk , knobbed	PW (mm)	All	All
windowpane flounder	TL (cm)	All	min of 50 / tow
winter flounder	TL (cm)	All	min of 100 / tow
winter skate	TL (cm)	All	All

Species aged	Structure	Subsample
bluefish	scales / otoliths	Collected each season. For each season, minimum of 50 scale and otolith samples collected from full length distribution. Spring collection may use other means of sampling to obtain the required minimum.
Menhaden	scales	Collected each season. For each season, minimum of 50 scale samples collected from full length distribution.
scup	scales	Collected every month. For each month scales are taken from the following: 3 fish/cm <20 cm; 5/cm from 20-29 cm; and all fish > 30 cm.
summer flounder	scales	all fish > = 60 cm
tautog	opercular bones	Collected from a minimum of 200 fish/year.
weakfish	scales / otoliths	Ageing/collections discontinued in October 2014
winter flounder	otoliths	Collected during April and May from two areas in the Sound: eastern-central and western. For each month and area, subsamples are taken as follows: in the eastern-central area 7 fish / cm < 30 cm, 14 / cm from 30-36 cm, all fish > 36 cm. In the western area 5 fish / cm < 30 cm, 10/cm from 30-36 cm, all fish > than 36 cm.

Notes: min = minimum; YOY = young-of-year; FL = fork length; TL = total length; CL = carapace length; ML = mantle length; PW = prosomal width.

Table 5.4. Number of Long Island Sound Trawl Survey (LISTS) samples taken by year and cruise.

In 1984, thirty-five sites per monthly cruise from April through November were scheduled for sampling. Starting in 1985, forty sites per cruise were scheduled. In 1991, the Trawl Survey was modified to a spring (April - June) and fall (September - October) format--July, August and November sampling was suspended. In 1993 and 1994, an additional cruise of 40 sites was added to the fall period. The additional fall cruise was suspended in 1995. One hundred twenty tows were conducted in 2006 due to delays in rebuilding the main engine on the R/V John Dempsey (spring) and mechanical failure/overhaul of the hydraulic power take-off (fall). Delays in overhauling the transmission in the fall of 2008 resulted in missing September sampling. The June cruise and all of fall sampling in 2010 were canceled for an engine replacement in the R/V John Dempsey. Due to delays in engine replacement, begun in 2010 but not completed until late April 2011, April sampling in 2011 was abbreviated.

Cruise	Year																													Total		
	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012		2013	2014
April	-	-	35	40	40	40	40	45*	-	40	40	40	40	40	40	40	40	40	40	40	40	40	-	40	40	40	40	12	40	40	40	
May	13	41	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	38	40	40	40	40	
June	19	5	41	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	39	40	40	40	40	40	-	40	40	40	40	
July	35	40	40	40	40	40	17	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
August	34	40	40	40	40	40	40	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
September	35	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	41**	40	40	40	40	40	40	40	40	-	40	-	40	40	40	
Sept/Oct	-	-	-	-	-	-	-	-	-	40	40	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
October	35	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	-	40	40	-	40	40	40	-	40	40	40	39	
November	29	40	40	40	40	40	40	-	-	-	-	-	-	-	-	-	-	-	-	40	-	-	-	-	-	-	-	-	-	-	-	
Total	200	246	316	320	320	320	297	205	160	240	240	200	200	200	200	201	200	200	200	200	199	200	120	200	160	200	78	172	200	200	199	6,593

Table 5.5. Station information for LISTS April 2014.

Standard LISTS tows in the spring begin with SP and fall begins with FA. Latitude (N) and Longitude (W) are displayed in decimal degrees. Surface and bottom temperature and salinity are labeled as S_ and B_, respectively. Area swept is estimated by assuming the effective sweep is 2/3rds of the footrope length.

Sample Number	Date	Site Number	Bottom Type	Depth Interval	Time Start	Duration (min)	Latitude	Longitude	S_Temp (sfc, C)	S_Salinity (sfc, ppt)	B_Temp (btm, C)	B_Salinity (btm, ppt)	Ave Speed (knots)	Distance (nm)	Area Swept (sqnm)
SP2014001	4/9/2014	1433	S	2	10:05	30	41.2465	-72.3514	4.1	26.3	4.3	27.4	1.4	0.6854	0.0035
SP2014002	4/9/2014	1737	T	1	12:06	30	41.2892	-72.1974	5.3	27.6	4.4	29.8	3.1	1.5507	0.0078
SP2014003	4/11/2014	1432	S	2	7:18	30	41.2336	-72.3997	4.4	21.8	4.3	27.8	3.6	1.7784	0.0090
SP2014004	4/11/2014	1336	T	4	9:24	30	41.2237	-72.2355	4.6	27.3	4.2	30.8	2.6	1.3099	0.0066
SP2014005	4/11/2014	831	S	4	11:58	30	41.1433	-72.4478	4.4	28.0	3.9	29.3	2.2	1.1114	0.0056
SP2014006	4/11/2014	929	S	3	13:09	30	41.1648	-72.5309	4.3	27.8	4.0	28.4	2.3	1.1378	0.0058
SP2014007	4/11/2014	1029	S	3	14:11	30	41.1648	-72.5875	4.6	27.9	3.9	28.1	2.9	1.4291	0.0072
SP2014008	4/21/2014	931	S	4	7:50	30	41.1593	-72.4492	5.8	27.4	5.6	27.9	1.9	0.9279	0.0047
SP2014009	4/21/2014	128	T	2	9:43	30	41.0303	-72.5820	6.3	27.5	6.1	27.4	2.9	1.4675	0.0074
SP2014010	4/21/2014	227	T	3	11:00	30	41.0458	-72.6033	6.4	27.5	5.7	27.6	3.2	1.6163	0.0082
SP2014011	4/21/2014	325	T	3	12:21	30	41.0570	-72.7542	6.6	27.6	4.6	27.8	2.7	1.3316	0.0067
SP2014012	4/21/2014	526	T	3	13:28	30	41.0901	-72.6899	6.7	27.7	5.5	27.6	2.4	1.2152	0.0061
SP2014013	4/21/2014	725	T	4	14:54	30	41.1200	-72.7455	6.3	27.6	5.5	27.6	2.6	1.2809	0.0065
SP2014014	4/21/2014	926	T	4	15:56	30	41.1503	-72.6949	6.8	25.4	5.8	27.5	2.6	1.2911	0.0065
SP2014015	4/22/2014	1328	T	2	8:20	30	41.2380	-72.5781	6.3	20.4	6.2	26.3	2.5	1.2717	0.0064
SP2014016	4/22/2014	1428	T	1	9:16	30	41.2472	-72.5800	6.6	23.8	6.2	26.0	2.8	1.4230	0.0072
SP2014017	4/22/2014	924	T	3	10:59	30	41.1720	-72.7795	6.7	26.6	5.9	27.1	3.1	1.5670	0.0079
SP2014018	4/22/2014	523	M	4	12:10	30	41.0908	-72.7973	7.5	27.6	5.4	27.7	3.2	1.6138	0.0082
SP2014019	4/22/2014	221	M	4	13:23	30	41.0425	-72.8856	7.6	27.5	4.7	27.8	3.1	1.5472	0.0078
SP2014020	4/28/2014	313	M	3	8:41	30	41.0633	-73.1986	7.1	26.3	6.6	27.1	3.4	1.6800	0.0085
SP2014021	4/28/2014	5911	M	3	10:49	30	40.9876	-73.3303	7.6	26.9	7.0	26.9	3.0	1.4951	0.0076
SP2014022	4/28/2014	212	M	3	11:54	30	41.0315	-73.2950	8.7	26.3	6.4	27.2	3.2	1.6102	0.0081
SP2014023	4/28/2014	511	M	2	13:08	17	41.0923	-73.3076	7.9	26.6	7.0	26.8	3.1	0.8877	0.0045
SP2014024	4/28/2014	512	M	2	14:00	20	41.0900	-73.3038	8.3	26.6	6.9	26.8	3.5	1.1527	0.0058
SP2014025	4/28/2014	417	T	3	15:48	30	41.0745	-73.0751	7.0	27.2	7.5	28.2	3.3	1.6498	0.0083
SP2014026	4/28/2014	519	M	3	16:41	21	41.0876	-73.0143	7.7	27.2	6.4	27.4	3.1	1.0850	0.0055
SP2014027	4/28/2014	717	M	2	18:10	30	41.1190	-73.0978	8.9	24.2	7.3	26.5	2.6	1.3077	0.0066
SP2014028	4/28/2014	918	T	2	19:07	30	41.1556	-73.0601	8.1	26.8	7.8	27.1	2.8	1.4054	0.0071
SP2014029	5/2/2014	1118	M	1	6:38	30	41.1816	-73.0493	8.3	26.0	7.6	26.9	3.4	1.7122	0.0087
SP2014030	5/2/2014	1019	T	2	7:38	16	41.1652	-73.0198	7.9	26.5	7.2	27.1	3.1	0.8309	0.0042
SP2014031	5/2/2014	1320	M	1	8:52	30	41.2023	-72.9907	9.0	25.1	7.5	26.9	3.2	1.5753	0.0080
SP2014032	5/2/2014	1221	T	2	9:56	23	41.2085	-72.9211	8.2	25.9	7.8	26.6	2.8	1.0607	0.0054
SP2014033	5/2/2014	922	M	3	11:06	30	41.1663	-72.8511	7.5	27.2	7.1	27.1	3.0	1.4798	0.0075
SP2014034	5/2/2014	521	M	4	12:33	30	41.0883	-72.9145	8.1	27.2	7.3	26.4	2.7	1.3267	0.0067
SP2014035	5/2/2014	622	M	4	13:20	30	41.0937	-72.8865	8.4	27.3	7.2	27.3	2.8	1.4011	0.0071
SP2014036	5/2/2014	623	M	4	14:16	30	41.1005	-72.8486	8.1	27.3	7.0	27.3	3.1	1.5524	0.0078
SP2014037	5/2/2014	724	T	4	15:08	30	41.1115	-72.7933	8.3	27.3	7.1	27.3	3.2	1.6115	0.0081
SP2014038	5/2/2014	1427	T	1	16:41	30	41.2358	-72.6572	7.8	25.8	7.2	26.7	3.9	1.9450	0.0098
SP2014039	5/2/2014	1332	S	1	18:01	30	41.2267	-72.4588	7.5	26.1	7.0	27.4	3.8	1.9085	0.0096
SP2014040	5/2/2014	1333	S	1	18:49	21	41.2317	-72.3876	7.6	25.8	7.4	26.9	2.0	0.6895	0.0035

Table 5.6. Station information for LISTS May 2014.

Standard LISTS tows in the spring begin with SP and fall begins with FA. Latitude (N) and Longitude (W) are displayed in decimal degrees. Surface and bottom temperature and salinity are labeled as S_ and B_ respectively. Area swept is estimated by assuming the effective sweep is 2/3rds of the footrope length.

Sample Number	Date	Site Number	Bottom Type	Depth Interval	Time Start	Duration (min)	Latitude	Longitude	S_Temp (sfc, C)	S_Salinity (sfc, ppt)	B_Temp (btm, C)	B_Salinity (btm, ppt)	Ave Speed (knots)	Distance (nm)	Area Swept (sq.nm)
SP2014041	5/12/2014	1336	T	4	7:30	30	41.2116	-72.2923	10.0	26.3	7.7	30.3	1.8	0.9110	0.0046
SP2014042	5/12/2014	1028	T	4	9:46	30	41.1765	-72.5742	8.5	28.0	8.4	28.2	3.2	1.6135	0.0082
SP2014043	5/12/2014	825	T	4	11:25	30	41.1493	-72.7036	10.3	26.3	8.0	27.7	2.8	1.3788	0.0070
SP2014044	5/12/2014	827	T	3	12:42	30	41.1317	-72.6664	11.6	26.2	8.2	27.6	3.6	1.7964	0.0091
SP2014045	5/13/2014	1534	T	1	7:10	18	41.2567	-72.3640	9.6	25.7	9.2	27.1	1.6	0.4938	0.0025
SP2014046	5/13/2014	128	T	2	9:47	30	41.0303	-72.5826	11.8	26.6	9.5	27.3	3.5	1.7352	0.0088
SP2014047	5/13/2014	5824	S	1	11:27	30	40.9810	-72.7325	12.4	26.4	8.4	27.0	3.0	1.5075	0.0076
SP2014048	5/13/2014	124	M	4	12:51	30	41.0192	-72.8021	12.1	26.3	7.5	27.3	3.0	1.5065	0.0076
SP2014049	5/13/2014	125	T	4	13:50	30	41.0108	-72.7428	12.1	26.3	7.7	27.2	3.1	1.5559	0.0079
SP2014050	5/14/2014	1434	S	1	7:08	30	41.2308	-72.4005	10.5	18.3	9.3	27.0	1.7	0.8282	0.0042
SP2014051	5/14/2014	929	S	3	8:43	30	41.1638	-72.5291	10.4	27.3	9.3	27.9	3.6	1.7834	0.0090
SP2014052	5/14/2014	428	S	3	10:03	30	41.0822	-72.5785	10.0	27.4	9.4	27.7	3.7	1.8284	0.0092
SP2014053	5/14/2014	327	T	3	11:17	30	41.0510	-72.6855	12.3	26.4	8.6	27.3	2.8	1.3943	0.0070
SP2014054	5/14/2014	629	S	4	12:44	30	41.1027	-72.5544	11.0	27.2	9.3	27.9	2.9	1.4609	0.0074
SP2014055	5/14/2014	931	S	4	13:58	30	41.1493	-72.4987	10.0	27.8	9.1	28.3	3.2	1.6174	0.0082
SP2014056	5/15/2014	1127	T	3	8:56	30	41.1945	-72.5922	10.7	27.1	9.1	27.7	3.8	1.9022	0.0096
SP2014057	5/15/2014	924	T	3	9:54	30	41.1708	-72.7811	11.7	26.8	10.3	26.6	2.7	1.3274	0.0067
SP2014058	5/15/2014	722	M	3	11:11	30	41.1310	-72.8453	12.7	26.0	8.4	27.2	3.3	1.6520	0.0083
SP2014059	5/15/2014	821	M	3	12:22	30	41.1063	-72.9141	12.7	26.0	8.2	27.1	3.2	1.5863	0.0080
SP2014060	5/15/2014	1022	M	2	13:38	30	41.1808	-72.8455	12.6	26.7	9.8	26.8	3.3	1.6709	0.0084
SP2014061	5/15/2014	1020	T	2	15:12	30	41.1825	-72.9109	13.0	25.9	9.8	26.7	2.6	1.2892	0.0065
SP2014062	5/19/2014	514	M	2	8:24	30	41.0981	-73.1593	12.0	24.6	9.3	26.5	2.9	1.4688	0.0074
SP2014063	5/19/2014	511	M	2	9:32	30	41.1012	-73.2620	12.5	25.5	9.5	26.4	3.2	1.6219	0.0082
SP2014064	5/19/2014	210	T	2	10:40	30	41.0461	-73.3156	12.7	25.4	11.4	25.9	3.2	1.5777	0.0080
SP2014065	5/19/2014	5709	S	2	12:08	18	40.9488	-73.4070	13.1	25.7	12.9	25.8	2.7	0.8108	0.0041
SP2014066	5/19/2014	5513	S	2	14:52	30	40.9236	-73.2488	13.6	25.8	13.2	25.9	3.1	1.5641	0.0079
SP2014067	5/19/2014	5713	T	2	15:53	30	40.9551	-73.2549	13.6	25.8	10.7	26.3	3.2	1.5968	0.0081
SP2014068	5/20/2014	517	T	3	8:10	30	41.1048	-73.0273	12.6	25.9	8.4	26.8	2.6	1.3132	0.0066
SP2014069	5/20/2014	112	M	4	9:58	30	41.0265	-73.2382	13.2	25.0	8.5	26.7	3.1	1.5374	0.0078
SP2014070	5/20/2014	114	M	4	11:11	30	41.0092	-73.2231	13.2	24.8	8.5	26.8	2.7	1.3546	0.0068
SP2014071	5/20/2014	5918	M	3	12:51	30	40.9848	-73.0361	13.1	26.2	10.3	26.5	2.6	1.3189	0.0067
SP2014072	5/20/2014	5921	M	3	14:05	30	40.9880	-72.9135	12.9	26.3	12.2	26.4	2.6	1.2800	0.0065
SP2014073	5/21/2014	1118	M	1	7:41	30	41.1915	-73.0179	11.6	26.3	10.9	26.4	2.8	1.4229	0.0072
SP2014074	5/21/2014	719	M	3	10:01	30	41.1252	-72.9684	13.1	26.0	8.7	26.7	2.9	1.4694	0.0074
SP2014075	5/21/2014	119	M	4	11:31	30	41.0200	-73.0071	13.8	26.1	8.4	27.0	2.7	1.3282	0.0067
SP2014076	5/21/2014	121	M	4	13:20	30	41.0138	-72.9335	14.0	26.3	8.6	26.9	2.5	1.2750	0.0064
SP2014077	5/22/2014	1225	T	2	8:49	30	41.1961	-72.7768	12.7	27.0	11.2	27.1	3.8	1.8910	0.0096
SP2014078	5/22/2014	1425	M	1	10:18	7	41.2375	-72.7283	11.9	27.0	11.8	27.0	2.7	0.3135	0.0016
SP2014079	5/22/2014	1427	T	1	12:20	30	41.2368	-72.6615	11.9	27.0	11.8	27.0	3.2	1.6075	0.0081
SP2014080	5/22/2014	1428	T	1	13:33	30	41.2355	-72.6403	12.0	26.8	11.6	26.9	2.6	1.2907	0.0065

Table 5.7. Station information for LISTS June 2014.

Standard LISTS tows in the spring begin with SP and fall begins with FA. Latitude (N) and Longitude (W) are displayed in decimal degrees. Surface and bottom temperature and salinity are labeled as S_ and B_, respectively. Area swept is estimated by assuming the effective sweep is 2/3rds of the footrope length.

Sample Number	Date	Site Number	Bottom Type	Depth Interval	Time Start	Duration (min)	Latitude	Longitude	S_Temp (sfc, C)	S_Salinity (sfc, ppt)	B_Temp (btm, C)	B_Salinity (btm, ppt)	Ave Speed (knots)	Distance (nm)	Area Swept (sq.nm)
SP2014081	6/9/2014	1738	T	2	7:58	30	41.2860	-72.1976	15.2	29.1	14.7	29.6	3.2	1.5993	0.0081
SP2014082	6/9/2014	1235	T	4	9:48	30	41.2126	-72.2692	15.8	26.9	13.3	30.5	2.2	1.1180	0.0057
SP2014083	6/9/2014	330	S	1	12:21	30	41.0618	-72.4973	16.9	26.4	15.8	26.7	2.4	1.1868	0.0060
SP2014084	6/9/2014	5825	S	1	14:24	30	40.9848	-72.7281	17.3	25.7	17.0	25.8	3.3	1.6747	0.0085
SP2014085	6/10/2014	831	S	4	7:46	30	41.1440	-72.4481	15.9	27.1	13.4	28.8	3.5	1.7717	0.0090
SP2014086	6/10/2014	426	T	3	9:39	30	41.0677	-72.6955	17.8	25.9	13.1	27.6	3.2	1.5853	0.0080
SP2014087	6/10/2014	529	S	3	10:55	30	41.0905	-72.5854	17.6	26.2	13.6	28.4	3.3	1.6424	0.0083
SP2014088	6/10/2014	430	T	3	12:11	30	41.0793	-72.5393	17.0	26.4	13.6	28.3	3.8	1.8760	0.0095
SP2014089	6/11/2014	1432	S	2	7:11	30	41.2328	-72.3953	14.3	28.3	14.4	28.3	3.6	1.8028	0.0091
SP2014090	6/11/2014	931	S	4	8:34	30	41.1610	-72.4445	15.8	27.0	13.8	28.8	3.9	1.9598	0.0099
SP2014091	6/11/2014	729	S	3	10:21	21	41.1147	-72.5811	16.5	26.5	14.2	28.3	2.4	0.8573	0.0043
SP2014092	6/11/2014	229	T	2	12:06	30	41.0363	-72.6064	15.7	26.4	13.5	27.7	3.1	1.5425	0.0078
SP2014093	6/16/2014	1228	T	3	8:02	30	41.2125	-72.5608	15.3	27.6	15.2	27.8	2.8	1.4032	0.0071
SP2014094	6/16/2014	725	T	4	11:19	30	41.1278	-72.6992	16.6	27.1	13.0	27.1	.	.	.
SP2014095	6/16/2014	623	M	4	12:37	30	41.1107	-72.8003	15.1	27.0	13.5	27.5	3.6	1.8107	0.0091
SP2014096	6/16/2014	422	M	4	14:08	30	41.0760	-72.8477	15.2	27.0	13.9	27.5	3.2	1.5869	0.0080
SP2014097	6/17/2014	714	T	1	8:24	30	41.1303	-73.1396	18.2	25.1	17.6	25.5	3.0	1.4751	0.0075
SP2014098	6/17/2014	210	T	2	11:40	30	41.0395	-73.3650	17.9	25.7	13.6	26.1	2.5	1.2475	0.0063
SP2014099	6/17/2014	512	M	2	12:50	30	41.0988	-73.2570	18.7	25.7	15.3	26.0	2.9	1.4280	0.0072
SP2014100	6/17/2014	511	M	2	14:18	20	41.1010	-73.2673	20.3	25.6	15.0	26.0	3.5	1.1522	0.0058
SP2014101	6/18/2014	617	T	2	8:00	30	41.1126	-73.0411	17.0	26.0	12.5	26.5	2.3	1.1749	0.0059
SP2014102	6/18/2014	312	M	3	9:50	30	41.0637	-73.2346	19.2	25.8	13.6	26.2	2.9	1.4416	0.0073
SP2014103	6/18/2014	11	M	4	11:59	30	41.0077	-73.3401	17.8	25.6	12.7	26.3	2.7	1.3375	0.0068
SP2014104	6/18/2014	5513	S	2	13:24	30	40.9268	-73.2478	18.9	25.6	17.8	25.6	2.1	1.0519	0.0053
SP2014105	6/18/2014	5912	M	3	14:49	30	40.9875	-73.2957	18.8	25.6	14.4	26.0	2.8	1.4052	0.0071
SP2014106	6/19/2014	1320	M	1	7:38	30	41.2067	-72.9930	17.0	26.2	14.6	26.4	3.5	1.7342	0.0088
SP2014107	6/19/2014	519	M	3	9:21	30	41.0970	-72.9711	17.6	26.2	12.7	26.8	2.9	1.4378	0.0073
SP2014108	6/19/2014	21	M	3	10:59	30	41.0105	-72.8778	18.1	25.8	13.4	27.3	3.2	1.5928	0.0080
SP2014109	6/19/2014	5922	M	3	12:30	30	40.9918	-72.8373	18.0	25.8	13.9	26.7	3.4	1.6844	0.0085
SP2014110	6/19/2014	118	M	4	13:56	30	41.0305	-73.0023	17.9	25.9	13.7	27.4	3.2	1.5948	0.0081
SP2014111	6/20/2014	321	M	4	8:33	30	41.0530	-72.9257	18.3	26.2	14.4	27.4	3.6	1.7844	0.0090
SP2014112	6/20/2014	924	T	3	10:02	30	41.1353	-72.7675	18.1	26.3	15.3	27.5	3.1	1.5424	0.0078
SP2014113	6/20/2014	1026	T	4	11:43	30	41.1757	-72.6481	17.7	26.5	15.5	27.6	2.7	1.3272	0.0067
SP2014114	6/23/2014	1022	M	2	8:25	30	41.1705	-72.8828	18.7	26.4	14.7	27.1	3.2	1.5870	0.0080
SP2014115	6/23/2014	925	T	4	9:49	30	41.1637	-72.7236	19.3	26.6	16.2	27.8	3.1	1.5672	0.0079
SP2014116	6/23/2014	1125	T	3	12:06	30	41.1591	-72.7081	19.9	26.5	16.1	27.8	3.1	1.5346	0.0078
SP2014117	6/23/2014	1325	T	2	13:25	30	41.2278	-72.7246	18.2	27.2	17.3	27.2	2.8	1.3867	0.0070
SP2014118	6/24/2014	1322	T	1	8:19	30	41.2213	-72.8828	18.8	26.8	17.2	27.0	2.9	1.4662	0.0074
SP2014119	6/24/2014	1425	M	1	10:39	20	41.2372	-72.7314	18.1	27.2	18.1	27.4	3.4	1.1412	0.0058
SP2014120	6/24/2014	1427	T	1	11:51	30	41.2363	-72.6602	18.2	27.5	17.7	27.7	3.9	1.9445	0.0098

Table 5.8. Station information for LISTS September 2014.

Standard LISTS tows in the spring begin with SP and fall begins with FA. Latitude (N) and Longitude (W) are displayed in decimal degrees. Surface and bottom temperature and salinity are labeled as S_ and B_, respectively. Area swept is estimated by assuming the effective sweep is 2/3rds of the footrope length.

Sample Number	Date	Site Number	Bottom Type	Depth Interval	Time Start	Duration (min)	Latitude	Longitude	S_Temp (sfc, C)	S_Salinity (sfc, ppt)	B_Temp (btm, C)	B_Salinity (btm, ppt)	Ave Speed (knots)	Distance (nm)	Area Swept (sq.nm)
FA2014001	9/9/2014	1738	T	2	8:43	30	41.2828	-72.2036	21.2	30.4	21.0	30.4	2.4	1.2122	0.0061
FA2014002	9/9/2014	1840	T	1	11:21	30	41.3298	-72.0856	20.8	29.8	20.2	30.5	3.1	1.5656	0.0079
FA2014003	9/11/2014	830	S	4	7:55	30	41.1480	-72.4853	22.1	28.6	21.8	28.9	3.2	1.6082	0.0081
FA2014004	9/11/2014	328	T	3	9:27	30	41.0616	-72.5876	22.3	28.5	22.2	28.6	3.8	1.9190	0.0097
FA2014005	9/11/2014	123	M	4	10:59	30	41.0375	-72.7970	22.7	27.7	21.7	28.4	3.3	1.6614	0.0084
FA2014006	9/11/2014	5823	S	1	12:23	30	40.9810	-72.8245	22.1	28.1	22.0	28.1	3.3	1.6389	0.0083
FA2014007	9/11/2014	226	T	3	13:48	30	41.0373	-72.6905	22.6	28.3	22.3	28.5	3.3	1.6290	0.0082
FA2014008	9/11/2014	128	T	2	15:01	30	41.0201	-72.6385	22.5	28.3	22.3	28.5	3.8	1.8838	0.0095
FA2014009	9/12/2014	1336	T	4	7:22	30	41.2253	-72.2426	21.0	29.6	20.7	30.0	2.3	1.1568	0.0058
FA2014010	9/12/2014	629	S	4	9:36	30	41.1135	-72.5006	22.0	28.7	22.0	28.7	3.3	1.6496	0.0083
FA2014011	9/12/2014	325	T	3	11:09	30	41.0641	-72.7070	22.3	28.1	22.0	28.4	3.7	1.8269	0.0092
FA2014012	9/12/2014	525	T	4	12:24	30	41.0840	-72.7667	22.6	28.0	22.1	28.3	2.4	1.2036	0.0061
FA2014013	9/12/2014	828	S	3	13:56	30	41.1373	-72.6146	22.2	28.9	22.0	28.9	2.7	1.3411	0.0068
FA2014014	9/12/2014	1029	S	3	15:06	30	41.1638	-72.5888	22.3	28.8	21.7	29.1	3.2	1.5949	0.0081
FA2014015	9/15/2014	1228	T	3	8:11	30	41.2137	-72.5510	21.1	28.6	20.7	28.9	2.3	1.1747	0.0059
FA2014016	9/15/2014	623	M	4	12:13	30	41.1111	-72.7953	21.9	27.9	21.5	28.1	3.5	1.7453	0.0088
FA2014017	9/15/2014	523	M	4	13:35	30	41.0805	-72.8508	22.0	27.8	21.7	28.2	2.5	1.2390	0.0063
FA2014018	9/15/2014	824	T	4	15:04	30	41.1275	-72.8065	22.3	28.2	21.8	28.4	2.6	1.2811	0.0065
FA2014019	9/16/2014	1332	S	1	7:22	30	41.2311	-72.3991	20.3	29.1	20.3	29.3	1.8	0.9218	0.0047
FA2014020	9/16/2014	926	T	4	9:27	30	41.1645	-72.6320	21.4	28.4	21.5	28.6	2.4	1.2161	0.0061
FA2014021	9/16/2014	1024	T	3	10:42	30	41.1837	-72.7308	21.5	28.1	21.5	28.1	2.7	1.3576	0.0069
FA2014022	9/16/2014	1123	M	2	11:52	30	41.1915	-72.7881	21.4	27.9	21.4	27.9	3.1	1.5250	0.0077
FA2014023	9/16/2014	920	T	2	13:25	30	41.1642	-72.9235	21.0	27.6	20.9	27.6	3.4	1.6838	0.0085
FA2014024	9/16/2014	919	T	2	14:43	30	41.1507	-72.9905	21.4	27.6	21.2	27.6	2.8	1.3858	0.0070
FA2014025	9/17/2014	615	M	2	8:20	30	41.0905	-73.1498	21.2	27.5	21.2	27.6	2.9	1.4633	0.0074
FA2014026	9/17/2014	611	M	1	9:45	30	41.1108	-73.2733	20.9	27.2	20.9	27.2	3.2	1.5805	0.0080
FA2014027	9/17/2014	5709	S	2	11:40	30	40.9468	-73.4084	21.6	27.1	21.4	27.1	3.1	1.5585	0.0079
FA2014028	9/17/2014	5912	M	3	13:17	30	40.9853	-73.3021	21.8	27.4	21.8	27.6	3.0	1.5182	0.0077
FA2014029	9/17/2014	5513	S	2	14:48	30	40.9248	-73.2523	22.0	27.1	20.9	27.0	3.1	1.5731	0.0079
FA2014030	9/17/2014	5914	M	4	16:06	27	40.9930	-73.2029	22.3	27.4	21.9	27.8	2.7	1.2285	0.0062
FA2014031	9/18/2014	1020	T	2	8:35	30	41.1815	-72.9251	20.6	27.6	20.6	27.7	3.1	1.5347	0.0078
FA2014032	9/18/2014	620	M	3	9:51	30	41.1158	-72.9310	21.4	28.0	21.6	28.2	2.8	1.4152	0.0072
FA2014033	9/18/2014	5917	M	3	11:43	30	40.9923	-73.0558	21.6	27.8	21.6	28.2	3.3	1.6447	0.0083
FA2014034	9/18/2014	118	M	4	13:19	20	41.0230	-73.0510	22.0	27.9	21.6	28.3	2.9	0.9541	0.0048
FA2014035	9/18/2014	518	M	3	14:44	30	41.0903	-73.0566	22.5	27.8	21.5	28.6	3.0	1.4934	0.0075
FA2014036	9/18/2014	719	M	3	15:57	30	41.1153	-73.0173	21.5	27.7	21.5	28.1	2.9	1.4431	0.0073
FA2014037	9/23/2014	1119	M	2	7:43	30	41.1883	-73.0049	20.4	27.6	20.3	27.6	2.9	1.4466	0.0073
FA2014038	9/23/2014	1320	M	1	9:15	30	41.2075	-72.9825	20.3	27.5	20.1	27.5	3.2	1.5818	0.0080
FA2014039	9/23/2014	1322	T	1	10:37	30	41.2228	-72.8733	20.0	27.6	20.1	27.6	3.1	1.5253	0.0077
FA2014040	9/23/2014	1428	T	1	12:28	30	41.2375	-72.6310	20.5	28.6	20.3	28.8	4.0	1.9970	0.0101

Table 5.9. Station information for LISTS October 2014.

Standard LISTS tows in the spring begin with SP and fall begins with FA. Latitude (N) and Longitude (W) are displayed in decimal degrees. Surface and bottom temperature and salinity are labeled as S_ and B_, respectively. Area swept is estimated by assuming the effective sweep is 2/3rds of the footrope length

Sample Number	Date	Site Number	Bottom Type	Depth Interval	Time Start	Duration (min)	Latitude	Longitude	S_Temp (sfc, C)	S_Salinity (sfc, ppt)	B_Temp (btm, C)	B_Salinity (btm, ppt)	Ave Speed (knots)	Distance (nm)	Area Swept (sq.nm)
FA2014041	10/14/2014	1332	S	1	7:37	30	41.2320	-72.3956	17.9	29.0	18.1	29.7	2.2	1.0957	0.0055
FA2014042	10/14/2014	1027	T	4	9:44	30	41.1808	-72.6466	18.3	28.4	18.2	28.8	2.7	1.3613	0.0069
FA2014043	10/14/2014	624	T	4	11:12	30	41.1187	-72.7563	18.6	28.4	18.4	28.8	3.2	1.5909	0.0080
FA2014044	10/14/2014	5921	M	3	12:56	30	41.0000	-72.8583	18.6	28.2	18.6	28.2	2.8	1.3879	0.0070
FA2014045	10/15/2014	1118	M	1	7:48	30	41.1898	-73.0222	17.6	27.6	17.6	27.6	2.9	1.4472	0.0073
FA2014046	10/15/2014	15	T	4	9:48	30	41.0083	-73.1256	18.8	27.9	18.8	28.5	2.6	1.2945	0.0065
FA2014047	10/15/2014	5613	T	2	11:57	30	40.9493	-73.1891	18.2	27.5	19.0	28.0	3.0	1.4966	0.0076
FA2014048	10/15/2014	5513	S	2	13:47	30	40.9280	-73.2540	18.5	27.4	18.0	27.5	3.1	1.5696	0.0079
FA2014049	10/16/2014	5920	M	2	9:19	30	40.9851	-72.9490	19.0	28.0	19.0	28.0	3.4	1.7108	0.0086
FA2014050	10/16/2014	5923	M	3	10:42	30	40.9870	-72.8038	18.8	27.6	18.7	28.3	3.2	1.6217	0.0082
FA2014051	10/16/2014	5925	T	1	12:29	30	41.0020	-72.7085	18.8	28.1	18.8	28.2	3.3	1.6267	0.0082
FA2014052	10/17/2014	714	T	1	8:17	30	41.1313	-73.1349	18.4	27.2	18.5	27.6	2.6	1.2754	0.0064
FA2014053	10/17/2014	612	M	1	9:39	30	41.1100	-73.2678	18.6	27.5	18.8	27.8	2.7	1.3608	0.0069
FA2014054	10/17/2014	511	M	2	10:53	30	41.1017	-73.2643	18.6	27.6	18.5	27.8	2.8	1.3767	0.0070
FA2014055	10/17/2014	412	M	2	12:21	23	41.0660	-73.3085	18.7	27.8	18.7	27.9	3.0	1.1341	0.0057
FA2014056	10/20/2014	211	T	2	9:09	30	41.0498	-73.3044	18.1	27.8	18.3	28.1	2.8	1.3760	0.0070
FA2014057	10/20/2014	5911	M	3	11:18	30	41.0010	-73.2738	18.1	27.6	18.4	28.0	2.7	1.3368	0.0068
FA2014058	10/20/2014	17	M	4	13:37	30	41.0067	-73.0760	18.1	27.8	18.3	28.5	2.8	1.4113	0.0071
FA2014059	10/21/2014	118	M	4	8:40	30	41.0210	-73.0478	18.1	28.0	18.4	28.6	2.6	1.2822	0.0065
FA2014060	10/21/2014	222	M	4	10:36	30	41.0322	-72.8888	17.7	28.2	18.1	28.8	2.9	1.4692	0.0074
FA2014061	10/21/2014	5919	M	3	13:53	30	40.9900	-73.0358	18.0	27.9	18.1	28.1	3.4	1.6762	0.0085
FA2014062	10/28/2014	1432	S	2	7:26	30	41.2338	-72.3904	16.2	28.5	16.5	29.2	2.5	1.2679	0.0064
FA2014063	10/28/2014	1126	T	3	9:33	30	41.1995	-72.6628	16.0	28.7	16.1	28.8	3.3	1.6286	0.0082
FA2014064	10/28/2014	419	M	4	11:51	30	41.0758	-72.9691	16.8	28.1	16.7	28.2	3.4	1.7195	0.0087
FA2014065	10/28/2014	1029	S	3	15:04	30	41.1587	-72.5986	16.7	29.0	16.8	29.1	3.0	1.5082	0.0076
FA2014066	10/29/2014	1737	T	1	7:42	30	41.2903	-72.1965	16.6	30.5	16.6	30.6	3.2	1.6059	0.0081
FA2014067	10/29/2014	1533	S	1	9:46	30	41.2563	-72.3738	16.3	27.7	16.6	29.0	2.6	1.3042	0.0066
FA2014068	10/29/2014	630	S	4	12:06	30	41.1071	-72.4960	16.6	28.6	16.5	28.9	3.0	1.4764	0.0075
FA2014069	10/29/2014	526	T	3	14:29	30	41.1002	-72.6331	16.9	28.4	16.7	28.9	3.4	1.6863	0.0085
FA2014070	10/30/2014	1235	T	4	7:27	30	41.2140	-72.2636	16.4	29.9	16.5	30.9	1.8	0.8953	0.0045
FA2014071	10/30/2014	430	T	3	9:54	30	41.0862	-72.4915	16.0	28.4	16.4	28.9	2.8	1.3869	0.0070
FA2014072	10/30/2014	325	T	3	11:40	30	41.0652	-72.7001	16.3	28.8	16.3	29.0	3.3	1.6604	0.0084
FA2014073	10/30/2014	623	M	4	13:19	24	41.0977	-72.8477	16.4	28.3	16.7	28.7	2.5	0.9834	0.0050
FA2014074	10/31/2014	1124	T	2	9:07	30	41.2011	-72.7343	15.7	28.8	15.7	28.8	2.8	1.3912	0.0070
FA2014075	10/31/2014	821	M	3	10:55	30	41.1088	-72.9043	16.3	28.3	16.3	28.3	3.1	1.5325	0.0077
FA2014076	10/31/2014	1224	T	2	12:54	30	41.1958	-72.8143	16.0	28.0	16.1	28.3	2.9	1.4546	0.0074
FA2014077	11/4/2014	228	T	2	8:36	30	41.0427	-72.5558	14.3	28.6	14.3	28.6	2.5	1.2545	0.0063
FA2014078	11/4/2014	527	T	3	10:23	30	41.0918	-72.6603	15.0	28.5	14.9	28.5	3.7	1.8696	0.0094
FA2014079	11/4/2014	828	S	3	12:26	28	41.1393	-72.6155	15.0	28.6	14.9	28.7	3.0	1.4177	0.0072

Table 5.10. Samples with non-standard tow durations and reasons for incomplete tows, spring and fall 2014.*Standard LISTS tows begin with SP (spring) or FA (fall).*

Sample	Date	Site	Bottom Type	Depth Interval	Time	Duration	Reason	Comments
APRIL								
SP2014023	4/28/2014	0511	M	2	13:08	17	speed drop	speed dropped but net seemed to pop off hang during haul-back; one old pot in net
SP2014024	4/28/2014	0512	M	2	14:00	20	pots	snagged string of pots with submerged buoy weighted down by mussels
SP2014026	4/28/2014	0519	M	3	16:41	21	pots	snagged string of old gear (2011 trap tag)
SP2014030	5/2/2014	1019	T	2	7:38	16	hang	hang damaged the net; had to change nets
SP2014032	5/2/2014	1221	T	2	9:56	23	hang	came off during haul-back; no damage to net
SP2014040	5/2/2014	1333	S	1	18:49	21	speed drop	speed dropped but no gear or debris in net
MAY								
SP2014045	5/13/2014	1534	T	1	7:10	18	speed drop	net loaded up with debris and algae
SP2014065	5/19/2014	5709	S	2	12:08	18	pots	buoy on port door, active and ghost gear; tree limbs in net; damage to net
SP2014078	5/22/2014	1425	M	1	10:18	7	hangs	2 attempts; kept hanging up on bottom; snagged large section of rigid pipe; significant damage to net; had to change nets
JUNE								
SP2014091	6/11/2014	0729	S	3	10:21	21	pots	2 attempts; 1st part speed dropped but no gear or debris in net; 2nd attempt buoy tangled with door, no pots onboard
SP2014100	6/17/2014	0511	M	2	14:18	20	pots	ghost pots in net (trap tags from 2004)
SP2014119	6/24/2014	1425	M	1	10:39	20	hangs	2 attempts; hung on bottom; damage to net; changed net
SEPT								
FA2014030	9/17/2014	5914	M	4	16:06	27	speed drop	speed drop but no gear or debris on net
FA2014034	9/18/2014	0118	M	4	13:19	20	speed drop	speed drop but no gear or debris on net
OCT								
FA2014055	10/17/2018	0412	M	2	12:21	23	speed drop	speed drop but no gear or debris on net
FA2014073	10/30/2014	0623	M	4	13:19	24	pots	snagged active gear set N-S (2014 trap tags); re-tied pots
FA2014079	11/4/2014	0828	S	3	12:26	28	rough bottom	felt rough bottom just before boost, so hauled back
FA2014080	11/4/2014	1133	S	4	15:00	0	hnag / dunes	3 attempts; on third attempt substantial damage to net & ran out of daylight

Table 5.11. Data requests by month, 2014.

MONTH	REQUEST	ORGANIZATION OR PURPOSE
January	LISTS sturgeon catch data, 1984-2013	ASMFC
	LISTS squid indices & biological info	university
	LISTS BADD index for LISS	EPA
	LISTS sampling locations for dump site EIS (GIS)	EPA
February	LISTS BSB length frequencies	CT DEEP / ASMFC
	LISTS methodology	LeHigh University
	LISTS winter flounder age matrix & ESS indices, 1984-2013	CT DEEP
	LISTS winter flounder age matrix, 1984-2013	Dominion
	LISTS winter flounder age matrix	NMFS
	LISTS towpaths (GIS)	CT DOA Aquaculture
	LISTS tow points (GIS)	EPA
March	menhaden catch & indices from LISTS & seine surveys	Dominion
	LISTS summer flounder counts & lengths	NY DEC
	LISTS species indicators for LISS	EPA
	summary of CT DEEP trawl surveys in LIS	NMFS
	CT DEEP trawl survey data for SFL, WFL	NMFS
	LISTS BSB indices at length	ASMFC
	menhaden indices for LISTS, CT & Th riverine surveys	ASMFC
	LISTS site grid (GIS)	ASMFC
April	LISTS tautog age matrix, 1984-2012	ASMFC
	LISTS time-series warm/cold, lobster indices & species richness	CT CEQ
	LISTS count & biomass indices, 1984-2013	Normandeau Assoc.
	LISTS scup & fluke indices & age keys, 1984-2013	NMFS
	LISTS site grid (GIS)	ASMFC
May	LISTS catch data for BLF,BSB,PGY,RED,SFL,SQI,WFL,WHI,WPF	NOAA
	LISTS indices and SMD data, 1984-2013	NOAA
	LISTS site grid (GIS)	Uconn / CT DEEP
June		
July	LISTS lobster catch & length data, 2014	CT DEEP / media
August	LISTS lobster catch data, 1984-2013	Univ of Maine
September	LISTS BSB indices at length	ASMFC
	LISTS species distributions for 13 Priority GCN species (GIS)	CT DEEP
October		
November	LISTS abundance indices	CT news media
	black sea bass	CT DEEP / ASMFC
December	LISTS lobster catch & length data, 2014	CT DEEP / ASMFC
	LISTS tows, counts, lengths (1984-2013) flat files	NMFS NEFSC

Table 5.12. Sample requests by month, 2014.

MONTH	REQUEST	ORGANIZATION OR PURPOSE
April	squid & various finfish specimens for dissection class	Putnam High School
	various species for decomposition studies	UNH
	various species for microscopic tissue structure studies for spp ID	UNH
	dogfish for parasite studies	Uconn
	channeled and knobbed whelk (conch)	NY DEC
May	squid & various finfish specimens for dissection class	Putnam High School
	tautog tissue samples for DNA study	VIMS
	dogfish for parasite studies	Uconn
	channeled and knobbed whelk (conch)	NY DEC
	various species for decomposition studies	UNH
	various species for microscopic tissue structure studies for spp ID	UNH
	various critters for environmental outreach program	M.E.N.
June	channeled and knobbed whelk (conch)	NY DEC
	hermit crabs	UConn
July		
August		
September	channeled and knobbed whelk (conch)	NY DEC
	various species for fish biology lab	Uconn
October	channeled and knobbed whelk (conch)	NY DEC
	squid	CT DEEP - CARE
November	various fish species	CT DEEP - Encon Police

Table 5.13. List of finfish species observed in 2014.

Sixty finfish species were observed in 2014. (Bold type indicates new species). Since 1984, one hundred-six species of finfish have been identified in LISTS (see Appendix 5.1 for the full list of species).

Common Name	Scientific Name	Common Name	Scientific Name
anchovy, bay	Anchoa mitchilli	lookdown	Selene vomer
black sea bass	Centropristis striata	mackerel, Atlantic	Scomber scombrus
blue runner	Caranx crysos	menhaden, Atlantic	Brevoortia tyrannus
bluefish	Pomatomus saltatrix	moonfish	Selene setapinnis
butterfish	Peprilus triacanthus	perch, silver	Bairdiella chrysoura
cod, Atlantic	Gadus morhua	perch, white	Morone americana
cornetfish, blue spotted	Fistularia tabacaria	pipefish, northern	Syngnathus fuscus
croaker, Atlantic	Micropogonias undulatus	puffer, northern	Sphoeroides maculatus
cunner	Tautoglabrus adspersus	rockling, fourbeard	Enchelyopus cimbrius
cusck-eel, striped	Ophidion marginatum	rudderfish, banded	Seriola zonata
dogfish, smooth	Mustelus canis	sand lance, American	Ammodytes americanus
dogfish, spiny	Squalus acanthias	scad, mackerel	Decapterus macarellus
drum, black	Pogonias cromis	scad, rough	Trachurus lathami
filefish, planehead	Monacanthus hispidus	scad, round	Decapterus punctatus
flounder, fourspot	Paralichthys oblongus	scup	Stenotomus chrysops
flounder, smallmouth	Etropus microstomus	sea raven	Hemitripterus americanus
flounder, summer	Paralichthys dentatus	searobin, northern	Prionotus carolinus
flounder, windowpane	Scophthalmus aquosus	searobin, striped	Prionotus evolans
flounder, winter	Pseudopleuronectes american	shad, American	Alosa sapidissima
goatfish, red	Mullus auratus	shad, hickory	Alosa mediocris
hake, red	Urophycis chuss	silverside, Atlantic	Menidia menidia
hake, silver	Merluccius bilinearis	skate, clearnose	Raja eglanteria
hake, spotted	Urophycis regia	skate, little	Leucoraja erinacea
herring, Atlantic	Clupea harengus	skate, winter	Leucoraja ocellata
herring, alewife	Alosa pseudoharengus	spot	Leiostomus xanthurus
herring, blueback	Alosa aestivalis	striped bass	Morone saxatilis
hogchoker	Trinectes maculatus	sturgeon, Atlantic	Acipenser oxyrinchus
jack, crevalle	Caranx hippos	tautog	Tautoga onitis
kingfish, northern	Menticirrhus saxatilis	toadfish, oyster	<i>Opsanus tau</i>
lizardfish, inshore	Synodus foetens	weakfish	<i>Cynoscion regalis</i>

Names taken from: Common and Scientific Names of Fishes from the United States, Canada and Mexico, American Fisheries Society, Sixth ed., 2004.

Table 5.14. List of invertebrates observed in 2014.

In 2014, forty-two invertebrate "species" were identified. In most cases, invertebrates are identified to species; however, species that are very similar are identified to genus, and in difficult cases, to a higher taxon.

Common Name	Scientific Name	Common Name	Scientific Name
Tubularia hydroids	Tubularia, spp.	mussel, blue	Mytilus edulis
anemones	anemomes spp.	mussel, ribbed	Geukensia demissa
arks	Noetia-Anadara spp.	northern moon snail	Lunatia heros
bryozoan, bushy	Phylum Bryozoa	oyster, common	Crassostrea virginica
bryozoan, rubbery	Alcyonidium verrilli	sea grape	Molgula spp.
clam, common razer	Ensis directus	sea urchin, purple	Arbacia punctulata
clam, hard clams	Artica-Mercinaria-Pitar sp.	shrimp, brown	Penaeus aztecus
clam, surf	Spisula solidissima	shrimp, coastal mud	Upogebia affinis
coral, star	Astrangia poculata	shrimp, ghost	Gilvossius setimanus
crab, mud	Family Xanthidae	shrimp, mantis	Squilla empusa
crab, blue	Callinectes sapidus	shrimp, sand	Crangon septemspinosa
crab, flat claw hermit	Pagurus pollicaris	slipper shell, common	Crepidula fornicata
crab, horseshoe	Limulus polyphemus	sponge spp.	sponge spp.
crab, lady	Ovalipes ocellatus	sponge, boring	Cliona celate
crab, rock	Cancer irroratus	sponge, deadman's fingers	Haliclona spp.
crab, spider	Libinia emarginata	sponge, red bearded	Microciona prolifera
hydroid spp.	hydroid spp.	squid, longfin inshore	Loligo pealeii
jelly, comb	Phylum Ctenophora	starfish spp.	Asteriid spp.
jelly, water	Rhacostoma atlanticum	tunicates, misc	misc. class ascidiacea
jellyfish, lion's mane	Cyanea capillata	whelk, channeled	Busycotypus canaliculatus
lobster, American	Homarus americanus	whelk, knobbed	Busycon carica

Names taken from: A Field Guide to the Atlantic Seashore, Peterson Field Guide Series, 1978 (Gosner, 1978).

Table 5.15. Total number and weight (kg) of finfish and invertebrates caught in 2014.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring and American sand lance are not quantified. Number of tows (sample size)=199.

species	count	%	weight	%	species	count	%	weight	%	
butterfish	69,372	45.3	1,707.6	10.6	1					
scup	45,705	29.9	5,161.4	31.9	2					
weakfish	10,477	6.8	334.8	2.1	3					
bluefish	4,457	2.9	522.7	3.2	4					
northern searobin	2,584	1.7	225.9	1.4	5					
striped searobin	2,544	1.7	1,020.8	6.3	6					
moonfish	2,200	1.4	23.2	0.1	7					
windowpane flounder	2,191	1.4	365.6	2.3	8					
Atlantic herring	1,838	1.2	91.2	0.6	9	longfin inshore squid	13,436	86.3	582.3	37.9
bay anchovy	1,424	0.9	9.4	0.1	10	horseshoe crab	261	1.7	497.3	32.4
winter flounder	1,372	0.9	459.7	2.8	11	spider crab	nc		145.6	9.5
black sea bass	1,295	0.8	543.3	3.4	12	blue mussel	nc		52.2	3.4
smooth dogfish	1,197	0.8	2,799.2	17.3	13	lion's mane jellyfish	1,262	8.1	48.2	3.1
summer flounder	859	0.6	567.4	3.5	14	American lobster	178	1.1	31.5	2.1
fourspot flounder	820	0.5	145.0	0.9	15	bushy bryozoan	nc		24.8	1.6
little skate	770	0.5	428.2	2.6	16	mixed sponge species	nc		20.6	1.3
Atlantic menhaden	723	0.5	267.8	1.7	17	common slipper shell	nc		18.8	1.2
alewife	555	0.4	43.2	0.3	18	mantis shrimp	332	2.1	14.4	0.9
spotted hake	505	0.3	59.5	0.4	19	flat claw hermit crab	nc		14.0	0.9
red hake	398	0.3	33.5	0.2	20	knobbed whelk	34	0.2	12.3	0.8
silver hake	323	0.2	10.6	0.1	21	lady crab	nc		9.3	0.6
striped bass	255	0.2	407.5	2.5	22	sea grape	nc		7.3	0.5
hogchoker	246	0.2	27.8	0.2	23	channeled whelk	29	0.2	5.9	0.4
tautog	194	0.1	192.5	1.2	24	hydroid spp.	nc		5.3	0.3
American shad	162	0.1	12.3	0.1	25	rock crab	nc		4.8	0.3
smallmouth flounder	152	0.1	6.0	0.0	26	northern moon snail	nc		4.6	0.3
clearnose skate	104	0.1	207.7	1.3	27	Tubularia, spp.	nc		4.6	0.3
winter skate	82	0.1	133.8	0.8	28	boring sponge	nc		4.3	0.3
blueback herring	58	0.0	4.2	0.0	29	sand shrimp	nc		4.1	0.3
northern kingfish	51	0.0	3.2	0.0	30	blue crab	18	0.1	3.0	0.2
hickory shad	30	0.0	10.5	0.1	31	arks	nc		2.7	0.2
inshore lizardfish	30	0.0	2.8	0.0	32	mud crabs	nc		2.6	0.2
spot	20	0.0	1.8	0.0	33	starfish spp.	2	0.0	1.6	0.1
spiny dogfish	15	0.0	62.2	0.4	34	ribbed mussel	nc		1.6	0.1
Atlantic sturgeon	13	0.0	272.4	1.7	35	comb jelly spp	nc		1.4	0.1
American sand lance	12	0.0	0.2	0.0	36	star coral	nc		0.7	0.0
blue runner	10	0.0	0.9	0.0	37	purple sea urchin	4	0.0	0.6	0.0
northern puffer	10	0.0	1.3	0.0	38	surf clam	4	0.0	0.5	0.0
striped cusk-eel	6	0.0	0.6	0.0	39	coastal mud shrimp	1	0.0	0.3	0.0
Atlantic cod	5	0.0	0.3	0.0	40	rubbery bryzoan	nc		0.3	0.0
rough scad	5	0.0	0.5	0.0	41	tunicates, misc	nc		0.3	0.0
planehead filefish	4	0.0	0.4	0.0	42	anemones	5	0.0	0.2	0.0
fourbeard rockling	4	0.0	0.4	0.0	43	brown shrimp	2	0.0	0.2	0.0
crevalle jack	2	0.0	0.2	0.0	44	common razor clam	1	0.0	0.2	0.0
Atlantic croaker	2	0.0	0.2	0.0	45	hard clams	nc		0.2	0.0
cunner	2	0.0	0.2	0.0	46	common oyster	nc		0.2	0.0
Atlantic mackerel	2	0.0	0.2	0.0	47	red bearded sponge	nc		0.1	0.0
silver perch	2	0.0	0.2	0.0	48	deadman's fingers sponge	nc		0.1	0.0
oyster toadfish	2	0.0	0.6	0.0	49	ghost shrimp	1	0.0	0.1	0.0
Atlantic silverside	1	0.0	0.1	0.0	50	water jelly	1	0.0	0.1	0.0
black drum	1	0.0	0.1	0.0	51	Total	15,571		1,529.2	
blue spotted cornetfish	1	0.0	0.1	0.0	52	Note: nc= not counted				
lookdown	1	0.0	0.1	0.0	53					
mackerel scad	1	0.0	0.1	0.0	54					
northern pipefish	1	0.0	0.1	0.0	55					
round scad	1	0.0	0.1	0.0						
red goatfish	1	0.0	0.1	0.0						
banded rudderfish	1	0.0	0.4	0.0						
sea raven	1	0.0	1.5	0.0						
white perch	1	0.0	0.2	0.0						
Total	153,100		16,173.8							

Table 5.16. Total counts and weight (kg) of finfish taken in the spring and fall sampling periods, 2014.

Species are listed in order of descending count.. Young-of-year bay anchovy, striped anchovy, Atlantic herring and American sand lance are not included. Number of tows (sample sizes): Spring = 120 and Fall=79.

Spring					Fall				
species	count	%	weight	%	species	count	%	weight	%
scup	15,531	45.8	3,647.2	41.4	butterfish	66,020	55.4	1,491.2	20.3
butterfish	3,352	9.9	216.4	2.5	scup	30,174	25.3	1,514.2	20.6
Atlantic herring	1,835	5.4	90.9	1.0	weakfish	10,454	8.8	316.5	4.3
northern searobin	1,756	5.2	201.8	2.3	bluefish	4,451	3.7	515.3	7.0
windowpane flounder	1,746	5.1	309.3	3.5	moonfish	2,200	1.8	23.2	0.3
striped searobin	1,690	5.0	698.2	7.9	striped searobin	854	0.7	322.6	4.4
winter flounder	1,244	3.7	431.9	4.9	northern searobin	828	0.7	24.1	0.3
black sea bass	1,058	3.1	412.2	4.7	smooth dogfish	800	0.7	1,820.8	24.7
bay anchovy	892	2.6	5.3	0.1	bay anchovy	532	0.4	4.1	0.1
summer flounder	675	2.0	412.2	4.7	Atlantic menhaden	454	0.4	158.1	2.1
fourspot flounder	659	1.9	136.4	1.5	windowpane flounder	445	0.4	56.3	0.8
little skate	522	1.5	292.8	3.3	little skate	248	0.2	135.4	1.8
alewife	450	1.3	39.2	0.4	black sea bass	238	0.2	131.1	1.8
smooth dogfish	397	1.2	978.4	11.1	spotted hake	187	0.2	34.3	0.5
spotted hake	318	0.9	25.2	0.3	summer flounder	184	0.2	155.2	2.1
silver hake	309	0.9	9.4	0.1	fourspot flounder	161	0.1	8.6	0.1
red hake	275	0.8	18.5	0.2	winter flounder	128	0.1	27.8	0.4
Atlantic menhaden	269	0.8	109.7	1.2	red hake	123	0.1	15.0	0.2
striped bass	206	0.6	298.4	3.4	smallmouth flounder	121	0.1	4.0	0.1
hogchoker	165	0.5	18.8	0.2	alewife	105	0.1	4.0	0.1
tautog	158	0.5	167.4	1.9	hogchoker	81	0.1	9.0	0.1
American shad	121	0.4	9.8	0.1	clearnose skate	62	0.1	136.4	1.9
winter skate	68	0.2	110.1	1.2	northern kingfish	51	0.0	3.2	0.0
blueback herring	48	0.1	3.7	0.0	striped bass	49	0.0	109.1	1.5
clearnose skate	42	0.1	71.3	0.8	American shad	40	0.0	2.5	0.0
smallmouth flounder	32	0.1	2.0	0.0	tautog	36	0.0	25.1	0.3
hickory shad	24	0.1	8.0	0.1	inshore lizardfish	30	0.0	2.8	0.0
weakfish	23	0.1	18.3	0.2	spot	20	0.0	1.8	0.0
spiny dogfish	15	0.0	62.2	0.7	silver hake	14	0.0	1.2	0.0
American sand lance	12	0.0	0.2	0.0	winter skate	14	0.0	23.7	0.3
bluefish	6	0.0	7.4	0.1	Atlantic sturgeon	13	0.0	272.4	3.7
striped cusk-eel	6	0.0	0.6	0.0	blueback herring	10	0.0	0.5	0.0
Atlantic cod	5	0.0	0.3	0.0	blue runner	10	0.0	0.9	0.0
fourbeard rockling	4	0.0	0.4	0.0	northern puffer	9	0.0	1.2	0.0
silver perch	2	0.0	0.2	0.0	hickory shad	6	0.0	2.5	0.0
Atlantic silverside	1	0.0	0.1	0.0	rough scad	5	0.0	0.5	0.0
northern puffer	1	0.0	0.1	0.0	planehead filefish	4	0.0	0.4	0.0
sea raven	1	0.0	1.5	0.0	Atlantic herring	3	0.0	0.3	0.0
white perch	1	0.0	0.2	0.0	crevalle jack	2	0.0	0.2	0.0
Total	33,919		8,816.0		Atlantic croaker	2	0.0	0.2	0.0
					cunner	2	0.0	0.2	0.0
					Atlantic mackerel	2	0.0	0.2	0.0
					oyster toadfish	2	0.0	0.6	0.0
					black drum	1	0.0	0.1	0.0
					blue spotted cornetfish	1	0.0	0.1	0.0
					lookdown	1	0.0	0.1	0.0
					mackerel scad	1	0.0	0.1	0.0
					northern pipefish	1	0.0	0.1	0.0
					round scad	1	0.0	0.1	0.0
					red goat fish	1	0.0	0.1	0.0
					banded rudderfish	1	0.0	0.4	0.0
					Total	119,182		7,357.8	

Table 5.17. Total catch of invertebrates taken in the spring and fall sampling periods, 2014.
Species are ranked by total weight (kg). Number of tows (sample sizes): Spring = 120 and Fall=79.

species	Spring				species	Fall			
	count	%	weight	%		count	%	weight	%
horseshoe crab	175	5.2	329.4	37.8	longfin inshore squid	11,813	96.5	408.8	61.5
longfin inshore squid	1,623	48.7	173.5	19.9	horseshoe crab	86	0.7	167.9	25.3
spider crab	nc		136.1	15.6	blue mussel	nc		10.1	1.5
blue mussel	nc		42.1	4.8	mantis shrimp	239	2.0	10.1	1.5
lion's mane jellyfish	1,228	36.8	41.9	4.8	spider crab	nc		9.5	1.4
American lobster	167	5.0	27.8	3.2	knobbed whelk	22	0.2	8.7	1.3
bushy bryozoan	nc		23.5	2.7	common slipper shell	nc		8.7	1.3
mixed sponge species	nc		20.5	2.4	lady crab	nc		7.6	1.1
common slipper shell	nc		10.1	1.2	lion's mane jellyfish	34	0.3	6.3	0.9
flat claw hermit crab	nc		9.4	1.1	flat claw hermit crab	nc		4.6	0.7
sea grape	nc		7.2	0.8	boring sponge	nc		4.3	0.6
northern moon snail	nc		4.6	0.5	American lobster	11	0.1	3.7	0.6
channeled whelk	22	0.7	4.5	0.5	blue crab	15	0.1	2.7	0.4
Tubularia, spp.	nc		4.5	0.5	hydroid spp.	nc		2.2	0.3
mantis shrimp	93	2.8	4.3	0.5	channeled whelk	7	0.1	1.4	0.2
sand shrimp	nc		4.0	0.5	bushy bryozoan	nc		1.3	0.2
rock crab	nc		4.0	0.5	arks	nc		1.2	0.2
knobbed whelk	12	0.4	3.6	0.4	rock crab	nc		0.8	0.1
hydroid spp.	nc		3.1	0.4	mud crabs	nc		0.7	0.1
mud crabs	nc		1.9	0.2	starfish spp.	nc		0.5	0.1
lady crab	nc		1.7	0.2	surf clam	3	0.0	0.3	0.0
ribbed mussel	nc		1.6	0.2	brown shrimp	2	0.0	0.2	0.0
arks	nc		1.5	0.2	star coral	nc		0.2	0.0
comb jelly spp	nc		1.3	0.1	common razor clam	1	0.0	0.2	0.0
starfish spp.	2	0.0	1.1	0.1	common oyster	nc		0.2	0.0
star coral	nc		0.5	0.1	coastal mud shrimp	1	0.0	0.1	0.0
purple sea urchin	3	0.1	0.5	0.1	comb jelly spp	nc		0.1	0.0
blue crab	3	0.1	0.3	0.0	sand shrimp	nc		0.1	0.0
tunicates, misc	nc		0.3	0.0	deadman's fingers sponge	nc		0.1	0.0
anemones	5	0.2	0.2	0.0	rubbery bryozoan	nc		0.1	0.0
coastal mud shrimp	nc		0.2	0.0	sea grape	nc		0.1	0.0
hard clams	nc		0.2	0.0	mixed sponge species	nc		0.1	0.0
rubbery bryozoan	nc		0.2	0.0	Tubularia, spp.	nc		0.1	0.0
surf clam	1	0.0	0.2	0.0	purple sea urchin	1	0.0	0.1	0.0
red bearded sponge	nc		0.1	0.0	water jelly	1	0.0	0.1	0.0
ghost shrimp	1	0.0	0.1	0.0					
Total	3,335		866.0		Total	12,236		663.2	

Note: nc= not counted

Table 5.22. Bluefish indices of abundance, 1984-2014.

Using September and October length data, the geometric mean catch per tow was calculated for two age groups of bluefish: age-0 and all fish age 1 and older. Age-0 was defined as bluefish less than 30 cm fork length.

Year	Fall			
	age 0 count / tow	age 0 kg / tow	ages 1+ count / tow	ages 1+ kg / tow
1984	20.34	2.51	1.61	2.03
1985	11.27	1.64	4.16	6.25
1986	8.05	1.13	3.77	5.96
1987	9.01	0.88	3.11	4.85
1988	10.73	1.59	2.20	4.43
1989	21.07	3.17	1.92	3.80
1990	12.82	2.09	6.14	8.92
1991	22.57	2.75	5.59	8.49
1992	9.23	1.27	8.44	14.88
1993	11.61	1.96	3.34	7.11
1994	24.85	2.54	3.07	6.09
1995	16.85	2.48	4.07	5.32
1996	13.85	2.27	2.34	4.09
1997	31.26	2.56	2.35	3.68
1998	25.89	2.08	1.65	2.70
1999	39.19	5.43	0.86	1.61
2000	14.67	2.97	2.18	3.75
2001	19.04	2.11	2.62	3.87
2002	12.35	2.25	3.63	4.81
2003	16.85	3.16	2.16	3.31
2004	13.30	2.39	10.38	13.96
2005	12.10	2.39	2.65	5.04
2006	12.43	1.49	2.14	2.74
2007	23.98	4.14	2.44	4.22
2008	6.14	0.82	4.52	8.18
2009	11.65	1.16	3.18	5.09
2010	-	-	-	-
2011	8.21	1.34	1.40	2.36
2012	13.11	1.86	0.97	1.67
2013	7.86	0.87	0.96	1.82
2014	16.53	2.22	0.88	1.47
84-13				
mean	15.87	2.18	3.24	5.21

Table 5.23. Scup indices-at-age, 1984-2014.

Spring (May and June) and fall (September and October) catch and age data were used to determine the geometric mean indices-at-age¹. The spring and fall age keys were used to expand length frequencies to age frequencies and then the spring and fall overall indices were proportioned by the percentage of fish in each age. The 0-10+ index represents the overall index (sum of ages 0-10+), and the adult 2+ index is provided as the sum of ages 2-10+ index. Fish older than age 9 were included in the age 10+ index².

Spring (May-June)														
Year	0-10+	2+	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10+	
1984	2.797	2.308	0	0.489	1.311	0.577	0.307	0.074	0.004	0.002	0	0	0.034	
1985	5.648	2.707	0	2.941	2.002	0.327	0.244	0.047	0.025	0.050	0	0.004	0.008	
1986	7.230	2.785	0	4.444	1.651	0.988	0.137	0.003	0.003	0.003	0	0	0.003	
1987	2.186	1.758	0	0.428	1.646	0.071	0.034	0.007	0	0	0	0	0	
1988	2.061	0.893	0	1.168	0.309	0.502	0.054	0.026	0	0	0	0	0.003	
1989	6.249	0.615	0	5.634	0.563	0.034	0.016	0.000	0.001	0.001	0	0	0	
1990	4.867	2.345	0	2.521	2.098	0.206	0.037	0.005	0	0	0	0	0	
1991	7.046	2.795	0	4.251	1.436	1.258	0.086	0.012	0.002	0	0	0	0	
1992	1.749	1.360	0	0.389	1.212	0.093	0.052	0.002	0	0.002	0	0	0	
1993	2.530	2.492	0	0.038	2.286	0.189	0.006	0.006	0.002	0.002	0	0	0	
1994	3.892	3.093	0	0.799	2.038	0.931	0.100	0.015	0.003	0.007	0	0	0	
1995	13.587	0.645	0	12.943	0.387	0.199	0.052	0.003	0.003	0	0	0	0	
1996	7.766	2.562	0	5.204	2.477	0.074	0.004	0.006	0.002	0	0	0	0	
1997	7.558	4.394	0	3.164	2.610	1.679	0.063	0.009	0.023	0.005	0.005	0	0	
1998	10.826	0.761	0	10.065	0.578	0.115	0.063	0.005	0	0	0	0	0	
1999	4.732	2.021	0	2.711	1.755	0.162	0.074	0.030	0	0	0	0	0	
2000	146.224	21.711	0	124.513	17.184	4.237	0.195	0.064	0.030	0	0	0	0	
2001	22.486	20.837	0	1.649	18.988	1.575	0.252	0.018	0.003	0.001	0	0	0	
2002	257.914	208.764	0	49.150	66.611	123.248	17.437	1.294	0.099	0.035	0.040	0	0	
2003	13.116	12.980	0	0.136	4.047	3.284	4.964	0.608	0.069	0.005	0.005	0	0	
2004	26.915	26.902	0	0.014	3.965	8.956	4.904	8.207	0.764	0.079	0.018	0.009	0	
2005	8.483	7.325	0	1.157	1.278	1.055	1.511	1.269	1.944	0.223	0.045	0	0	
2006	59.052	40.570	0	18.482	23.719	5.629	2.072	2.557	3.160	2.897	0.529	0.007	0	
2007	32.802	25.288	0	7.514	15.865	5.845	1.489	0.548	0.536	0.541	0.385	0.073	0.007	
2008	92.100	75.143	0	16.957	40.620	27.815	4.936	0.911	0.158	0.303	0.236	0.148	0.016	
2009	104.454	72.840	0	31.614	28.228	28.413	12.491	2.498	0.613	0.215	0.134	0.250	0.000	
2010	68.138	67.717	0	0.421	24.265	21.998	14.002	6.019	1.187	0.118	0.058	0.041	0.029	
2011	36.112	33.985	0	2.127	3.285	11.378	9.812	4.116	3.391	1.421	0.248	0.071	0.263	
2012	114.410	65.371	0	49.039	25.925	11.982	9.231	9.567	4.671	2.755	0.871	0.144	0.226	
2013	57.922	53.309	0	4.613	29.415	8.721	3.150	4.982	4.451	1.545	0.758	0.169	0.117	
2014	60.483	45.822	0	14.661	10.635	23.833	5.069	1.504	2.323	1.486	0.608	0.319	0.045	
84-13	Mean	37.695	25.543	0.000	12.153	10.925	9.051	2.926	1.430	0.705	0.340	0.111	0.030	0.023

Fall (Sept-Oct)														
Year	0-10+	2+	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10+	
1984	10.721	1.692	7.986	1.043	0.783	0.519	0.280	0.092	0.018	0	0	0	0	
1985	30.972	1.277	24.914	4.781	0.425	0.587	0.190	0.044	0.030	0.002	0	0	0	
1986	25.761	2.519	12.863	10.379	2.277	0.219	0.013	0.005	0.005	0	0	0	0	
1987	18.544	2.063	12.468	4.013	1.405	0.579	0.058	0.009	0.009	0.004	0	0	0	
1988	39.699	2.092	31.687	5.920	1.818	0.242	0.032	0	0	0	0	0	0	
1989	65.087	1.596	40.920	22.571	1.501	0.083	0.012	0	0	0	0	0	0	
1990	69.477	7.396	54.350	7.731	6.946	0.398	0.034	0.005	0.008	0	0	0.005	0	
1991	311.570	2.953	291.568	17.050	1.759	1.040	0.147	0.008	0	0	0	0	0	
1992	83.731	6.244	50.971	26.516	5.540	0.398	0.287	0.013	0.007	0	0	0	0	
1993	77.057	1.165	74.061	1.831	1.019	0.121	0.012	0.010	0	0	0.003	0	0	
1994	92.523	0.657	90.778	1.088	0.457	0.185	0.012	0.003	0	0	0	0	0	
1995	59.136	0.150	32.465	26.521	0.144	0.006	0	0	0	0	0	0	0	
1996	61.459	1.400	51.497	8.562	1.365	0.029	0	0.005	0	0	0	0	0	
1997	41.276	0.809	31.791	8.677	0.630	0.172	0.008	0	0	0	0	0	0	
1998	103.272	0.628	90.404	12.240	0.537	0.069	0.022	0	0	0	0	0	0	
1999	537.683	8.574	498.180	30.930	8.349	0.195	0.019	0.011	0	0	0	0	0	
2000	521.103	9.265	250.391	261.446	8.323	0.794	0.140	0.008	0	0	0	0	0	
2001	177.641	20.239	140.506	16.897	18.421	1.607	0.186	0.025	0	0	0	0	0	
2002	348.703	41.179	259.902	47.623	23.321	16.812	0.665	0.325	0.048	0	0.007	0	0	
2003	152.227	83.963	52.910	15.354	32.065	22.394	26.440	2.493	0.539	0.016	0.016	0	0	
2004	291.458	36.277	251.052	4.129	8.338	15.082	5.978	6.245	0.534	0.072	0.008	0.021	0	
2005	424.063	18.183	373.318	32.562	8.144	2.437	4.015	1.505	1.689	0.332	0.060	0	0	
2006	116.755	13.575	52.164	51.016	9.525	2.341	0.257	0.351	0.377	0.681	0.044	0	0	
2007	475.295	37.346	319.893	118.056	29.335	5.929	0.896	0.226	0.302	0.313	0.313	0.033	0	
2008	303.256	24.478	243.679	35.099	11.921	7.044	3.556	1.055	0.502	0.137	0.124	0.140	0	
2009	139.380	31.506	67.486	40.388	20.786	6.934	2.615	0.735	0.214	0.131	0.068	0.022	0	
2010	-	-	-	-	-	-	-	-	-	-	-	-	-	
2011	198.226	40.786	119.032	38.409	8.157	14.894	9.669	3.922	3.225	0.586	0.167	0.025	0.140	
2012	223.522	15.983	153.235	54.305	9.963	2.846	2.063	0.567	0.137	0.323	0.076	0.007	0	
2013	40.683	16.235	17.744	6.704	9.187	4.069	0.807	1.058	0.746	0.237	0.090	0.031	0.011	
2014	182.583	14.003	144.702	23.878	4.325	6.505	1.188	0.426	0.808	0.476	0.193	0.051	0.032	
84-13	Mean	173.803	14.835	127.525	31.443	8.015	3.725	2.014	0.646	0.289	0.098	0.034	0.010	0.005

(1) In 1984, 1985, 2003, 2004, 2006, 2008, 2010, 2011, and 2014 less than the number of scheduled tows were conducted in some months (Table 5.4).
 (2) Fish in the age 10+ group include: 6 fish taken 1984-1988, 8 fish taken 2002-2010, 81 taken in 2011, 28 taken in 2012, 26 taken in 2013, and 15 taken in 2014. The oldest scup aged were four 14-year-old fish taken in 1985 and 2013 and 2014 (2).

Table 5.24. Age frequency of striped bass taken in spring, 1984-2014.

Ages were derived from trawl survey length data using the average of Hudson River and Chesapeake Bay von Bertalanffy parameters.

Age	Year																														
	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
1	0	0	0	0	0	0	0	0	0	2	0	0	3	0	0	0	1	0	2	1	1	0	0	2	11	5	0	1	11	0	0
2	0	0	0	2	1	5	28	11	4	3	6	98	12	36	119	41	113	47	150	30	15	220	3	46	20	84	3	2	46	49	4
3	0	0	0	0	1	3	8	7	8	7	10	26	97	116	122	87	20	41	76	38	38	54	25	109	15	54	7	2	13	33	94
4	0	0	0	2	4	1	2	3	13	16	20	8	37	40	68	42	22	15	48	23	18	59	15	44	48	130	17	29	13	21	73
5	0	0	0	2	0	1	1	5	5	14	18	7	14	17	28	95	22	28	45	39	21	33	22	44	41	64	24	50	19	12	20
6	0	0	0	2	1	1	3	0	1	8	8	6	7	14	20	46	32	36	52	41	22	28	11	28	11	34	11	44	12	16	6
7	0	0	0	0	0	0	0	2	0	7	1	1	8	9	3	17	12	13	25	23	14	16	10	9	7	10	6	29	5	10	1
8	0	0	0	0	0	0	0	1	2	1	1	3	2	4	1	4	4	2	12	5	3	9	4	3	3	1	2	7	3	15	5
9	0	0	0	0	0	0	0	2	1	1	1	0	3	2	1	0	1	2	3	7	2	1	3	1	1	0	0	1	2	1	1
10	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	1	2	0	1	0	0	0	3	3	2	0	0	0	0	2	1
11	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	0	0	0	0	0	0	1	0
12	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	0	0	0	8	7	11	43	32	34	59	65	150	184	238	362	334	229	184	414	207	135	421	97	289	159	382	70	166	125	160	205

Note: number of fish taken but not measured = one in 1984, one in 1988, two in 1990.

Table 5.25. Striped bass indices-at-age, 1984-2014.

Spring length data was converted to ages using the average of Hudson River and Chesapeake Bay von Bertalanffy parameters (Vic Crecco, pers comm). Indices-at-age were then determined by apportioning the spring indices (from Table 5.18) by the percentage of fish in each age.

Year	Index	Spring											
		Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12
1984	0.02	0	0	0	0	0	0	0	0	0	0	0	0
1985	0.00	0	0	0	0	0	0	0	0	0	0	0	0
1986	0.00	0	0	0	0	0	0	0	0	0	0	0	0
1987	0.05	0	0.0125	0	0.0125	0.0125	0.0125	0	0	0	0	0	0
1988	0.04	0	0.0057	0.0057	0.0229	0	0.0057	0	0	0	0	0	0
1989	0.06	0	0.0273	0.0164	0.0055	0.0055	0.0055	0	0	0	0	0	0
1990	0.16	0	0.1042	0.0298	0.0074	0.0037	0.0112	0	0	0	0.0037	0	0
1991	0.15	0	0.0516	0.0328	0.0141	0.0234	0	0.0094	0.0047	0.0094	0.0047	0	0
1992	0.22	0	0.0259	0.0518	0.0841	0.0324	0.0065	0	0.0129	0.0065	0	0	0
1993	0.27	0.0093	0.0140	0.0326	0.0745	0.0652	0.0372	0.0326	0.0047	0.0047	0	0	0
1994	0.30	0	0.0277	0.0462	0.0923	0.0831	0.0369	0.0046	0.0046	0.0046	0	0	0
1995	0.59	0	0.3855	0.1023	0.0315	0.0275	0.0236	0.0039	0.0118	0	0.0039	0	0
1996	0.63	0.0103	0.0411	0.3321	0.1267	0.0479	0.0240	0.0274	0.0068	0.0103	0	0.0034	0
1997	0.85	0	0.1286	0.4143	0.1429	0.0607	0.0500	0.0321	0.0143	0.0071	0	0	0
1998	0.97	0	0.3189	0.3269	0.1822	0.0750	0.0536	0.0080	0.0027	0.0027	0	0	0
1999	1.10	0	0.1346	0.2857	0.1379	0.3119	0.1510	0.0558	0.0131	0	0.0033	0.0033	0
2000	0.84	0.0037	0.4163	0.0737	0.0811	0.0811	0.1179	0.0442	0.0147	0.0037	0.0074	0	0
2001	0.61	0	0.1558	0.1359	0.0497	0.0928	0.1193	0.0431	0.0066	0.0066	0	0	0
2002	1.30	0.0063	0.4722	0.2392	0.1511	0.1416	0.1637	0.0787	0.0378	0.0094	0.0031	0	0
2003	0.87	0.0042	0.1267	0.1605	0.0971	0.1647	0.1732	0.0971	0.0211	0.0296	0	0	0
2004	0.56	0.0042	0.0627	0.1588	0.0752	0.0878	0.0919	0.0585	0.0125	0.0084	0	0.0042	0
2005	1.17	0	0.6100	0.1497	0.1636	0.0915	0.0776	0.0444	0.0250	0.0028	0	0.0028	0
2006	0.61	0	0.0189	0.1572	0.0943	0.1384	0.0692	0.0629	0.0252	0.0189	0.0189	0.0063	0
2007	1.02	0.0071	0.1629	0.3860	0.1558	0.1558	0.0992	0.0319	0.0106	0.0035	0.0106	0	0
2008	0.57	0.0394	0.0717	0.0538	0.1721	0.1470	0.0394	0.0251	0.0108	0.0036	0.0072	0	0
2009	0.60	0.0078	0.1316	0.0846	0.2037	0.1003	0.0533	0.0157	0.0016	0	0	0	0
2010	0.40	0	0.0169	0.0394	0.0958	0.1352	0.0620	0.0338	0.0113	0	0	0	0
2011	0.48	0.0029	0.0058	0.0058	0.0839	0.1446	0.1272	0.0839	0.0202	0.0029	0	0	0.0029
2012	0.43	0.0381	0.1595	0.0451	0.0451	0.0659	0.0416	0.0173	0.0104	0.0069	0	0.0035	0
2013	0.67	0	0.2052	0.1382	0.0879	0.0503	0.0670	0.0419	0.0628	0.0042	0.0084	0.0042	0
2014	0.41	0	0.0080	0.1880	0.1460	0.0400	0.0120	0.0020	0.0100	0.0020	0.0020	0	0
84-13													
mean		0.0044	0.1298	0.1168	0.0830	0.0782	0.0573	0.0284	0.0115	0.0049	0.0024	0.0009	0.0001

Table 5.26. Summer flounder indices-at-age, 1984-2014.

Year and season specific age keys obtained from the NMFS spring and fall surveys were used to convert LISTs length frequencies to ages. Starting in 2000 LISTs ageing data (60 cm and over) were added to the age key to supplement the older age groups. Indices-at-age were determined for each season by apportioning the spring and fall overall indices (from Table 5.18 and Table 5.19) by the percentage of fish in each age.

Spring														
Year	0-11	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12
1984	0.6291	0	0.3236	0.2610	0.0445	0	0	0	0	0	0	0	0	0
1985	0.4410	0	0.0166	0.3168	0.0489	0.0587	0	0	0	0	0	0	0	0
1986	0.9510	0	0.7700	0.0892	0.0742	0.0126	0.0050	0	0	0	0	0	0	0
1987	1.0572	0	0.9515	0.0793	0.0202	0.0036	0.0026	0	0	0	0	0	0	0
1988	0.4986	0	0.2317	0.2232	0.0352	0.0085	0	0	0	0	0	0	0	0
1989	0.1016	0	0.0111	0.0550	0.0191	0.0164	0	0	0	0	0	0	0	0
1990	0.3475	0	0.3053	0.0201	0.0156	0.0065	0	0	0	0	0	0	0	0
1991	0.6391	0	0.3892	0.2059	0.0205	0.0235	0	0	0	0	0	0	0	0
1992	0.5546	0	0.3182	0.1906	0.0229	0	0.0229	0	0	0	0	0	0	0
1993	0.5074	0	0.3216	0.1504	0.0101	0.0152	0.0101	0	0	0	0	0	0	0
1994	0.8601	0	0.4959	0.3136	0.0324	0	0	0	0.0182	0	0	0	0	0
1995	0.2796	0	0.2023	0.0608	0.0110	0	0	0	0.0055	0	0	0	0	0
1996	0.9609	0	0.6216	0.2370	0.0868	0	0.0052	0	0.0103	0	0	0	0	0
1997	0.9991	0	0.4481	0.4461	0.0740	0.0121	0.0134	0.0054	0	0	0	0	0	0
1998	1.3067	0	0.0734	0.5952	0.4693	0.1167	0.0324	0.0197	0	0	0	0	0	0
1999	1.4401	0	0.3263	0.5563	0.3521	0.1110	0.0696	0.0248	0	0	0	0	0	0
2000	1.7898	0	0.3805	0.7853	0.4240	0.0538	0.1316	0.0092	0	0.0054	0	0	0	0
2001	1.7468	0	0.8408	0.3395	0.3653	0.1073	0.0488	0.0333	0.0067	0.0051	0	0	0	0
2002	3.1851	0	1.0571	1.2637	0.4646	0.2233	0.0930	0.0362	0.0236	0.0145	0.0091	0	0	0
2003	3.4211	0	1.6080	1.0159	0.3949	0.2316	0.0851	0.0462	0.0327	0.0025	0.0042	0	0	0
2004	1.8381	0	0.2592	0.8180	0.4100	0.1878	0.0338	0.0817	0.0302	0.0145	0.0029	0	0	0
2005	0.8038	0	0.2523	0.2641	0.1495	0.0334	0.0364	0.0393	0.0196	0.0046	0.0046	0	0	0
2006	0.6129	0	0.0383	0.3597	0.0676	0.0654	0.0337	0.0263	0.0168	0.0051	0	0	0	0
2007	2.5073	0	1.1569	0.2053	0.5595	0.3163	0.1150	0.0888	0.0428	0.0152	0.0065	0.0010	0	0
2008	1.6145	0	0.6008	0.2912	0.2374	0.2633	0.1165	0.0622	0.0236	0.0033	0.0054	0.0054	0.0054	0
2009	1.9295	0	0.7772	0.3770	0.2905	0.1804	0.1949	0.0700	0.0258	0.0101	0.0036	0	0	0
2010	2.6878	0	1.8671	0.2805	0.2113	0.1439	0.0944	0.0416	0.0244	0.0142	0.0052	0.0052	0	0
2011	3.8479	0	1.0024	1.0839	0.8014	0.3820	0.3159	0.1098	0.0628	0.0580	0.0171	0.0146	0	0
2012	3.0620	0	0.4684	0.6283	0.9746	0.6346	0.2044	0.0754	0.0333	0.0224	0.0050	0.0113	0.0043	0
2013	3.2359	0	0.8843	0.6681	0.6637	0.6734	0.2047	0.0818	0.0201	0.0184	0.0041	0.0044	0.0129	0
2014	2.9996	0	0.9709	0.7062	0.4847	0.4325	0.2977	0.0465	0.0369	0.0126	0.0072	0.0022	0.0022	0.0022
84-13														
Mean	1.4619	0.0000	0.5667	0.4060	0.2450	0.1294	0.0623	0.0284	0.0132	0.0064	0.0023	0.0014	0.0008	0.0000
Fall														
Year	0-11	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8	Age 9	Age 10	Age 11	Age 12
1984	0.9888	0	0.5648	0.3269	0.0713	0.0140	0.0042	0.0042	0.0034	0	0	0	0	0
1985	1.1931	0.2453	0.3605	0.4984	0.0804	0	0.0085	0	0	0	0	0	0	0
1986	1.7157	0.1738	1.1902	0.2681	0.0817	0.0019	0	0	0	0	0	0	0	0
1987	1.3963	0.0749	1.0573	0.2309	0.0305	0.0027	0	0	0	0	0	0	0	0
1988	1.4159	0.0150	0.8739	0.4782	0.0366	0.0122	0	0	0	0	0	0	0	0
1989	0.1363	0	0.0227	0.1051	0.0085	0	0	0	0	0	0	0	0	0
1990	0.8678	0.0321	0.6720	0.1214	0.0339	0.0042	0.0042	0	0	0	0	0	0	0
1991	1.2557	0.0363	0.8141	0.3457	0.0432	0.0082	0.0041	0.0041	0	0	0	0	0	0
1992	1.0178	0.0131	0.5685	0.3578	0.0561	0.0134	0.0089	0	0	0	0	0	0	0
1993	1.1113	0.0842	0.8371	0.1490	0.0362	0.0029	0	0.0019	0	0	0	0	0	0
1994	0.5517	0.1325	0.3008	0.0957	0.0138	0.0089	0	0	0	0	0	0	0	0
1995	0.5408	0.0424	0.3812	0.1043	0.0090	0.0039	0	0	0	0	0	0	0	0
1996	2.1914	0.0840	1.0394	1.0276	0.0375	0.0029	0	0	0	0	0	0	0	0
1997	2.4980	0.0693	0.8494	1.2261	0.3016	0.0321	0.0099	0.0084	0.0012	0	0	0	0	0
1998	1.7153	0	0.3251	1.0456	0.2867	0.0392	0.0187	0	0	0	0	0	0	0
1999	2.6787	0.0482	0.8000	1.4412	0.2963	0.0823	0.0084	0.0023	0	0	0	0	0	0
2000	1.9134	0.1151	0.5117	0.8244	0.2971	0.1122	0.0433	0.0067	0	0.0029	0	0	0	0
2001	4.4181	0.0208	2.6891	1.1372	0.4342	0.1095	0.0153	0.0078	0	0.0042	0	0	0	0
2002	6.1211	0.4415	3.0870	1.9304	0.4769	0.1216	0.0429	0.0168	0.0040	0	0	0	0	0
2003	3.3879	0	1.4584	1.3192	0.4069	0.0873	0.0908	0.0164	0.0089	0	0	0	0	0
2004	1.9537	0.2545	0.3848	0.7551	0.4398	0.0804	0.0241	0.0150	0	0	0	0	0	0
2005	2.4099	0.0671	1.0930	0.7441	0.3554	0.0866	0.0316	0.0123	0.0166	0.0032	0	0	0	0
2006	1.3148	0.0976	0.2170	0.5915	0.2299	0.0957	0.0435	0.0214	0.0182	0	0	0	0	0
2007	1.8880	0.1295	0.5669	0.3869	0.4676	0.2012	0.0778	0.0408	0.0087	0.0043	0	0	0.0043	0
2008	3.0853	0.7816	0.4848	0.9581	0.4458	0.3256	0.0804	0.0090	0	0	0	0	0	0
2009	3.1169	0.4054	0.6606	0.8883	0.6241	0.3182	0.1330	0.0437	0.0244	0.0070	0.0122	0.0000	0.0000	0
2010	-	-	-	-	-	-	-	-	-	-	-	-	-	0
2011	2.5578	0.1173	0.6933	0.9333	0.5641	0.1232	0.0543	0.0275	0.0130	0.0130	0.0061	0.0052	0.0075	0
2012	3.7358	0.1633	0.4592	0.8283	1.4239	0.5848	0.1836	0.0631	0.0296	0	0	0	0	0
2013	3.0664	0.2181	0.5709	0.6080	0.8049	0.6328	0.1789	0.0291	0.0139	0.0016	0	0.0082	0	0
2014	1.7086	0.1231	0.4034	0.3945	0.3620	0.2825	0.0823	0.0294	0.0205	0.0078	0	0.0031	0	0
84-13														
Mean	2.0774	0.1332	0.8115	0.6802	0.2894	0.1072	0.0368	0.0114	0.0049	0.0012	0.0006	0.0005	0.0004	0.0000

note: 1984-1999 indices-at-age were run using a GT 60cm group in the age key.

Table 5.27. Tautog indices-at-age, 1984-2014.

Year and season specific age keys obtained from the LISTS spring and fall surveys were used to convert LISTS length frequencies to ages. Indices-at-age were then determined for each season by apportioning the spring and fall overall indices (from Table 5.18 and Table 5.19) by the percentage of fish in each age, and then summing the spring and fall indices-at-age. The age 1-20+ index is the sum of indices ages 1 – 20+. The age 20+ category includes 36 fish ranging from 20 to 30 years of age.

Year	Age										
	1 - 20+	1	2	3	4	5	6	7	8	9	10
1984	3.4693	0.0109	0.0816	0.1898	0.3030	0.4590	0.4955	0.2892	0.2851	0.3105	0.3532
1985	1.7968	0	0.0191	0.0936	0.1922	0.1667	0.1279	0.1836	0.3005	0.2021	0.0902
1986	1.7200	0.0015	0.0273	0.0933	0.0495	0.1037	0.2019	0.2409	0.2452	0.2864	0.1017
1987	1.2129	0.0242	0.0799	0.0592	0.0602	0.1003	0.1341	0.1908	0.1349	0.0957	0.0523
1988	0.9006	0.0031	0.0327	0.0466	0.0721	0.0447	0.0401	0.0755	0.1008	0.1641	0.0790
1989	1.2590	0	0.0426	0.0683	0.1370	0.0893	0.1154	0.1495	0.1600	0.1046	0.0817
1990	1.1615	0.0113	0.0840	0.1546	0.1122	0.1142	0.0493	0.0500	0.1247	0.0875	0.0622
1991	1.1468	0.0057	0.0235	0.0582	0.1189	0.1241	0.1487	0.0931	0.1253	0.1071	0.1067
1992	1.0254	0.0197	0.0490	0.0709	0.0412	0.0491	0.1229	0.1323	0.0849	0.0632	0.0636
1993	0.5694	0.0034	0.0211	0.0505	0.0313	0.0166	0.0605	0.0595	0.0423	0.0489	0.0522
1994	0.5839	0.0093	0.0362	0.0322	0.0684	0.0558	0.0551	0.0555	0.0799	0.0516	0.0312
1995	0.2529	0.0034	0.0091	0.0092	0.0297	0.0602	0.0269	0.0212	0.0346	0.0150	0.0219
1996	0.5627	0.0073	0.0518	0.0305	0.0086	0.0762	0.0452	0.0654	0.0712	0.0667	0.0608
1997	0.5079	0	0.0390	0.0675	0.0568	0.0574	0.0639	0.0491	0.0556	0.0486	0.0101
1998	0.6442	0	0.0425	0.0281	0.0701	0.0821	0.0876	0.0875	0.0848	0.0465	0.0575
1999	0.7614	0.0498	0.0792	0.0583	0.0666	0.1015	0.1379	0.0748	0.0843	0.0431	0.0203
2000	0.8004	0.0012	0.0466	0.0578	0.0829	0.0740	0.1402	0.1376	0.0897	0.0392	0.0467
2001	0.8946	0.0062	0.0304	0.0863	0.0830	0.1294	0.1197	0.1193	0.1058	0.0715	0.0454
2002	1.1666	0.0101	0.0247	0.0585	0.1012	0.1748	0.1972	0.1895	0.2091	0.0739	0.0419
2003	0.8978	0.0033	0.0124	0.0083	0.0598	0.1485	0.2385	0.1596	0.0893	0.0778	0.0185
2004	0.6934	0.0075	0.0205	0.0150	0.0361	0.0710	0.1930	0.1096	0.0494	0.0812	0.0440
2005	0.7596	0.0100	0.0367	0.0618	0.0261	0.0922	0.1437	0.1576	0.1064	0.0303	0.0268
2006	0.8405	0	0.0334	0.0345	0.1039	0.1274	0.1140	0.1196	0.1521	0.0620	0.0479
2007	0.6136	0.0038	0.0126	0.0167	0.0460	0.0478	0.0608	0.0919	0.0936	0.0966	0.0532
2008	0.7269	0.0066	0.0279	0.0428	0.0620	0.0848	0.1164	0.0708	0.0649	0.0831	0.0640
2009	0.4822	0.0150	0.0355	0.0074	0.0026	0.0394	0.0681	0.1013	0.0658	0.0319	0.0324
2010	0.2472	0	0.0053	0.0455	0.0093	0.0053	0.0315	0.0503	0.0294	0.0096	0.0093
2011	0.4456	0.0180	0.0401	0.0532	0.0303	0.0301	0.0612	0.0630	0.0415	0.0267	0.0167
2012	0.5809	0.027	0.1148	0.0919	0.0808	0.0635	0.0389	0.0384	0.0499	0.0489	0.0115
2013*	0.5779	0.0083	0.0575	0.0726	0.0668	0.0714	0.1041	0.0893	0.0441	0.023	0.0105
2014*	0.6960	0.0134	0.0911	0.0757	0.0804	0.0991	0.1237	0.0887	0.0475	0.0325	0.011
84-13											
Mean	0.8218	0.0088	0.0392	0.0543	0.0657	0.0828	0.1050	0.1044	0.1007	0.0754	0.0469

Year	Age									
	11	12	13	14	15	16	17	18	19	20+
1984	0.1261	0.2281	0.0933	0.0507	0.0449	0.0322	0.0469	0.0156	0.0006	0.0531
1985	0.1595	0.0982	0.0226	0.0994	0	0.0249	0.0039	0.0124	0	0
1986	0.1423	0.0863	0.0374	0.0522	0.0232	0.0071	0.0114	0.0003	0.0023	0.0061
1987	0.0607	0.0543	0.0479	0.0313	0.0246	0.0265	0.0105	0.0004	0.0048	0.0203
1988	0.0469	0.0395	0.0295	0.0225	0.0493	0.0086	0.0063	0.0055	0.0052	0.0286
1989	0.0569	0.0932	0.0430	0.0404	0.0348	0.0172	0.0067	0.0048	0	0.0136
1990	0.0979	0.0375	0.0567	0.0397	0.0221	0.0250	0.0088	0.0170	0.0035	0.0033
1991	0.0610	0.0258	0.0399	0.0361	0.0217	0.0005	0.0160	0.0117	0.0080	0.0148
1992	0.0599	0.0512	0.0440	0.0581	0.0236	0.0208	0.0167	0.0298	0.0167	0.0078
1993	0.0368	0.0351	0.0351	0.0129	0.0157	0.0152	0.0129	0.0097	0.0097	0
1994	0.0234	0.0238	0.0071	0.0118	0.0118	0.0096	0.0024	0.0047	0.0070	0.0071
1995	0.0036	0.0036	0.0073	0	0	0	0.0036	0	0	0.0036
1996	0.0230	0.0127	0.0103	0.0048	0.0099	0.0090	0.0086	0.0004	0.0001	0.0002
1997	0.0072	0.0119	0.0144	0.0048	0.0121	0.0071	0	0.0024	0	0
1998	0.0192	0.0164	0.0055	0.0055	0	0.0027	0.0055	0	0	0.0027
1999	0.0191	0.0090	0.0087	0.0029	0	0	0.0030	0.0029	0	0
2000	0.0213	0.0130	0.0123	0.0101	0.0084	0.0104	0.0023	0	0.0027	0.0040
2001	0.0407	0.0161	0.0152	0.0004	0.0053	0.0105	0.0036	0.0001	0.0026	0.0031
2002	0.0257	0.0185	0.0107	0.0070	0.0147	0.0039	0	0	0	0.0052
2003	0.0274	0.0088	0.0059	0.0184	0.0029	0.0124	0	0.0029	0	0.0031
2004	0.0204	0.0221	0.0119	0.0003	0.0028	0.0031	0.0026	0.0002	0	0.0027
2005	0.0347	0.0257	0.0039	0.0037	0	0	0	0	0	0
2006	0.0183	0.0200	0.0037	0	0.0037	0	0	0	0	0
2007	0.0294	0.0156	0.0194	0.0108	0.0019	0.0116	0	0.0019	0	0
2008	0.0322	0.0225	0.0228	0.0163	0.0098	0	0	0	0	0
2009	0.0343	0.0064	0.0091	0.0217	0.0070	0.0032	0.0011	0	0	0
2010	0.0192	0.0139	0.0048	0.0046	0.0046	0	0	0	0.0046	0
2011	0.0167	0.0161	0.0080	0.0080	0.0040	0.0000	0.0040	0.0080	0.0000	0.0000
2012	0	0.0077	0.0038	0	0.0038	0	0	0	0	0
2013*	0.0091	0.0056	0.004	0.0005	0.0111				0	0
2014*	0.0078	0.0152	0.002	0.0019	0.0027			0.0033	0	0
84-13										
Mean	0.0424	0.0346	0.0213	0.0192	0.0125	0.0090	0.0061	0.0045	0.0023	0.0060

* 2013 and 2014 - ageing not complete so used a 2010-2012 pooled age key

Table 5.28. Weakfish age 0 and age 1+ indices of abundance, 1984-2014.

Using spring (May, June) and fall (September, October) length data, the geometric mean catch per tow was calculated for three groups of weakfish: fall age-0, spring - all fish age 1 and older (1+), and fall - all fish age 1 and older (1+). Weakfish less than 30 cm fork length in the fall were defined as age-0.

Year	Fall		Fall		Spring	
	age 0 count / tow	age 0 kg / tow	ages 1+ count / tow	age 1+ kg / tow	ages 1+ count / tow	ages 1+ kg / tow
1984	1.00	0.14	0.53	0.84	0.02	0.15
1985	6.19	0.74	0.24	0.46	0.00	0.10
1986	13.16	0.91	0.24	0.51	0.10	0.33
1987	0.63	0.13	0.11	0.16	0.02	0.11
1988	3.49	0.30	0.06	0.13	0.05	0.17
1989	8.69	0.94	0.02	0.10	0.04	0.16
1990	5.56	0.56	0.08	0.13	0.07	0.13
1991	11.95	1.44	0.31	0.41	0.28	0.26
1992	3.05	0.31	0.18	0.24	0.12	0.22
1993	4.08	0.46	0.12	0.18	0.10	0.15
1994	11.19	1.23	0.06	0.13	0.04	0.12
1995	5.22	0.84	0.70	0.64	0.18	0.16
1996	15.23	1.49	0.56	0.52	0.19	0.19
1997	12.38	1.03	0.89	0.81	0.42	0.34
1998	5.02	0.76	0.28	0.36	0.37	0.41
1999	30.93	3.21	0.39	0.51	0.45	0.59
2000	63.31	3.34	0.30	0.32	0.18	0.28
2001	40.09	2.20	0.52	0.54	0.27	0.26
2002	41.35	2.85	0.16	0.26	0.16	0.26
2003	49.41	1.77	0.07	0.17	0.04	0.14
2004	58.98	2.99	0.21	0.25	0.15	0.16
2005	25.86	2.50	0.12	0.18	0.27	0.23
2006	1.05	0.20	0.29	0.30	0.14	0.22
2007	63.93	3.86	0.06	0.14	0.11	0.22
2008	9.03	1.17	0.08	0.14	0.05	0.12
2009	6.48	0.57	0.30	0.22	0.08	0.16
2010	-	-	-	-	0.02	0.12
2011	11.64	0.87	0.68	0.55	0.10	0.15
2012	21.96	1.47	0.73	0.69	0.62	0.56
2013	7.01	0.59	0.52	0.52	0.52	0.44
2014	41.53	2.27	0.08	0.12	0.17	0.23
84-13						
mean	18.55	1.34	0.30	0.36	0.17	0.23

**TABLES 5.30 - 5.65
LENGTH FREQUENCIES
LISTS**

Table 5.30. Alewife length frequencies, spring and fall, 1 cm intervals, 1989–2014.

From 1989 - 1990, lengths were recorded from the first three tows of each day; since 1991, lengths have been recorded from every tow.

length	Spring																									
	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
6	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
7	0	0	0	0	2	0	0	0	0	0	0	4	0	0	1	0	1	0	0	4	0	2	1	0	0	0
8	0	0	0	0	18	3	3	0	0	2	9	16	0	3	1	2	0	0	4	1	10	0	1	3	2	
9	0	0	2	0	15	9	6	1	6	0	6	21	32	1	18	6	16	0	0	4	6	10	0	3	7	
10	0	0	0	1	11	19	18	2	22	7	6	28	23	5	32	55	32	0	8	5	11	23	5	6	16	
11	0	0	5	4	10	44	11	2	64	11	20	52	14	6	27	87	26	29	13	32	10	9	22	8	11	
12	6	0	4	7	6	83	17	8	127	12	32	43	5	29	25	100	55	44	34	131	17	6	54	27	19	
13	1	0	4	4	47	122	48	16	63	44	42	99	4	70	11	83	61	15	38	193	24	12	48	98	18	
14	0	0	9	7	77	172	35	26	69	61	56	234	7	139	28	63	37	9	37	178	51	6	50	187	14	
15	3	0	8	5	68	140	54	32	56	51	120	334	6	157	25	33	50	49	85	86	101	8	59	123	12	
16	2	0	8	5	84	159	38	86	44	50	144	320	4	86	26	31	74	25	128	46	106	7	37	56	5	
17	5	4	4	16	63	108	32	203	28	34	330	85	5	82	21	33	73	78	161	47	142	5	7	27	10	
18	4	4	9	8	59	81	7	254	32	22	136	15	4	15	19	18	71	93	182	25	196	2	11	17	21	
19	6	7	7	2	37	33	7	180	9	11	99	20	3	6	26	42	59	86	122	49	215	7	11	24	22	
20	3	1	7	2	27	24	10	161	17	17	82	22	9	17	13	30	26	76	105	38	137	7	9	19	10	
21	1	0	3	1	13	17	14	107	34	22	72	27	12	28	22	50	21	40	71	21	53	18	9	18	28	
22	4	2	8	2	10	26	12	103	48	18	47	41	18	46	25	48	18	18	41	14	29	22	10	24	34	
23	5	1	8	6	3	12	12	76	44	16	47	90	36	63	40	36	7	5	28	16	13	12	16	27	39	
24	7	0	3	2	1	12	7	34	28	14	21	58	45	49	42	13	6	1	10	7	14	4	7	18	15	
25	3	2	1	0	3	5	2	9	9	2	11	11	23	12	29	11	3	1	3	0	11	2	4	11	4	
26	1	0	1	2	1	5	1	3	1	2	2	1	5	7	17	5	2	0	2	0	1	0	2	3	3	
27	2	0	1	0	0	1	0	0	0	0	0	1	2	1	2	2	1	0	0	0	0	0	0	1	0	
28	1	0	0	0	1	1	0	0	0	1	0	0	0	1	0	2	1	0	0	1	0	0	2	0		
29	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
31	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
32	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	56	21	93	74	556	1,076	334	1,304	701	395	1,275	1,515	274	820	452	749	642	569	1,068	901	1,138	172	364	698	291	

length	Fall																									
	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
6	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	-	0	0	0	0
7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
8	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	1	0	0	1	0	-	0	0	0	
9	0	0	0	0	3	1	0	0	1	0	0	1	6	1	1	0	1	0	3	2	0	-	1	0	0	
10	0	0	0	0	5	1	4	1	1	0	1	4	23	0	7	1	7	0	8	2	1	-	1	0		
11	0	0	0	0	27	30	5	5	6	1	3	5	59	0	33	6	14	0	22	1	2	-	9	0		
12	0	0	0	1	120	82	9	25	12	9	6	9	86	4	64	7	8	0	44	0	2	-	22	2		
13	0	0	3	0	88	84	14	21	21	7	9	17	72	0	4	12	17	0	87	5	10	-	14	3		
14	0	0	2	4	16	36	11	30	31	0	11	10	23	3	3	16	15	0	134	14	10	-	22	0		
15	0	0	1	8	21	31	0	9	53	0	5	8	24	3	5	28	15	2	118	4	8	-	28	2		
16	3	0	3	10	53	14	4	1	110	1	25	2	36	17	20	30	12	4	31	0	1	-	14	1		
17	2	0	0	12	25	33	1	2	194	4	34	0	27	8	19	12	3	0	8	3	1	-	19	2		
18	3	0	0	9	13	24	1	1	62	3	11	1	5	0	0	1	5	0	6	0	1	-	17	0		
19	0	0	0	2	1	11	0	0	0	1	4	1	0	1	0	0	0	0	7	1	0	-	1	0		
20	0	0	0	0	0	2	0	0	0	0	2	0	0	0	0	0	0	0	0	0	1	-	0	0		
21	0	0	0	0	3	1	1	0	0	1	2	0	0	0	0	0	0	0	0	0	0	-	0	0		
22	0	1	0	0	2	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	-	0	0		
23	0	0	0	0	0	1	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0	-	0	0		
24	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0		
25	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0		
26	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0		
27	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0		
Total	8	1	9	46	377	354	50	95	492	27	117	58	364	38	156	113	98	6	468	33	37	0	148	10		

Table 5.31. American shad length frequencies, spring and fall, 2.0 cm intervals (midpoint given), 1989-2014.

From 1989 - 1990, lengths were recorded from the first three tows of each day; since 1991, lengths have been recorded from every tow.

length	Spring																									
	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
7	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1
9	0	0	0	0	8	2	17	0	6	9	5	5	2	13	6	1	6	0	0	0	1	0	0	0	0	7
11	0	0	1	3	7	2	16	5	24	27	20	46	1	101	12	8	11	0	5	26	12	12	5	3	48	41
13	4	0	10	8	4	4	11	9	59	85	31	29	2	87	11	14	10	0	20	78	36	21	28	34	38	32
15	49	1	82	17	6	22	22	191	177	108	65	21	2	41	0	45	25	38	54	180	66	77	100	106	20	9
17	29	8	49	23	10	72	68	154	319	97	52	32	4	49	3	6	4	14	44	51	40	47	25	45	11	3
19	5	5	4	33	6	374	40	47	62	32	20	13	0	17	0	2	0	5	8	11	15	5	3	5	2	1
21	1	3	10	25	6	158	6	9	2	1	35	1	0	4	4	2	6	0	3	3	3	2	1	0	1	1
23	0	3	31	20	5	18	2	16	5	8	50	4	0	7	7	4	7	0	4	3	4	0	0	10	8	16
25	0	2	10	7	1	6	0	15	1	7	14	2	3	4	0	0	3	0	7	0	0	1	0	22	1	2
27	0	1	1	0	0	2	0	5	0	1	1	1	0	0	0	0	2	0	4	0	0	0	0	4	0	2
29	0	0	0	0	0	1	0	0	0	0	0	0	0	5	0	0	0	0	0	0	0	0	0	0	0	0
31	0	0	0	0	0	0	0	0	0	1	0	0	0	2	0	0	1	0	0	0	0	0	0	0	0	0
33	0	0	0	0	0	0	0	0	1	3	0	3	0	1	0	0	1	0	2	0	0	0	0	0	0	0
35	0	1	1	1	0	1	0	0	0	0	0	0	0	1	0	1	0	0	0	1	1	0	0	0	0	1
37	0	0	0	2	0	1	0	0	4	0	1	0	0	1	0	0	1	1	0	0	0	0	0	0	0	1
39	1	0	0	3	2	2	1	0	2	0	4	0	0	2	0	0	0	1	1	0	0	0	0	0	0	1
41	1	0	1	5	2	3	2	0	3	0	3	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0
43	0	0	1	4	2	1	0	0	1	1	6	0	0	2	0	0	0	0	0	0	0	0	0	1	0	0
45	1	0	1	7	2	3	2	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
47	0	0	0	2	0	1	2	0	1	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	2	1
49	0	0	0	2	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
51	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	91	24	202	163	61	675	189	452	669	378	313	157	14	337	43	83	79	60	152	353	178	165	162	231	142	120

length	Fall																									
	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
7	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	2	0	0	0	0	0	-	0	0	0	1
9	0	0	7	1	2	6	7	0	6	1	5	0	1	1	4	5	4	0	2	4	0	-	4	4	0	9
11	0	1	4	5	23	26	16	1	20	14	27	0	4	1	14	6	3	0	19	4	27	-	4	4	0	2
13	0	0	7	21	54	208	24	7	28	13	44	0	1	0	22	4	5	0	26	3	22	-	2	2	1	2
15	0	0	4	2	33	245	14	2	5	4	6	0	0	0	2	0	0	13	0	36	-	2	0	2	5	
17	0	0	22	7	10	20	2	0	12	64	13	2	5	11	15	77	3	1	2	0	3	-	6	2	8	0
19	32	34	93	41	53	57	84	0	67	290	130	16	47	199	121	155	23	6	5	6	42	-	35	5	31	9
21	129	143	22	102	466	229	335	15	99	123	251	104	34	44	80	21	46	0	8	28	88	-	42	52	32	9
23	30	27	0	30	394	197	83	19	12	0	179	39	3	0	6	0	14	1	8	7	25	-	14	21	5	1
25	0	0	0	1	24	50	3	4	0	0	17	0	1	0	0	1	0	0	0	0	0	-	0	0	0	2
27	0	0	0	3	2	7	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
29	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
31	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
33	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
35	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
37	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
39	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
41	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
43	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
45	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
47	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
49	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
51	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
Total	192	205	159	214	1,061	1,047	568	48	251	509	674	161	96	256	262	273	98	8	83	52	243	-	109	90	79	40

Table 5.37. Atlantic menhaden length frequency, spring and fall, 1 cm intervals, 1996-2014.

Menhaden are scheduled to be measured from every tow. However, the following numbers of menhaden were not measured: 5 juveniles and 4 adults in 1996, and 7 adults in 1997.

length	Spring																		
	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
8	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
9	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0
10	0	0	0	0	0	0	4	0	0	0	0	0	0	0	0	0	7	0	0
11	0	0	0	1	0	0	13	0	0	0	0	0	0	0	0	0	3	0	0
12	0	0	0	0	0	0	10	0	0	0	0	0	0	0	0	0	15	0	0
13	0	0	0	0	0	0	6	0	0	0	2	0	0	0	0	0	8	0	0
14	0	0	0	0	0	0	5	0	0	0	0	0	0	0	0	0	5	0	0
15	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	8	0	0
16	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1	0
17	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
18	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
19	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0
20	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
21	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0
22	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
23	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
24	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0
25	0	0	0	0	0	0	0	0	0	0	0	5	0	0	0	0	1	0	0
26	0	0	0	0	0	0	1	0	0	0	0	4	0	0	0	0	2	3	6
27	0	0	0	0	0	0	1	0	0	0	0	6	2	3	1	4	14	25	46
28	0	1	0	0	1	0	1	0	0	0	0	5	4	9	5	10	33	32	81
29	0	1	0	0	1	0	0	1	3	0	1	5	2	2	1	18	53	59	79
30	0	1	0	0	0	0	1	1	0	0	0	4	1	5	0	10	28	27	34
31	0	3	0	0	0	0	0	0	1	0	2	4	1	0	0	1	12	13	19
32	0	0	0	0	1	0	3	0	0	0	0	0	0	0	0	0	1	0	1
33	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0
34	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0
35	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0
Total	0	6	0	1	9	0	47	2	5	1	5	33	10	19	7	43	195	162	266

length	Fall																		
	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
4	0	0	0	0	0	0	0	1	0	0	0	0	0	0	-	0	0	0	0
5	0	0	0	0	0	0	0	2	0	0	0	1	0	0	-	0	0	0	0
6	0	0	0	0	0	0	0	17	1	0	0	24	0	0	-	0	1	1	0
7	1	0	0	20	12	0	2	32	26	0	1	39	2	0	-	0	0	0	0
8	0	1	18	51	73	0	6	22	178	11	0	32	2	2	-	0	0	0	0
9	0	11	53	152	128	0	8	9	135	22	0	12	6	0	-	0	0	0	0
10	1	5	120	471	125	1	9	1	143	19	0	34	3	3	-	0	1	0	2
11	0	6	49	337	51	25	14	1	47	13	2	51	2	4	-	0	0	0	1
12	0	11	44	25	35	30	10	1	18	9	8	24	1	5	-	6	0	4	5
13	0	0	20	2	15	16	14	4	1	1	1	49	0	4	-	7	1	5	0
14	0	2	0	0	6	7	20	2	0	3	2	7	0	3	-	9	0	4	0
15	0	0	0	0	2	4	24	0	0	1	0	1	1	5	-	6	1	1	0
16	0	0	0	0	2	0	8	0	0	2	1	1	4	4	-	3	0	1	0
17	0	0	0	0	3	0	12	0	0	0	0	0	3	0	-	0	1	0	0
18	0	0	0	0	0	0	17	0	0	0	0	0	0	1	-	0	2	0	0
19	0	0	0	0	0	0	16	0	0	0	0	0	0	1	-	0	2	0	0
20	0	0	0	1	0	0	2	0	0	0	0	0	0	0	-	0	2	0	0
21	0	0	0	1	0	0	1	0	0	1	0	0	0	0	-	0	1	0	0
22	0	0	0	0	0	0	1	0	0	0	0	0	0	0	-	0	0	0	0
23	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
24	0	0	0	1	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
25	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	1	7
26	0	0	0	0	0	0	1	0	0	0	0	3	0	0	-	0	7	2	2
27	2	0	0	0	0	0	1	0	0	1	0	21	9	4	-	4	27	6	68
28	3	1	0	3	0	0	2	0	3	4	0	35	2	7	-	18	68	13	164
29	23	17	0	6	1	0	18	5	10	21	2	31	1	1	-	48	66	12	132
30	30	25	0	28	3	0	29	8	44	54	2	18	0	5	-	30	35	14	63
31	11	17	1	42	7	1	39	8	65	43	2	7	0	2	-	4	11	5	2
32	2	6	1	27	12	0	27	3	51	21	1	2	0	0	-	2	0	1	9
33	0	1	0	19	4	2	25	2	10	5	0	0	0	0	-	0	0	0	0
34	0	0	0	1	4	0	9	1	7	2	1	0	0	0	-	0	0	0	0
35	0	0	0	0	1	0	5	0	1	1	0	0	0	0	-	0	0	0	0
Total	73	103	306	1,187	484	86	320	119	740	234	23	392	36	51	-	137	226	162	455

Table 5.40. Blueback herring length frequencies, spring and fall, 1 cm intervals, 1989-2014.*From 1989 - 1990, lengths were recorded from the first three tows of each day; since 1991, lengths have been recorded from every tow.*

length	Spring																										
	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	
6	0	0	0	0	0	0	1	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	
7	0	0	2	0	2	7	2	0	0	2	0	4	1	0	3	2	1	0	0	1	0	4	0	0	0	5	1
8	0	0	3	0	2	76	20	4	0	5	0	10	7	12	7	9	8	1	0	8	0	1	0	0	9	8	
9	0	0	2	0	3	114	11	5	21	15	0	14	5	9	23	23	14	8	1	11	7	4	3	3	9	3	
10	0	0	5	10	7	74	9	19	45	45	0	18	2	9	26	47	6	23	9	14	19	19	5	18	5	1	
11	0	0	3	4	9	41	9	10	258	48	0	28	1	6	11	39	10	2	3	12	25	38	9	12	8	2	
12	3	0	5	0	2	9	5	3	4	16	0	18	2	3	4	20	12	0	5	2	27	8	3	5	1	2	
13	0	0	0	4	0	13	5	2	0	2	0	12	1	1	1	12	3	1	3	4	17	10	6	1	1	0	
14	0	0	0	15	0	5	3	1	1	1	0	3	0	0	0	0	7	0	1	1	5	4	2	0	0	0	
15	0	0	1	27	1	3	4	7	0	0	1	2	0	4	0	0	8	1	2	2	9	1	0	0	0	0	
16	0	0	0	65	0	8	3	7	0	3	5	1	1	1	4	4	13	2	23	1	30	4	2	2	7	0	
17	0	0	1	11	3	9	1	10	4	0	5	3	10	7	4	4	11	2	37	7	64	2	12	2	5	6	
18	0	1	0	2	0	3	0	4	2	0	0	5	15	2	3	3	1	2	7	3	49	1	3	2	3	11	
19	0	0	0	0	1	2	4	3	2	0	0	0	3	0	0	3	2	1	3	2	17	2	1	0	1	4	
20	0	0	0	4	0	1	1	0	0	0	0	2	1	1	0	0	5	2	0	1	2	0	1	0	1	3	
21	2	1	2	0	0	1	1	3	0	0	0	1	3	0	0	3	2	3	2	0	1	1	0	0	7	2	
22	1	0	0	1	0	3	0	4	0	1	0	3	0	0	1	0	1	0	1	1	0	1	0	0	5	2	
23	0	0	3	2	0	3	2	3	1	0	0	5	0	1	0	1	0	0	1	1	0	1	0	0	0	0	
24	0	1	2	0	0	0	0	2	0	0	0	3	0	0	0	0	0	0	2	0	0	1	0	0	1	0	
25	0	0	0	1	0	1	1	1	0	0	0	1	0	0	2	0	0	1	1	0	0	0	0	0	0	0	
26	0	0	0	1	0	0	1	0	0	0	0	1	0	0	0	3	0	0	0	0	0	0	0	0	0	0	
27	0	0	0	0	0	0	0	2	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	6	3	29	147	30	373	83	90	338	140	11	136	52	56	89	173	104	49	101	71	272	102	47	45	68	47	

length	Fall																									
	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
5	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
6	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
7	0	0	0	0	0	0	5	0	2	0	0	0	0	0	0	1	0	0	0	0	0	-	0	0	0	0
8	0	0	0	0	0	0	33	0	2	0	0	0	0	0	0	0	0	0	1	0	0	-	0	0	0	0
9	0	0	0	0	0	0	21	3	2	2	1	0	0	0	0	0	0	0	1	0	2	-	0	0	0	0
10	0	0	0	0	0	1	3	0	8	1	0	1	0	0	0	0	0	0	0	0	0	-	0	0	0	0
11	0	0	0	0	3	13	4	0	3	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
12	0	0	3	9	8	227	14	0	12	1	1	0	7	0	0	2	0	0	0	0	0	-	0	0	0	1
13	38	1	4	11	24	225	48	0	117	18	0	0	36	2	0	15	2	2	0	0	0	-	0	1	0	1
14	77	0	1	6	18	247	40	1	111	28	1	0	117	7	0	17	3	8	1	1	3	-	4	0	0	2
15	24	0	0	1	20	94	3	3	34	16	0	3	52	3	4	6	2	4	14	2	5	-	9	0	0	3
16	0	0	0	0	2	14	0	0	0	5	2	1	10	0	4	0	0	0	31	0	2	-	9	0	0	1
17	0	0	0	0	0	2	0	0	0	1	1	2	2	0	1	0	0	0	7	0	1	-	3	0	0	2
18	1	0	0	0	0	1	0	0	0	0	0	1	3	0	0	0	0	0	0	5	-	0	0	0	0	0
19	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
20	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	-	0	0	0	0
21	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	-	0	0	0	0
22	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
23	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
24	0	1	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
25	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
Total	140	2	9	27	76	827	172	7	292	72	8	8	227	12	9	42	8	14	55	3	18	0	25	1	0	10

Table 5.46. Fourspot flounder length frequencies, spring and fall, 2 cm intervals (midpoint given), 1989, 1990, 1996-2014.

Fourspot lengths were recorded from the first three tows of each day.

length	Spring																				
	1989	1990	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
13	2	0	0	0	0	1	0	1	0	0	0	0	1	0	0	0	1	0	1	0	0
15	5	2	0	0	5	5	0	0	3	0	3	0	0	0	0	0	0	0	0	0	0
17	21	8	1	3	8	12	1	2	17	2	13	0	0	6	0	0	6	2	5	1	1
19	19	19	8	16	14	61	22	5	89	8	8	0	6	7	7	4	2	1	24	2	6
21	17	42	31	60	13	28	26	4	99	6	4	1	18	11	9	10	3	10	42	11	5
23	11	341	198	161	16	32	239	42	33	8	4	14	24	9	17	6	5	45	56	20	9
25	56	528	279	353	105	72	422	181	84	124	26	71	29	44	39	37	33	157	258	185	64
27	103	225	208	456	209	97	256	300	199	228	82	75	33	105	81	91	55	150	441	209	172
29	120	139	193	392	233	81	201	245	191	187	129	64	44	170	108	127	55	107	461	189	179
31	89	60	117	192	137	66	139	153	175	163	178	68	61	121	94	90	69	93	303	139	107
33	51	27	54	76	60	60	81	45	89	88	113	52	36	52	70	51	36	49	92	100	78
35	8	33	15	22	16	25	39	11	26	47	35	31	13	43	34	31	24	27	31	27	29
37	2	12	6	3	4	7	12	8	7	12	5	11	4	9	11	7	9	9	4	16	8
39	0	4	3	0	2	1	1	2	3	6	2	3	1	7	2	0	4	5	0	0	0
41	0	0	0	0	0	0	0	0	0	0	0	2	1	0	0	1	0	0	0	0	1
43	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	1	0	0
45	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	504	1,440	1,113	1,734	822	548	1,439	999	1,015	879	602	394	271	585	472	455	302	655	1,719	899	659

length	Fall																				
	1989	1990	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
5	0	0	0	0	0	0	0	1	0	0	0	0	2	0	0	1	-	0	0	0	0
7	0	1	0	1	4	0	0	1	0	0	1	0	2	0	0	1	-	1	0	1	1
9	5	0	0	23	19	0	2	2	0	4	1	0	2	1	1	7	-	4	0	0	3
11	9	4	2	46	27	5	4	17	5	2	12	4	5	0	7	16	-	17	3	1	11
13	10	15	5	68	22	24	6	25	3	3	9	9	13	2	8	59	-	28	4	11	26
15	6	17	35	55	21	42	5	15	9	0	13	17	4	5	11	45	-	22	13	10	47
17	0	0	42	16	3	16	1	0	3	0	1	26	3	2	16	20	-	4	12	2	49
19	0	0	22	0	0	4	1	0	1	0	0	2	0	0	7	6	-	0	0	4	5
21	0	0	0	2	2	3	2	0	2	0	1	0	0	1	0	0	-	0	0	1	0
23	1	2	9	2	5	0	17	1	5	0	0	0	1	1	0	1	-	0	0	0	1
25	0	3	42	7	16	5	58	3	7	3	4	1	0	6	1	2	-	2	3	0	1
27	0	7	41	10	22	4	77	5	13	7	6	5	0	7	1	6	-	1	9	2	4
29	0	3	24	5	22	5	54	10	18	11	13	5	0	20	6	8	-	1	11	2	4
31	0	1	20	3	6	3	25	1	18	4	30	6	0	12	5	6	-	1	6	2	8
33	0	0	6	1	1	1	7	1	13	7	19	2	1	3	1	11	-	3	6	0	0
35	0	0	4	0	1	0	5	0	6	5	6	7	0	4	4	1	-	2	2	2	1
37	0	0	0	0	0	0	2	1	3	0	2	0	0	0	0	1	-	1	0	0	0
39	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	-	0	0	0	0
Total	31	53	252	239	171	112	266	83	106	46	118	85	33	64	68	192	-	87	69	38	161

Table 5.48. Horseshoe crab length frequencies by sex, spring, 1 cm intervals, 1998-2014.
Horseshoe crabs were measured (prosomal width) from every tow.

Sex	length	1998*	Spring																
			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	
F	13		1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
F	14		1	3	0	1	2	0	1	0	0	0	0	0	0	0	0	0	0
F	15		0	0	0	1	1	0	0	0	0	1	0	0	0	0	0	0	0
F	16		1	0	0	3	2	1	1	0	0	1	0	0	0	1	0	2	2
F	17		1	0	2	2	1	4	1	0	1	1	0	0	0	1	1	0	0
F	18		2	1	0	3	2	4	0	0	2	1	1	0	0	0	2	3	3
F	19		4	1	2	2	5	5	0	0	3	4	1	0	0	2	0	5	5
F	20		5	2	0	7	1	2	3	0	3	2	0	0	1	2	0	4	4
F	21		8	2	1	8	6	2	1	0	3	8	1	0	3	5	4	5	5
F	22		8	6	4	13	10	7	2	0	10	4	6	0	3	3	2	3	3
F	23		14	15	18	19	22	17	3	2	9	14	4	3	4	9	7	14	14
F	24		15	7	15	32	29	25	5	4	15	11	12	6	3	15	19	13	13
F	25		15	10	23	25	22	20	8	5	11	16	10	9	9	14	19	11	11
F	26		23	13	28	26	22	23	3	2	16	12	10	4	16	14	17	26	26
F	27		15	9	18	18	18	18	8	4	10	9	9	5	18	11	8	22	22
F	28		8	6	9	6	7	4	2	2	5	4	10	3	8	10	13	9	9
F	29		3	0	3	4	4	4	0	3	5	1	3	4	1	3	2	3	3
F	30		1	0	3	2	0	0	3	2	0	2	1	1	4	0	1	1	1
F	31		0	0	0	0	4	0	0	0	0	1	1	0	0	0	0	0	0
F	32		0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	1
M	14		0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0
M	15		0	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0
M	16		0	0	0	2	5	2	0	1	2	0	0	2	0	0	0	0	0
M	17		5	2	4	7	9	9	0	0	3	2	3	0	1	5	0	1	1
M	18		11	8	12	19	24	21	2	0	17	10	3	2	5	7	6	9	9
M	19		22	13	32	42	25	33	3	0	19	12	10	7	7	8	16	17	17
M	20		15	16	30	20	33	31	7	0	21	10	11	7	15	13	10	13	13
M	21		18	5	13	14	16	10	1	0	6	12	5	3	3	9	6	6	6
M	22		4	5	7	6	7	6	2	0	4	2	1	1	4	5	3	1	1
M	23		1	0	3	1	4	2	1	0	0	1	1	0	0	0	2	1	1
M	24		2	1	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0
M	25		0	0	0	0	0	1	2	0	0	0	0	0	0	0	0	3	3
M	26		0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0
M	27		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
M	28		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
M	29		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
M	30		0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
U	22		1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total			51	204	125	228	285	285	251	60	25	166	141	104	57	105	138	138	173

Table 5.49. Horseshoe crab length frequencies by sex, fall, 1 cm intervals, 1998-2014.
Horseshoe crabs were measured (prosomal width) from every tow.

Sex	length	Fall																
		1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
F	13	0	0	2	0	0	0	3	0	1	0	0	0	-	0	0	0	0
F	14	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
F	15	0	0	0	0	2	0	0	0	0	0	0	0	-	0	0	0	0
F	16	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
F	17	1	1	0	0	2	1	0	1	1	0	1	0	-	0	0	0	0
F	18	0	2	0	1	0	1	1	1	0	0	0	0	-	0	0	0	0
F	19	3	2	2	2	0	1	0	0	1	0	1	1	-	0	0	0	2
F	20	5	1	1	4	4	2	3	0	2	0	0	2	-	0	0	0	0
F	21	3	2	2	3	1	4	6	3	1	1	1	0	-	0	0	0	1
F	22	3	8	13	13	10	3	9	4	1	2	6	6	-	6	0	2	2
F	23	8	15	15	12	8	8	13	10	7	7	6	14	-	6	2	3	4
F	24	7	19	30	27	21	9	24	10	6	17	14	22	-	18	10	12	8
F	25	17	12	20	31	33	13	19	6	12	26	17	17	-	19	9	11	11
F	26	19	23	33	31	18	9	29	12	10	22	15	24	-	25	16	27	10
F	27	14	7	21	22	18	7	22	8	3	17	11	28	-	16	5	15	10
F	28	2	4	10	8	13	6	15	5	4	8	11	22	-	11	3	10	6
F	29	2	3	2	5	2	3	8	2	0	4	1	5	-	2	4	2	3
F	30	0	1	1	2	0	2	1	2	0	2	0	2	-	0	1	2	0
F	31	0	1	0	0	1	0	0	2	0	0	0	1	-	0	0	0	1
F	32	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
F	33	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
F	34	0	0	0	0	0	1	0	0	0	0	0	0	-	0	0	0	0
M	11	0	0	0	1	0	0	0	0	0	0	0	0	-	0	0	0	0
M	12	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
M	13	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
M	14	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
M	15	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
M	16	0	0	2	1	5	3	0	0	1	1	0	0	-	1	0	0	0
M	17	6	5	7	6	3	5	11	0	1	3	1	2	-	3	0	1	1
M	18	12	14	28	18	14	15	21	3	9	3	9	18	-	13	4	2	5
M	19	10	20	39	27	31	11	39	13	4	12	21	14	-	9	4	6	13
M	20	20	23	35	32	22	8	30	12	9	19	23	31	-	10	1	17	4
M	21	6	11	18	15	9	4	15	4	2	10	6	13	-	7	1	7	6
M	22	5	3	8	4	6	0	10	2	5	6	2	5	-	6	0	5	0
M	23	0	0	3	2	6	1	1	0	2	3	1	3	-	0	1	2	0
M	24	0	0	1	3	0	0	1	0	1	2	0	2	-	0	0	0	0
M	25	0	0	2	0	0	0	0	0	0	0	0	1	-	0	0	1	0
M	26	2	0	0	3	0	0	0	0	1	0	0	1	-	0	0	0	0
M	27	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
M	28	0	0	0	0	0	0	0	1	0	0	0	0	-	0	0	0	0
M	29	0	0	0	1	0	0	0	0	0	0	0	0	-	0	0	0	0
Total		145	177	295	274	229	117	281	101	83	165	148	234	-	152	61	41	29

Table 5.51. Long-finned squid length frequencies, fall, 1 cm intervals, 1986-1990, 1992-2014.*Length frequencies of squid taken from the first three tows of each day.*

length	Fall																												
	1986	1987	1988	1989	1990	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	
0	0	0	0	5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0	
1	0	13	0	12	0	0	0	0	0	0	0	0	0	0	0	4	0	0	3	12	0	14	-	0	0	11	0		
2	0	31	0	1	0	49	0	9	25	24	6	20	29	2	0	11	0	1	10	74	9	33	90	-	12	10	67	6	
3	0	126	59	112	74	266	914	80	156	57	125	115	104	53	36	80	90	170	91	107	20	87	343	-	80	101	51	25	
4	0	320	212	468	278	1,507	2,336	477	460	598	491	642	362	384	230	261	886	693	763	249	420	294	939	-	618	469	127	517	
5	0	892	826	743	830	2,906	3,502	1,332	1,223	1,371	1,091	1,888	1,214	1,215	663	695	2,225	1,757	1,539	587	1,367	417	2,332	-	1,417	705	273	1443	
6	3	1,019	1,165	677	836	5,015	4,358	1,803	1,896	1,869	1,278	2,737	1,782	1,842	923	1,067	3,185	2,705	2,337	913	2,780	604	2,894	-	1,405	731	426	1814	
7	13	817	722	446	469	5,210	4,331	2,152	2,254	2,751	1,169	3,412	2,390	2,204	996	1,193	2,566	2,759	2,552	917	3,822	780	2,746	-	1,315	698	550	1560	
8	135	654	333	283	220	3,110	3,811	2,225	2,080	2,224	935	2,939	1,808	1,797	839	929	1,885	1,787	2,006	611	3,549	908	1,791	-	840	638	570	1394	
9	16	692	146	108	129	1,594	2,913	2,486	2,124	1,853	570	1,993	1,829	1,081	616	488	1,785	907	1,283	385	2,119	777	1,131	-	670	584	418	1366	
10	13	503	65	58	42	894	1,772	2,055	1,540	1,264	446	1,216	1,332	695	528	354	861	626	970	204	1,974	480	808	-	637	399	306	1198	
11	0	310	62	70	39	737	1,178	1,607	905	698	291	675	780	556	264	214	215	392	541	183	1,379	332	326	-	343	359	178	862	
12	0	165	21	38	24	284	737	843	387	579	153	368	423	380	154	145	58	144	307	85	728	193	222	-	211	232	123	574	
13	0	82	24	34	17	242	408	415	159	297	126	328	277	247	132	87	2	96	194	31	447	103	108	-	139	148	62	315	
14	0	77	9	17	6	40	278	329	110	160	44	199	235	204	68	53	1	103	64	26	253	47	41	-	40	97	53	253	
15	0	31	11	17	3	18	185	181	77	83	31	103	133	128	66	13	2	48	44	9	150	18	27	-	86	64	14	213	
16	0	4	11	13	2	0	53	99	33	46	15	90	111	73	32	10	0	43	30	8	159	7	14	-	18	35	2	106	
17	0	14	0	10	4	0	73	75	15	16	13	23	120	101	8	6	0	1	24	17	103	5	2	-	7	8	6	50	
18	0	1	23	6	1	0	20	31	2	6	10	16	82	34	3	0	0	8	2	11	82	74	0	1	-	25	12	4	53
19	0	1	0	0	0	0	3	12	0	1	0	1	34	9	2	4	0	1	1	11	2	0	0	-	0	7	0	37	
20	0	13	0	5	1	0	2	7	0	0	1	1	22	3	2	1	0	4	2	1	3	0	0	-	0	1	0	21	
21	0	15	0	4	0	0	0	3	0	0	0	0	22	9	1	0	0	0	0	0	1	0	0	-	0	5	2	6	
22	0	2	0	3	1	0	0	11	0	6	0	1	17	0	0	0	0	0	1	0	0	0	0	-	0	2	1	0	
23	0	0	0	3	0	0	0	2	1	0	0	0	4	0	0	0	0	0	1	0	0	0	0	-	1	0	0	0	
24	0	1	0	3	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	5	0	0	-	0	0	0	0	
25	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	-	0	0	0	0	
26	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0	
27	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0	
28	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0	
29	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0	
30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	1	0	0	
Total	180	5,783	3,689	3,136	2,976	21,872	26,877	16,233	13,446	13,903	6,795	16,767	13,111	11,018	5,563	5,615	13,761	12,245	12,765	4,441	19,364	5,085	13,829	-	7,864	5,306	3,244	11,813	

Table 5.61. Windowpane flounder length frequencies, spring, 1 cm intervals, 1989, 1990, 1994-2014.
Lengths were recorded from the first three tows of each day.

length	Spring																							
	1989	1990	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	
4	0	0	1	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	1	0	0	0	
5	4	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	2	0	0	2	
6	0	0	0	0	0	2	0	2	5	1	1	10	2	0	0	1	0	4	4	9	0	0	10	
7	0	0	0	0	1	4	2	4	17	2	7	22	3	0	0	7	3	8	9	9	5	0	7	
8	0	2	4	1	3	5	4	3	27	7	6	23	6	0	0	31	5	17	10	20	19	10	41	
9	0	40	16	3	2	9	5	2	11	10	21	20	11	0	0	18	6	10	13	24	16	4	31	
10	25	66	67	12	34	15	7	8	17	13	12	11	19	7	2	4	11	23	8	10	10	16	24	
11	69	96	169	86	79	37	19	20	5	29	8	3	24	12	1	4	11	8	7	11	10	20	8	
12	89	74	305	148	162	76	60	40	3	23	10	7	25	16	7	8	17	4	20	2	0	16	10	
13	337	53	362	259	288	136	131	37	10	29	5	9	58	25	12	22	13	6	72	9	3	8	15	
14	430	66	232	189	381	309	200	45	11	26	8	13	100	22	34	28	44	17	93	7	7	10	18	
15	414	124	152	180	487	362	211	96	24	43	15	13	101	23	96	42	60	51	37	107	15	32	19	15
16	305	180	126	89	310	606	177	123	27	55	12	15	72	37	36	107	119	62	117	19	64	16	21	
17	174	212	209	70	331	754	130	165	23	73	9	15	65	22	48	129	137	97	166	23	81	17	26	
18	78	178	372	99	339	588	165	160	32	94	24	23	56	4	45	132	116	90	104	58	133	20	37	
19	65	132	357	139	548	440	260	194	26	78	19	26	45	16	20	110	101	75	124	58	155	30	37	
20	174	144	289	143	604	366	362	386	75	89	15	31	60	13	24	130	76	51	76	47	135	40	71	
21	216	116	217	85	567	429	461	357	136	95	22	45	32	22	24	186	122	50	88	66	97	62	75	
22	299	143	139	82	401	438	311	301	166	232	45	50	42	29	27	246	155	63	172	75	97	121	102	
23	319	108	163	57	409	368	229	217	138	290	110	92	39	42	28	181	216	92	198	107	117	140	170	
24	270	103	147	54	280	323	227	217	125	245	141	123	66	36	41	158	132	84	199	122	128	166	229	
25	177	87	183	54	236	231	188	206	121	208	133	111	109	47	31	162	118	82	155	134	121	142	228	
26	189	103	184	70	235	191	178	136	106	126	114	76	100	52	52	186	103	67	161	120	118	138	175	
27	138	79	138	56	187	222	162	161	91	88	69	88	86	49	37	104	100	60	148	103	102	86	145	
28	148	38	70	44	117	145	138	97	56	83	62	68	71	29	38	100	111	45	103	69	100	55	111	
29	78	26	68	24	97	98	67	53	47	59	41	37	48	24	24	65	52	30	146	42	70	41	56	
30	99	35	42	27	66	75	58	42	37	39	42	35	51	20	14	33	46	24	51	24	45	27	27	
31	50	20	25	12	31	23	34	39	12	25	19	22	32	13	8	14	22	11	67	25	33	12	21	
32	8	15	13	4	25	12	13	26	16	21	17	9	16	5	2	23	19	6	21	7	7	6	11	
33	16	3	2	9	5	8	6	3	8	15	7	2	10	1	3	2	5	1	33	14	13	8	5	
34	0	5	5	0	4	1	1	1	2	5	4	4	9	3	0	4	5	2	20	11	11	4	9	
35	0	4	5	1	3	0	3	4	5	10	2	4	5	0	0	3	3	3	11	1	4	2	5	
36	0	4	2	2	1	1	0	0	1	2	0	5	0	2	0	0	1	0	0	0	1	0	1	
37	0	0	0	1	0	0	3	1	1	2	2	1	1	0	0	0	0	0	8	0	0	0	0	
38	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	
39	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	
40	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
41	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
42	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	4,171	2,256	4,064	2,001	6,234	6,274	3,812	3,147	1,381	2,118	1,002	1,015	1,365	571	600	2,258	1,920	1,129	2,511	1,244	1,734	1,236	1,744	

Table 5.62. Windowpane flounder length frequencies, fall, 1 cm intervals, 1989, 1990, 1994-2014.
Lengths were recorded from the first three tows of each day.

length	Fall																						
	1989	1990	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
6	1	0	1	0	0	0	0	0	3	1	0	0	3	0	0	0	0	1	-	0	0	0	0
7	5	0	5	0	6	0	1	0	0	0	0	2	0	0	0	0	4	-	1	0	0	0	0
8	8	3	18	5	24	15	1	0	6	9	0	5	11	14	5	4	0	15	-	4	2	2	1
9	25	2	28	6	70	17	2	2	2	2	0	21	15	49	2	6	2	15	-	2	3	1	4
10	18	11	78	10	165	50	2	4	3	9	1	20	22	67	1	14	5	17	-	9	6	7	9
11	15	9	60	22	227	75	31	11	7	14	0	13	27	111	5	18	3	24	-	19	1	7	13
12	16	12	50	15	270	107	33	6	9	9	1	6	16	155	2	26	15	29	-	31	5	6	7
13	23	6	30	10	285	173	47	3	11	9	6	0	14	145	8	44	43	19	-	19	10	10	14
14	33	14	11	13	306	154	48	5	23	6	0	4	8	109	3	36	58	27	-	36	14	10	14
15	58	23	23	9	250	110	39	6	18	3	5	8	3	62	2	37	38	25	-	43	18	11	10
16	140	38	15	16	181	60	34	3	11	3	5	9	3	33	0	30	28	31	-	41	19	13	24
17	188	44	35	26	112	78	33	11	30	7	14	4	9	12	7	21	20	35	-	72	37	13	19
18	91	53	47	48	101	119	54	11	15	12	8	11	2	8	19	19	16	47	-	70	19	19	28
19	46	46	49	47	145	179	95	44	29	6	10	7	11	20	32	26	10	45	-	52	44	31	12
20	49	28	39	48	131	213	96	67	30	13	9	6	18	30	39	39	31	24	-	41	50	29	18
21	21	11	23	24	125	165	69	38	52	18	9	11	35	50	25	36	40	28	-	35	87	23	27
22	14	14	16	19	65	123	37	18	28	22	21	2	25	48	25	42	25	26	-	51	58	28	34
23	3	10	20	6	67	63	32	12	37	30	39	6	10	14	12	32	27	20	-	47	79	30	43
24	9	4	7	9	25	49	13	11	33	19	39	11	15	13	9	19	32	23	-	40	45	15	55
25	4	3	6	3	22	28	9	6	18	19	25	14	8	10	10	6	9	9	-	16	24	29	50
26	2	0	8	3	19	29	9	4	16	9	10	18	4	3	4	8	16	6	-	18	22	17	29
27	6	2	3	1	11	17	8	3	5	11	12	17	4	5	3	4	5	4	-	7	14	16	21
28	2	1	4	1	3	12	1	1	4	5	6	9	2	3	3	3	2	7	-	9	1	13	7
29	2	2	0	1	2	17	0	1	6	3	1	4	2	3	1	3	2	1	-	2	0	2	4
30	2	1	2	1	0	5	0	0	1	2	2	2	0	1	1	0	0	0	-	3	1	2	2
31	0	0	0	0	0	0	0	0	0	1	0	3	1	2	0	0	2	1	-	0	0	1	1
32	1	0	0	1	0	0	0	0	0	0	0	2	0	1	0	0	0	1	-	0	1	0	0
33	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0
Total	782	337	578	344	2,613	1,858	694	267	397	242	223	215	268	968	218	473	429	484	-	668	560	335	446

THIS PAGE INTENTIONALLY LEFT BLANK

**FIGURES 5.1 - 5.19
LISTS**





Figure 5.1. Trawl Survey site grid. Each sampling site is 1x2 nmi (nautical miles). A four-digit number identifies the site: the first two digits are the row numbers (corresponding to minutes of latitude) and the last two digits are the column numbers (corresponding to two nautical miles in length on the longitudinal axis). Examples: site 1428 near Guilford and 0028 near Mattituck. (Note: The sites in column 16 are approximately 2x1 nmi. The grid was drawn on the Eastern and Western Long Island Sound 80,000:1 nautical charts, which overlap by the area in column 16.)


Figure 5.2. April 2014 sites selected and sampled. The red outlined rectangles are the sites selected for the cruise and the blue dots are the sites sampled. Samples collected from a different site than published in the "Notice to Fishermen" are noted in table below map.



Persistent unfavorable weather conditions forced nine sites to be moved this month:


Sample	Site Sampled	Sampled Strata	Site Selected	Selected Strata	Sample	Site Sampled	Sampled Strata	Site Selected	Selected Strata
SP2014030	1019	T2	0818	T2	SP2014037	0724	T4	0125	T4
SP2014033	0922	M3	0024	M3	SP2014038	1427	T1	1423	T1
SP2014034	0521	M4	0119	M4	SP2014039	1332	S1	5823	S1
SP2014035	0622	M4	0120	M4	SP2014040	1333	S1	5824	S1
SP2014036	0623	M4	0121	M4					

Figure 5.3. May 2014 sites selected and sampled. *The red outlined rectangles are the sites selected for the cruise and the blue dots are the sites sampled. Samples collected from a different site than published in the “Notice to Fishermen” are noted in table below map.*




Sample	Site Sampled	Sampled Strata	Site Selected	Selected Strata	Reason Moved
No sites were moved during this cruise.					

Figure 5.4. June 2014 sites selected and sampled. The red outlined rectangles are the sites selected for the cruise and the blue dots are the sites sampled. Samples collected from a different site than published in the “Notice to Fishermen” are noted in table below map.




Sample	Site Sampled	Sampled Strata	Site Selected	Selected Strata	Reason Moved
SP2013083	1133	S4	0830	S4	sampled different site (same strata) for EPA

Figure 5.5. September 2014 sites selected and sampled. The red outlined rectangles are the sites selected for the cruise and the blue dots are the sites sampled. Samples collected from a different site than published in the “Notice to Fishermen” are noted in table below map.



Sample	Site Sampled	Sampled Strata	Site Selected	Selected Strata	Reason Moved
No sites were moved during this cruise.					

Figure 5.6. October 2014 sites selected and sampled. The red outlined rectangles are the sites selected for the cruise and the blue dots are the sites sampled. Samples collected from a different site than published in the “Notice to Fishermen” are noted in table below map.



Sample	Site Sampled	Sampled Strata	Site Selected	Selected Strata	Reason Moved
One site was not sampled during this cruise.					

Figure 5.7. Number of finfish species observed annually, 1984-2014. *Note: there was no October sampling in 2006 and there was no June, September or October sampling in 2010. Average number of finfish species caught per year is 57.6 for the time-series. See Table 5.4 for details on number of tows completed each year.*






Figure 5.8. Plots of abundance indices for: black sea bass, bluefish (total, age 0, and ages 1+), butterfish, cunner, and dogfish (smooth and spiny).



Legend:
■ = count / tow
▲ = kg / tow
 ---- = mean count / tow


Figure 5.9. Plots of abundance indices for: flounders (fourspot, summer, windowpane, winter, and winter ages 4+) and hakes (red, silver, and spotted).



Legend:


- = count / tow
- ▲ = kg / tow
- = mean count / tow

Figure 5.10. Plots of abundance indices for: herrings (alewife, Atlantic, and blueback), hogchoker, Northern kingfish, Atlantic menhaden, moonfish, and ocean pout.




Legend:
■ = count / tow
▲ = kg / tow
 ---- = mean count / tow

Figure 5.11. Plots of abundance indices for: fourbeard rockling, rough scad, longhorn sculpin, sea raven, and scup (all ages, age 0, and ages 2+).




Legend:
■ = count / tow
▲ = kg / tow
 ---- = mean count / tow

Figure 5.12. Plots of abundance indices for: searobins (striped and northern), shad (American and hickory), skates (clearnose, little, and winter), and spot.




Legend:
■ = count / tow
▲ = kg / tow
 ---- = mean count / tow

Figure 5.13 Plots of abundance indices for: striped bass, Atlantic sturgeon, tautog, and weakfish (all ages, age 0, and ages 1+).



Legend:
■ = count / tow
▲ = kg / tow
 ---- = mean count / tow

Figure 5.14. Plots of abundance and biomass indices for: crabs (lady, rock, and spider), horseshoe crab, American lobster, and long-finned squid.



Legend for bottom four graphs:

- = count / tow
- ▲ = kg / tow
- = mean count / tow

Figure 5.15. Mean number of finfish species per sample, spring and fall, 1984-2014. This index measures the diversity of species supported within the Sound's various habitats.




Figure 5.16. Open water forage abundance, 1992-2014. The geometric mean is calculated as the aggregate sample biomass per tow of 14 of the most common forage species sampled in the survey. This index measures the available food base which supports both resident and migratory species. The average since 1992 is 13.74 kg/tow (red line).




Figure 5.17. Geometric mean biomass of finfish and invertebrates per sample, spring and fall, 1992-2014.
 This index measures the diversity of species supported within the Sound's various habitats.





Figure 5.18: Percent of sampled winter flounder that were sexually mature by length group for female and male flounder captured in LISTS over five time periods, 1990-2013.





Figure 5.19. Trends in the number of cold temperate versus warm temperate species per sample captured in spring and fall LIS Trawl Surveys. See Appendix 2.5 for list of species included in analysis.



THIS PAGE INTENTIONALLY LEFT BLANK

**APPENDICES
LISTS**

Appendix 5.1. List of finfish species identified by A Study of Marine Recreational Fisheries in Connecticut (F54R) and other CT DEP Marine Fisheries Division programs. LISTS has collected one hundred-six finfish species from 1984-2014.

This appendix contains a list of 147 species identified (Bold type indicates new species) from all sampling programs conducted since 1984. Species are listed alphabetically by common name (AFS 2004). Sampling program abbreviations, survey time periods and gear type are as follows:

Survey Abbreviation	Survey Description	Time Period	Gear Type
CTR	CT River Creel Survey	1997-1998	bus stop creel survey mainstem of CT River
EPA	cooperative sampling in western LIS with EPA	1986-1990	used LISTS net
ESS (F54R)	Estuarine Seine Survey	1988 to present	7.6m (25 ft) beach seine
IS (F54R)	Inshore Survey of Juvenile Winter Flounder	1990-1994	beam trawls (also a little data from 1995-1996)
ISS (F54R-starting 2008)	Inshore Seine Surveys in CT & TH rivers	1979 to present	15.2m (50 ft) bag seine set by boat
LISTS (F54R)	Long Island Sound Trawl Survey	1984 to present	14m (50 ft) trawls with 2" codend mesh
MISC	misc sampling conducted on R/V Dempsey	various	various
NCA	"inshore" EPA NCA C2K sampling	2000	skiff trawls
NRRWS	sampling in western end of LIS, the "Narrows"	2000-2007	14m (50 ft) trawls with 2" codend mesh
SNFH (F54R)	Study of Nearshore Finfish Habitat	1995-1996	plankton net
SS (F54R)	Summer Survey	1991-1993, 1996	14m (50 ft) trawls with codend liner in LIS
TN	Trap Net Survey	1997-1998	trap nets in rivers

Common Name	Scientific Name	Survey
anchovy, bay	<i>Anchoa mitchilli</i>	LISTS;NRRWS;ESS;ISS;IS; SS;NCA;MISC
anchovy, striped	<i>Anchoa hepsetus</i>	LISTS; ESS; IS; SS
banded rudderfish	<i>Seriola zonata</i>	LISTS; ESS
bass, largemouth	<i>Micropterus salmoides</i>	ISS; TN;CTR
bass, rock	<i>Ambloplites rupestris</i>	ISS; TN;CTR
bass, smallmouth	<i>Micropterus dolomieu</i>	ISS; TN;CTR
bass, striped	<i>Morone saxatilis</i>	LISTS;NRRWS;ESS;ISS; SS;NCA;MISC;EPA;TN;CTR
bigeye	<i>Priacanthus arenatus</i>	LISTS; IS
bigeye, short	<i>Pristigenys alta</i>	LISTS
black sea bass	<i>Centropristes striata</i>	LISTS;NRRWS;ESS; IS; SS;NCA;MISC;EPA
blenny, feather	<i>Hypsoblennius hentz</i>	LISTS
bluefish	<i>Pomatomus saltatrix</i>	LISTS;NRRWS;ESS;ISS; SS; MISC;EPA; CTR
bluegill	<i>Lepomis macrochirus</i>	TN;CTR
bonefish	<i>Albula vulpes</i>	ISS
bonito, Atlantic	<i>Sarda sarda</i>	LISTS; EPA
bullhead, brown	<i>Ameiurus nebulosus</i>	ISS; NCA; TN;CTR
burrfish, striped	<i>Chilomycterus schoepfi</i>	LISTS; ESS
burrfish, web	<i>Chilomycterus antillarum</i>	ESS
butterfish	<i>Peprilus triacanthus</i>	LISTS;NRRWS;ESS;ISS;IS; SS;NCA;MISC;EPA
carp	<i>Cyprinus carpio</i>	ISS; NCA; TN;CTR
catfish, channel	<i>Ictalurus punctatus</i>	ISS; NCA; TN;CTR
catfish, white	<i>Ameiurus catus</i>	NCA; TN;CTR
cod, Atlantic	<i>Gadus morhua</i>	LISTS; SS
cornetfish, bluespotted	<i>Fistularia tabacaria</i>	LISTS; ESS; IS
cornetfish, red	<i>Fistularia petimba</i>	LISTS; IS
crappie, black	<i>Pomoxis nigromaculatus</i>	ISS; NCA; TN;CTR
crappie, white	<i>Pomoxis annularis</i>	TN;CTR
croaker, Atlantic	<i>Micropogonias undulatus</i>	LISTS; IS
cunner	<i>Tautoglabrus adspersus</i>	LISTS;NRRWS;ESS;ISS;IS; SS; MISC;EPA
cusck-eel, fawn	<i>Lepophidium profundorum</i>	LISTS
cusck-eel, striped	<i>Ophidion marginatum</i>	LISTS; SS
darter, tessellated	<i>Etheostoma olmstedii</i>	ISS
dogfish, smooth	<i>Mustelus canis</i>	LISTS;NRRWS;ESS; IS; SS; MISC;EPA
dogfish, spiny	<i>Squalus acanthius</i>	LISTS;NRRWS; MISC
drum, black	<i>Pogonias cromis</i>	LISTS
eel, American	<i>Anguilla rostrata</i>	LISTS;NRRWS;ESS;ISS;IS;SNFH;SS;NCA; EPA;TN;CTR
eel, conger	<i>Conger oceanicus</i>	LISTS; IS; SS
fallfish	<i>Semotilus corporalis</i>	ISS
filefish, orange	<i>Aluterus schoepfi</i>	LISTS; IS; SS
filefish, planehead	<i>Monacanthus hispidus</i>	LISTS; EPA
filefish, scrawled	<i>Aluterus scriptus</i>	IS
flounder, American plaice	<i>Hippoglossoides platessoide</i>	LISTS
flounder, fourspot	<i>Paralichthys oblongus</i>	LISTS;NRRWS; IS; SS; MISC;EPA

Appendix 5.1 cont.

Common Name	Scientific Name	Survey
flounder, smallmouth	<i>Etropus microstomus</i>	LISTS;NRRWS;ESS; IS; SS;NCA;MISC
flounder, summer	<i>Paralichthys dentatus</i>	LISTS;NRRWS;ESS;ISS;IS; SS;NCA;MISC;EPA;TN;CTR
flounder, windowpane	<i>Scophthalmus aquosus</i>	LISTS;NRRWS;ESS;ISS;IS; SS;NCA;MISC;EPA;TN;CTR
flounder, winter	<i>Pseudopleuronectes americanus</i>	LISTS;NRRWS;ESS;ISS;IS;SNFH;SS;NCA;MISC;EPA;TN;CT
flounder, yellowtail	<i>Pleuronectes ferrugineus</i>	LISTS; IS
glasseye snapper	<i>Priacanthus cruentatus</i>	LISTS
goatfish, dwarf	<i>Upeneus parvus</i>	LISTS
goatfish, red	<i>Mullus auratus</i>	LISTS
goby, code	<i>Gobiosoma robustum</i>	IS
goby, naked	<i>Gobiosoma boscii</i>	LISTS; ESS;ISS;IS
goldfish	<i>Carassius auratus</i>	CTR
goosefish	<i>Lophius americanus</i>	LISTS; IS; SS; MISC
grubby	<i>Myoxocephalus aeneus</i>	LISTS; ESS;ISS;IS;SNFH;SS; EPA
gunnel, banded	<i>Pholis fasciata</i>	ESS; IS
gunnel, rock	<i>Pholis gunnellus</i>	LISTS; ESS;ISS;IS;SNFH;SS
gurnard, flying	<i>Dactylopterus volitans</i>	ESS
haddock	<i>Melanogrammus aeglefinus</i>	LISTS; SS
hake, red	<i>Urophycis chuss</i>	LISTS;NRRWS; IS; SS; MISC;EPA
hake, silver	<i>Merluccius bilinearis</i>	LISTS;NRRWS; SS; MISC;EPA
hake, spotted	<i>Urophycis regia</i>	LISTS;NRRWS; ESS; IS; SS; MISC;EPA
harvestfish	<i>Peprilus paru</i>	LISTS
herring, Atlantic	<i>Clupea harengus</i>	LISTS;NRRWS; IS;SNFH;SS; MISC;EPA
herring, alewife	<i>Alosa pseudoharengus</i>	LISTS;NRRWS;ESS;ISS; SNFH;SS; MISC;EPA;TN;CTR
herring, blueback	<i>Alosa aestivalis</i>	LISTS;NRRWS;ESS;ISS;IS;SNFH;SS; EPA;TN;CTR
herring, round	<i>Trimeus teres</i>	LISTS; EPA
hogchoker	<i>Trinectes maculatus</i>	LISTS;NRRWS;ESS;ISS;IS; SS; MISC;EPA;TN
jack, blue runner	<i>Caranx crysos</i>	LISTS; EPA
jack, crevalle	<i>Caranx hippos</i>	LISTS;NRRWS; ESS; ISS; EPA
jack, yellow	<i>Caranx bartholomaei</i>	LISTS;NRRWS; ESS; IS; MISC;EPA
killifish, rainwater	<i>Lucania parva</i>	ESS
killifish, striped	<i>Fundulus majalis</i>	ESS; IS
kingfish, northern	<i>Menticirrhus saxatilis</i>	LISTS;NRRWS;ESS;ISS;IS; SS; EPA
lamprey, sea	<i>Petromyzon marinus</i>	LISTS; IS; TN
lizardfish, inshore	<i>Synodus foetens</i>	LISTS;NRRWS;ESS;ISS;IS; SS; MISC
lookdown	<i>Selene vomer</i>	LISTS; ISS
lumpfish	<i>Cyclopterus lumpus</i>	LISTS; IS;SNFH
mackerel, Atlantic	<i>Scomber scombrus</i>	LISTS; ISS; SS; EPA
mackerel, Spanish	<i>Scomberomorus maculatus</i>	LISTS; SS; EPA
menhaden, Atlantic	<i>Brevoortia tyrannus</i>	LISTS;NRRWS;ESS;ISS;IS;SNFH;SS;NCA;MISC;EPA
minnow, sheepshead	<i>Cyrinodon variegatus</i>	ESS;ISS
moonfish	<i>Selene setapinnis</i>	LISTS;NRRWS; SS; MISC;EPA
mullet, white	<i>Mugil curema</i>	LISTS;ESS;ISS
mummichog	<i>Fundulus heteroclitus</i>	ESS; IS
needlefish, Atlantic	<i>Strongylura marina</i>	ESS;ISS
ocean pout	<i>Macrozoarces americanus</i>	LISTS;NRRWS; MISC;EPA
oyster toadfish	<i>Opsanus tau</i>	LISTS;NRRWS;ESS;ISS;IS;SNFH;SS; EPA
perch, white	<i>Morone americana</i>	LISTS;NRRWS;ESS;ISS;IS;SNFH; NCA; TN;CTR
perch, yellow	<i>Perca flavescens</i>	ISS; SNFH; TN;CTR
perch, silver	<i>Bairdiella chrysoura</i>	LISTS
pickerel, chain	<i>Esox niger</i>	ISS; TN
pike, northern	<i>Esox lucius</i>	ISS; TN;CTR
pinfish	<i>Lagodon rhomboides</i>	LISTS
pipefish, northern	<i>Syngnathus fuscus</i>	LISTS;NRRWS;ESS;ISS;IS;SNFH;SS;NCA; EPA
pollock	<i>Pollachius virens</i>	LISTS;NRRWS; SNFH;SS; EPA
pompano, African	<i>Alectis ciliaris</i>	LISTS; ISS
puffer, northern	<i>Sphoeroides maculatus</i>	LISTS;NRRWS;ESS;ISS;IS; SS
pumpkinseed	<i>Lepomis gibbosus</i>	ESS;ISS; NCA; TN;CTR
radiated shanny	<i>Ulvaria subbifurcata</i>	SNFH
ray, bullnose	<i>Myliobatis freminvillei</i>	LISTS

Appendix 5.1 cont.

Common Name	Scientific Name	Survey
ray, roughtail stingray	<i>Dasyatis centroura</i>	LISTS
rockling, fourbeard	<i>Enchelyopus cimbrius</i>	LISTS;NRRWS; IS;SNFH;SS; MISC;EPA
salmon, Atlantic	<i>Salmo salar</i>	LISTS; TN
sand lance, American	<i>Ammodytes americanus</i>	LISTS; ESS; IS;SNFH;SS
sandbar (brown) shark	<i>Carcharhinus plumbeus</i>	LISTS
scad, bigeye	<i>Selar crumenophthalmus</i>	LISTS; SS; MISC
scad, mackerel	<i>Decapterus macarellus</i>	LISTS; SS
scad, rough	<i>Trachurus lathami</i>	LISTS;NRRWS; SS; MISC;EPA
scad, round	<i>Decapterus punctatus</i>	LISTS;NRRWS
sculpin, longhorn	<i>Myoxocephalus octodecemspinosus</i>	LISTS;NRRWS; ISS; SNFH; MISC
scup	<i>Stenotomus chrysops</i>	LISTS;NRRWS;ESS;ISS;IS; SS;NCA;MISC;EPA
sea raven	<i>Hemitripterus americanus</i>	LISTS; SNFH; MISC;EPA
seahorse, lined	<i>Hippocampus erectus</i>	LISTS; ESS; IS
searobin, northern	<i>Prionotus carolinus</i>	LISTS;NRRWS;ESS; IS;SNFH;SS; MISC;EPA
searobin, striped	<i>Prionotus evolans</i>	LISTS;NRRWS;ESS;ISS;IS; SS;NCA;MISC;EPA
seasnail	<i>Liparis atlanticus</i>	LISTS; SNFH
sennet, northern	<i>Sphyaena borealis</i>	LISTS; ESS
shad, American	<i>Alosa sapidissima</i>	LISTS;NRRWS;ESS;ISS; SS; MISC;EPA;TN;CTR
shad, gizzard	<i>Dorosoma cepedianum</i>	LISTS;NRRWS; ISS; TN
shad, hickory	<i>Alosa mediocris</i>	LISTS;NRRWS; ISS; SS; MISC;EPA; CTR
sharksucker	<i>Echeneis naucrates</i>	LISTS
shiner, golden	<i>Notemigonus crysoleucas</i>	ISS; TN
shiner, spottail	<i>Notropis hudsonius</i>	ISS; NCA; TN;CTR
silverside, Atlantic	<i>Menidia menidia</i>	LISTS;NRRWS;ESS;ISS;IS;SNFH;SS; MISC;EPA
silverside, inland	<i>Menidia beryllina</i>	SNFH
skate, barndoor	<i>Dipturus laevis</i>	LISTS
skate, clearnose	<i>Raja eglanteria</i>	LISTS;NRRWS; IS
skate, little	<i>Leucoraja erinacea</i>	LISTS;NRRWS;ESS; IS; SS;NCA;MISC;EPA; CTR
skate, winter	<i>Leucoraja ocellata</i>	LISTS;NRRWS; SS; MISC
smelt, rainbow	<i>Osmerus mordax</i>	LISTS; ESS; IS;SNFH;SS; TN;CTR
snapper, grey	<i>Lutjanus griseus</i>	ESS; IS
spot	<i>Leiostomus xanthurus</i>	LISTS;NRRWS; ISS;IS; SS; MISC;EPA
stargazer, northern	<i>Astroscopus guttatus</i>	LISTS; ESS
stickleback, four-spine	<i>Apeltes quadracus</i>	ESS; IS
stickleback, nine-spine	<i>Pungitius pungitius</i>	ESS; IS
stickleback, three-spine	<i>Gasterosteus aculeatus</i>	ESS; IS; TN
sturgeon, Atlantic	<i>Acipenser oxyrinchus</i>	LISTS
sucker, white	<i>Catostomus commersoni</i>	ISS; NCA; TN;CTR
tautog	<i>Tautoga onitis</i>	LISTS;NRRWS;ESS;ISS;IS; SS;NCA;MISC;EPA
tomcod, Atlantic	<i>Microgadus tomcod</i>	LISTS;NRRWS;ESS;ISS;IS;SNFH;SS; EPA; CTR
triggerfish, gray	<i>Balistes capriscus</i>	LISTS
trout, brook	<i>Salvelinus fontinalis</i>	TN;CTR
trout, brown	<i>Salmo trutta</i>	CTR
walleye	<i>Sander vitreus</i>	TN
weakfish	<i>Cynoscion regalis</i>	LISTS;NRRWS;ESS;ISS;IS; SS;NCA;MISC;EPA

Appendix 5.2 cont.

Common name	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	Total	
(number of tows)	200	246	316	320	320	320	297	200	160	240	240	200	200	200	200	200	200	200	200	200	199	200	120	200	120	200	78	172	200	200	199	6,547	
jack, crevalle	0	1	0	1	4	0	0	0	0	6	8	1	0	3	0	8	0	0	1	2	2	2	0	2	0	1	0	4	2	0	2	50	
jack, yellow	0	0	0	0	0	41	8	11	2	2	6	32	6	2	6	20	3	3	13	1	1	28	0	0	0	1	0	0	0	0	0	186	
kingfish, northern	0	0	0	0	0	1	1	4	2	10	7	25	6	7	15	6	2	2	1	1	5	4	0	4	3	7	0	34	59	14	51	271	
lamprey, sea	0	0	0	1	1	0	1	1	0	2	0	0	1	1	0	0	0	0	0	1	0	0	0	1	1	0	0	0	0	0	0	11	
lizardfish, inshore	0	0	0	0	0	2	0	0	0	0	1	0	0	2	1	7	1	21	1	0	0	1	4	2	10	2	0	43	0	0	30	128	
lobster, American	5,995	3,549	4,924	6,923	6,032	7,645	9,696	8,524	8,160	12,583	9,123	9,944	9,490	16,467	16,211	13,922	10,481	5,626	3,880	2,923	1,843	1,389	748	1,648	1,096	853	293	230	349	144	178	180,868	
lookdown	0	0	0	0	0	0	2	0	0	0	3	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	7	
lumpfish	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	
mackerel, Atlantic	68	17	20	29	45	376	46	2	4	17	11	1	5	8	13	21	2	0	5	8	0	37	0	9	0	5	0	0	0	0	2	751	
mackerel, Spanish	0	0	0	0	0	11	0	2	1	233	106	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	355
menhaden, Atlantic	161	304	718	600	335	623	407	348	1,115	298	411	318	88	116	306	1,187	492	86	366	799	746	235	28	426	47	69	7	181	426	234	723	12,198	
moonfish	7	226	23	7	142	60	10	24	62	6	149	33	921	287	1,188	645	1,817	225	424	133	182	356	361	979	689	2,575	0	640	262	868	2,200	15,501	
mullet, white	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	2	
ocean pout	26	3	14	14	30	58	39	42	18	66	42	30	26	15	13	17	18	6	13	14	18	3	5	12	9	22	6	27	14	0	0	619	
perch, silver	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	3	
perch, white	0	0	0	0	0	2	0	0	0	4	1	0	1	4	0	1	1	0	0	8	2	0	0	0	4	1	0	1	1	0	1	32	
pinfish	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	
pipefish, northern	1	0	1	0	3	0	0	0	5	21	2	2	0	1	0	2	4	4	2	6	2	4	3	2	0	2	4	4	1	2	1	79	
pollock	5	0	3	8	6	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	18	2	5	0	1	0	56	
pompano, African	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	
puffer, northern	1	2	6	0	3	2	2	5	1	28	4	1	3	1	28	14	4	8	6	3	5	5	0	8	0	5	0	9	47	3	10	214	
ray, bullnose ray	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	2
ray, roughtail stingray	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	1	1	0	1	0	0	1	0	0	1	1	0	0	0	8	
rockling, fourbeard	376	89	184	312	563	686	393	163	150	242	93	169	109	199	133	233	185	251	106	113	173	106	14	87	81	47	35	43	43	3	4	5,385	
rudderfish, banded	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	2	
salmon, Atlantic	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
sand lance, American	nc	nc	nc	nc	nc	nc	nc	nc	nc	3	25	95	0	2	4	178	4	4	3	19	70	6	0	30	7,495	1,227	13,061	9,535	2	7	12	31,782	
sand lance, (yoy-est)	nc	nc	nc	nc	nc	nc	nc	nc	nc	0	1,000	5	0	0	100	1,075	0	430	0	0	0	0	5,444	2	3,750	7,932	0	15,600	0	0	0	35,338	
scad, bigeye	0	0	0	0	15	63	1	1	0	0	3	0	2	1	1	21	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	108
scad, mackerel	0	0	0	0	0	0	1	2	6	0	4	1	3	0	1	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	1	21	
scad, rough	34	32	19	89	180	81	41	1	0	100	13	0	35	65	0	0	0	10	10	12	14	62	14	13	0	59	0	150	19	28	5	1,087	
scad, round	0	0	0	0	0	0	0	0	0	0	0	0	2	4	1	2	0	0	4	11	12	0	3	0	1	0	1	0	1	1	1	43	
sculpin, longhorn	14	82	51	32	107	107	263	139	31	11	7	5	7	4	2	2	14	5	3	5	5	0	0	3	2	2	1	9	1	1	0	915	
scup	8,806	18,054	16,449	9,761	12,566	37,642	21,193	45,790	13,646	32,218	38,456	13,985	16,087	9,582	23,742	101,095	101,464	58,325	100,481	26,926	61,521	52,642	28,829	75,681	53,560	46,991	7,157	34,457	53,119	24,961	45,705	1,190,891	
sea raven	57	59	70	88	52	34	44	19	4	1	1	2	2	3	30	9	19	7	11	3	7	3	0	5	0	5	6	3	5	0	1	550	
seahorse, lined	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
searobin, northern	585	2,267	546	280	605	381	357	609	313	951	878	1,317	672	579	360	547	2,014	1,594	2,123	1,632	784	265	630	691	809	2,012	1,128	803	3,642	1,934	2,584	33,894	
searobin, striped	1,434	2,295	2,035	1,482	2,086	2,211	2,353	865	857	1,491	1,298	682	1,008	819	1,321	1,690	3,129	2,061	2,394	2,235	1,308	757	366	755	612	1,507	141	1,630	2,973	2,724	2,544	49,063	
seasnail	0	0	0	0	1	0	8	0	0	0	0	0	0	0	0	0	4	0	0	4	0	2	0	0	0	0	0	0	0	0	0	0	19
sennet, northern	1	0	0	0	0	1	0	0	0	2	0	0	0	0	0	6	0	1	2	0	0	8	0	2	0	5	0	1	3	0	0	32	
shad, American	1,852	425	642	1,036	3,208	4,007	550	361	380	1,142	1,723	755	501	922	901	987	316	109	593	689	356	177	68	236	405	422	165	271	321	222	162	23,902	
shad, gizzard	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	1	2	0	1	0	0	0	0	1	0	0	0	9
shad, hickory	71	4	7	6	4	40	2	1	12	10	31	6	29	25	40	56	42	14	45	41	39	136	75	37	5	13	2	8	42	33	30	906	
shark, sandbar (brown)	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
sharksucker	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
silverside, Atlantic	0	0	0	0	0	0	0	0	1	54	3	39	0	2	0	1	2	1	0	1	0	0	0	1	2	3	1	0	0	3	1	115	
skate, barndoor	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
skate, clearnose	0	0	3	2	1	1	3	2	8	8	1	4	1	4	20	22	18	65	59	68	22	102	36	97	37	69	1	56	280	218	104	1,311	
skate, little	2,751	4,614	4,303	3,847	9,471	9,349	11,902	6,479	3,495	6,051	6,714	2,372	6,203	4,068	4,305	3,686	3,340	4,311	4,242	4,071	3,044	1,317	593	1,277	682	709	281	674	1,406	583	770	116,908	
skate, winter	1	20	34	17	114	120	85	50	31	62	51	41	88	48	62	41	31	38	45	82	53	31	23	44	51	44	16	37	97	91	82	1,629	
smelt, rainbow	0	0	0	0	5	4	2	2	0	9	9	4	0	0	0	0	0																

Appendix 5.2 cont.

Total count of finfish, lobster and squid taken in the LISTS, 1984-2014.

Year	Tows	Total Count
1984	200	122,527
1985	246	152,574
1986	316	153,383
1987	320	136,139
1988	320	216,479
1989	320	294,026
1990	297	277,183
1991	200	174,235
1992	160	186,975
1993	240	230,301
1994	240	204,795
1995	200	163,532
1996	200	165,756
1997	200	170,761
1998	200	258,082
1999	200	392,831
2000	200	271,608
2001	200	172,622
2002	200	229,284
2003	200	131,812
2004	199	250,439
2005	200	200,991
2006	120	109,330
2007	200	215,638
2008	120	164,948
2009	200	239,154
2010	78	39,340
2011	172	146,254
2012	200	170,798
2013	200	102,413
2014	199	177,250
	<hr/>	<hr/>
	6,547	5,921,460

Appendix 5.3. Annual total weight (kg) of finfish, lobster and squid taken in LISTS, 1992-2014.

Weights include all tows – number of tows shown in second row. Refer to Appendix 5.4 for details on number of tows conducted per month. Note: nw = not weighed.

Common name (number of tows)	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	Total
anchovy, bay	nw	5.6	12.2	3.6	6.6	13.3	10.3	5.8	8.3	14.5	7.7	35.3	2.8	10.5	8.6	6.8	9.4	161.3						
anchovy, striped	nw	nw	nw	nw	0.2	0.0	0.0	6.1	0.0	1.2	0.0	0.1	0.0	0.0	0.0	0.1	0.1	0.4	0.0	0.1	0.2	0.1	0.0	8.6
anchovy, spp (yoy-est)	nw	0.5	4.5	0.8	1.5	2.0	3.0	1.5	0.6	0.8	5.1	0.7	0.0	1.0	0.4	1.3	2.6	26.3						
bigeye	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.2	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4
bigeye, short	0.0	0.1	0.1	0.0	0.3	0.2	0.0	0.0	0.0	0.1	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0
black sea bass	1.8	6.4	11.0	4.7	12.1	10.5	10.6	17.2	22.6	74.8	188.3	49.6	40.5	26.4	9.3	46.8	29.8	59.5	20.1	54.2	141.0	181.2	543.3	1,561.7
blenny, feather	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.2
blue runner	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.0	2.3	0.0	1.7	2.7	0.0	0.9	8.0
bluefish	2,462.9	2,226.1	2,341.7	1,156.1	1,118.2	977.6	899.0	1,218.0	1,408.0	751.2	1,099.7	791.6	2,140.6	1,333.8	358.6	1,801.3	641.4	1,157.4	6.1	584.7	532.7	517.7	522.7	26,047.1
bonito, Atlantic	0.0	6.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.4	0.0	0.0	0.0	3.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	12.0
burrfish, striped	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	1.0
butterfish	1,357.3	1,450.1	1,202.2	1,664.5	1,844.7	2,017.2	3,661.1	4,171.6	1,458.3	1,834.0	1,924.2	682.8	1,842.7	2,097.3	1,631.4	1,446.2	1,442.0	3,186.9	166.9	1,600.8	1,891.3	1,252.5	1,707.6	41,533.6
cod, Atlantic	0.0	0.0	0.0	0.1	0.0	0.3	0.0	0.0	0.1	0.0	0.0	2.8	4.7	0.9	0.0	0.0	0.0	1.0	2.1	9.2	0.0	0.0	0.3	21.5
Gadus spp. (yoy/larvae)	nw	1.5	0	0	1.8	0.3	0.4	0	0	0	0.4	4.4												
corsefish, red	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.2
corsefish, blue spotted	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.1	0.1
crab, horseshoe	514.1	807.9	463.1	116.8	717	472.4	489.4	634.1	689.4	870.7	862.9	751	873.4	304.2	205.8	596.4	496.8	645.8	112.2	505.2	385.8	531.8	497.3	12,543.5
croaker, Atlantic	0.0	2.5	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.2	0.0	0.1	0.2	3.4
cunner	3.7	6.2	2.1	4.4	2.6	4.1	8.1	5.9	5.3	5.9	7.2	6.7	3.7	4.1	1.3	3.0	3.6	1.8	1.3	1.9	2.8	1.8	0.2	87.7
cuskeel, fawn	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2
cuskeel, striped	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.0	0.0	0.0	0.6	1.0
dogfish, smooth	863.2	1,339.1	934.6	566.8	862.8	527.3	989.8	923.0	1,038.5	1,407.6	2,814.3	1,527.4	1,435.3	1,421.7	1,176.6	2,110.2	1,134.2	2,213.3	34.4	2,031.7	1,833.3	2,162.3	2,799.2	32,146.6
dogfish, spiny	30.7	58.4	199.6	0.0	2.1	13.7	44.5	51.1	9.9	128.6	48.0	239.5	104.7	102.0	47.0	122.3	127.7	545.7	16.2	203.5	62.8	91.5	62.2	2,311.7
drum, black	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.1
eel, American	0.0	1.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.1
eel, American (yoy)	nw	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.1												
eel, conger	0.1	0.2	0.0	1.2	0.1	0.0	0.0	0.5	0.0	0.3	0.0	1.1	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.3	1.2	0.0	0.0	6.1
eel, conger (yoy)	nw	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.2												
filefish, orange	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2
filefish, planehead	0.0	0.8	0.1	0.0	0.3	0.0	0.0	0.3	0.0	0.1	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.4	2.4
flounder, American plaice	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.3
flounder, fourspot	382.4	193.6	202.4	402.9	407.2	615.3	306.0	203.9	398.6	362.7	326.9	350.1	309.3	125.9	88.1	224.9	186.3	169.8	92.0	224.2	454.5	203.4	145.0	6,375.4
flounder, smallmouth	0.6	2.6	1.5	1.2	2.3	2.4	6.4	5.2	2.7	3.8	4.9	3.0	2.8	2.4	0.6	2.6	3.2	4.7	1.4	3.5	7.5	5.2	6.0	76.5
flounder, summer	142.1	193.1	173.0	79.6	266.4	326.0	431.3	459.8	471.3	628.1	989.3	845.7	627.2	406.1	180.5	590.9	398.0	694.4	229.6	713.0	718.5	726.6	567.4	10,857.9
flounder, windowpane	286.1	578.9	597.2	356.2	1,223.6	986.1	741.1	594.2	368.8	475.5	343.3	378.8	333.7	177.5	128.9	510.8	524.0	342.8	449.3	395.9	501.1	326.6	365.6	10,986.0
flounder, winter	1,344.8	1,898.0	2,060.9	1,614.7	3,335.0	2,439.4	2,450.3	2,011.7	1,921.4	1,993.6	1,584.1	1,421.9	839.9	566.1	271.2	951.3	751.9	524.0	450.5	613.8	604.9	576.8	459.7	30,685.9
flounder, yellowtail	0.0	0.0	0.0	0.1	0.0	0.3	0.0	0.0	0.1	0.2	0.0	0.0	0.0	0.0	0.4	1.0	0.4	0.2	0.0	0.3	0.0	0.0	0.0	3.0
glassseye snapper	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.1	0.7	0.1	0.6	0.0	0.0	0.0	0.1	0.0	1.8
goatfish, red	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.1	0.9
goby, naked	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
goosefish	2.5	0.5	2.0	3.3	0.1	1.6	3.2	0.3	0.2	0.4	0.6	0.0	0.1	0.7	1.2	0.0	0.0	0.0	0.0	0.8	0.0	0.0	0.0	17.5
grubby	0.0	0.0	0.3	0.1	0.2	0.7	0.3	0.2	0.0	0.0	0.1	0.1	0.0	0.2	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.0	2.4
gunnel, rock	0.0	0.0	0.1	0.0	0.2	0.0	0.0	0.0	0.2	0.1	0.1	0.4	0.2	0.6	0.1	0.1	0.2	0.2	0.5	0.2	0.1	0.0	0.0	3.3
haddock	0.0	0.0	0.0	0.2	0.0	0.1	0.5	0.1	0.0	0.0	0.0	1.3	0.6	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	3.4
hake, red	127.7	254.4	63.9	145.6	95.5	80.5	217.5	226.5	162.6	109.7	206.6	73.4	51.6	56.0	37.4	200.4	141.3	59.5	64.3	25.1	148.6	61.1	33.5	2,642.7
hake, silver	22.0	21.9	127.6	61.6	20.0	70.8	88.3	99.6	28.8	152.2	89.6	13.9	27.3	7.1	37.7	14.6	208.5	50.0	35.4	40.3	171.0	23.6	10.6	1,422.4
hake, spotted	10.3	55.9	32.4	6.5	42.6	19.0	12.2	38.8	92.3	34.9	48.2	70.4	37.8	17.4	24.3	23.9	65.8	32.1	15.8	76.8	64.2	66.8	59.5	947.9
harvestfish	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.0	0.3
herring, Atlantic	797.5	1,120.0	769.3	1,631.7	189.8	515.1	74.6	45.4	124.1	72.6	63.9	89.1	58.3	131.1	10.3	234.2	52.1	239.2	179.0	199.4	61.5	321.2	91.2	7,070.6
herring, Atlantic (yoy-est)	nw	1.5	1.9	2.8	2.4	1.2	0.2	4.2	0.4	1.9	0.3	0.5	1.2	7.3	0.5	26.3								
herring, alewife	9.2	54.5	83.2	24.6	134.6	81.3	35.1	107.6	96.0	41.7	70.2	55.3	56.1	47.6	49.5	101.3	51.1	96.0	14.3	29.8	47.0	34.1	43.2	1,363.3
herring, blueback	8.5	4.7	31.2	7.5	6.2	16.5	5.1	1.1	6.8	11.1	2.4	4.0	6.5	5.4	2.5	9.1	3.2	14.6	3.4	3.2	1.6	4.3	4.2	163.1
herring, round	0.2	0.3	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	1.5
hogchoker	5.6	7.3	3.9	1.7	5.4	1.8	1.9	5.0	5.9	10.5	13.3	8.6	9.5	8.7	3.2	11.4	5.6	4.5	4.4	16.8	30.7	27.2	27.8	220.7
jack, crevalle	0.0	0.5	0.5	0.1	0.0	0.6	0.0	0.7	0.0	0.0	0.1	0.2	0.2	0.2	0.0	0.1	0.0	0.1	0.0	0.4	0.2	0.0	0.2	4.1
jack, yellow	0.2	0.2	0.4	2.1																				

Appendix 5.3 cont.

Common name	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	Total	
(number of tows)	160	240	240	200	200	200	200	200	200	200	200	200	199	200	120	200	160	200	78	172	200	200	199	4,368	
mackerel, Spanish	1.5	5.3	6.4	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	2.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	15.5
menhaden, Atlantic	60.6	103.9	87.8	41.9	40.5	38.5	9.2	90.9	31.8	4.7	96.3	344.9	110.7	77.9	5.5	63.9	10.4	18.0	2.7	69.8	144.6	87.5	267.8	1,809.8	
moonfish	1.5	0.6	4.1	2.1	11.6	4.6	13.4	9.6	15.0	3.8	7.4	2.3	3.4	6.0	3.5	12.0	13.4	19.5	0.0	6.3	3.6	10.0	23.2	176.9	
mullet, white	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.2
ocean pout	7.7	16.4	9.1	6.5	7.2	4.8	2.7	3.9	4.9	2.3	4.3	2.9	5.4	0.7	0.9	3.2	2.1	4.8	1.4	4.5	2.0	0.0	0.0	0.0	97.7
perch, silver	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.3
perch, white	0.0	0.3	0.3	0.0	0.1	0.9	0.0	0.4	0.2	0.0	0.0	1.4	0.5	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.2	0.0	0.2	0.0	4.8
pinfish	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.2
pipefish, northern	0.4	0.6	0.2	0.1	0.0	0.1	0.0	0.1	0.2	0.3	0.2	0.4	0.2	0.3	0.2	0.2	0.0	0.2	0.3	0.3	0.1	0.2	0.1	0.2	4.7
pollock	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.8	0.1	0.5	0.0	0.1	0.0	0.0	2.0
pompano, African	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
puffer, northern	0.1	0.9	0.4	0.1	0.3	0.1	0.5	1.1	0.4	0.7	0.3	0.3	0.4	0.3	0.0	0.5	0.0	0.4	0.0	0.9	3.1	0.3	1.3	1.3	12.4
ray, bullnose ray	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5.7	0.0	5.7
ray, roughtail stingray	0.0	0.0	0.0	0.0	0.0	50.6	3.4	0.0	0.0	2.5	24.4	0.0	4.1	0.0	0.0	0.0	3.0	0.0	0.0	13.0	5.0	0.0	0.0	0.0	106.0
rockling, fourbeard	12.8	15.7	8.5	14.7	8.6	17.3	11.6	28.8	14.7	21.5	9.7	9.2	13.0	6.8	1.5	7.6	7.1	3.9	2.9	4.0	3.5	0.2	0.4	0.4	224.0
rudderfish, banded	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4
salmon, Atlantic	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
sand lance, American	nw	0.3	0.6	0.4	0.0	0.1	0.3	0.3	0.3	0.3	0.1	0.2	0.2	0.0	0.3	0.7	2.0	2.0	5.2	7.5	0.2	0.1	0.2	0.2	26.0
sand lance, (yoy - est)	nw	0.0	0.8	0.1	0.0	0.0	0.1	0.4	0.0	0.6	0.0	0.0	0.0	0.0	2.9	0.1	0.2	2.3	0.0	3.8	0.0	0.0	0.0	0.0	11.3
scad, bigeye	0.0	0.0	0.3	0.0	0.1	0.1	0.1	1.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.0
scad, mackerel	0.2	0.0	0.4	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	1.1
scad, rough	0.0	4.4	0.2	0.0	1.5	2.0	0.0	0.0	0.0	0.7	0.7	0.5	0.7	1.9	0.5	0.7	0.0	2.8	0.0	6.8	1.1	1.3	0.5	0.5	26.3
scad, round	0.0	0.0	0.0	0.0	0.0	0.2	0.3	0.1	0.2	0.0	0.0	0.3	0.3	0.3	0.0	0.3	0.0	0.1	0.0	0.1	0.0	0.1	0.1	0.1	2.4
sculpin, longhorn	9.0	3.2	1.6	1.3	2.1	0.8	1.0	0.3	5.0	1.5	0.9	2.0	3.4	0.0	0.0	0.8	0.3	0.3	0.4	2.0	0.2	0.4	0.0	0.0	36.5
scup	837.7	867.9	878.1	770.5	739.4	530.5	740.5	3,641.3	6,679.0	5,828.4	13,814.0	5,221.9	6,801.1	3,080.7	4,636.1	5,333.5	6,509.9	6,332.1	1,971.6	6,759.5	6,170.2	5,945.6	5,161.4	99,250.9	
sea raven	3.9	0.6	0.2	0.7	1.5	0.4	11.3	4.9	9.2	4.1	4.1	1.6	2.4	0.5	0.0	3.6	0.0	1.7	1.6	0.9	1.1	0.0	1.5	1.5	55.8
seahorse, lined	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
searobin, northern	35.6	97.9	66.7	166.9	57.4	60.4	39.4	52.0	251.2	222.7	267.3	252.2	112.0	21.3	74.5	74.2	58.8	194.3	149.5	85.5	405.2	161.7	225.9	3,132.6	
searobin, striped	305.1	260.0	208.6	277.5	278.7	230.5	509.7	497.0	1,036.1	861.0	1,065.0	805.1	465.4	183.7	113.5	217.0	263.0	471.8	66.4	558.7	1,086.4	1,112.5	1,020.8	11,893.5	
seasnail	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.2	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7
sennet, northern	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.1	0.2	0.0	0.0	0.7	0.0	0.2	0.0	0.4	0.0	0.1	0.3	0.0	0.0	0.0	2.7
shad, American	63.3	138.9	165.8	81.4	36.2	66.8	60.2	117.3	25.8	9.6	40.3	40.8	24.2	18.2	6.1	15.8	20.2	28.9	8.6	17.5	25.3	15.3	12.3	1,038.8	
shad, gizzard	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.1	0.1	0.2	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.9
shad, hickory	4.9	4.4	7.6	2.5	10.2	9.1	15.9	19.4	17.1	6.7	19.6	20.1	14.2	43.1	19.1	10.4	1.1	3.6	0.4	1.5	14.1	10.8	10.5	266.3	
sharksucker	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3
silverside, Atlantic	0.1	1.0	0.3	0.9	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.1	0.2	0.3	0.1	0.0	0.0	0.3	0.1	0.1	3.9
skate, barndoor	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4
skate, clearnose	10.3	11.3	1.8	11.0	1.7	7.4	36.8	39.4	37.9	132.4	107.3	130.8	48.2	187.1	52.4	193.3	78.1	148.5	4.5	109.8	491.7	387.0	207.7	2,436.4	
skate, little	1,389.0	2,534.8	3,091.5	1,055.3	2,801.8	1,945.8	2,085.5	1,829.6	1,604.7	2,022.6	2,121.9	2,187.3	1,689.8	682.5	310.6	697.0	327.4	390.0	148.3	359.4	657.9	317.8	428.2	30,678.7	
skate, winter	105.3	220.9	139.2	89.2	212.7	109.7	180.7	89.8	66.5	112.2	133.5	162.1	100.3	59.9	60.0	117.8	140.8	108.5	37.7	101.2	179.8	111.2	133.8	2,772.8	
smelt, rainbow	0.0	0.6	0.6	0.3	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7
spot	0.0	10.6	4.3	0.3	14.1	1.1	0.0	5.7	17.8	1.3	7.2	0.1	0.9	0.0	1.2	0.0	21.3	0.2	0.0	0.7	107.5	195.4	1.8	391.5	
squid, long-finned	844.9	1,629.1	965.4	796.4	720.4	515.2	767.0	826.4	582.3	346.2	279.9	573.2	953.4	683.5	326.0	773.6	330.1	648.4	161.4	370.7	333.9	170.8	582.3	14,180.5	
stargazer, northern	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.2
striped bass	89.4	210.3	198.6	185.3	373.5	509.9	484.2	815.4	602.6	472.5	855.2	770.3	811.8	675.1	418.7	888.0	456.3	897.4	173.2	721.9	278.0	421.0	407.5	11,716.1	
sturgeon, Atlantic	244.8	633.6	848.6	145.5	19.9	37.8	189.7	498.6	79.0	270.6	275.3	550.2	117.6	152.7	368.7	336.4	111.3	286.6	5.6	181.9	154.2	98.0	272.4	5,879.0	
tautog	508.3	320.0	373.9	95.1	225.9	271.8	347.1	326.6	463.5	491.2	921.1	346.0	353.7	269.2	301.4	551.4	309.								

Appendix 5.4. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1984.

Finfish species are in order of descending count. Number of tows (sample size)=102.

species	count	%	weight	%	species	count	%	weight	%
butterfish	18,700	31.0	.	.	Atlantic mackerel	48	0.1	.	.
windowpane flounder	13,746	22.8	.	.	spotted hake	46	0.1	.	.
winter flounder	6,847	11.4	.	.	sea raven	32	0.1	.	.
bluefish	6,738	11.2	.	.	ocean pout	25	0	.	.
scup	3,225	5.4	.	.	rough scad	22	0	.	.
fourspot flounder	1,868	3.1	.	.	longhorn sculpin	12	0	.	.
little skate	1,491	2.5	.	.	black sea bass	11	0	.	.
red hake	1,323	2.2	.	.	moonfish	7	0	.	.
American shad	982	1.6	.	.	Atlantic sturgeon	6	0	.	.
blueback herring	925	1.5	.	.	round herring	5	0	.	.
striped searobin	697	1.2	.	.	spiny dogfish	4	0	.	.
silver hake	575	1.0	.	.	American eel	2	0	.	.
smooth dogfish	534	0.9	.	.	striped bass	2	0	.	.
tautog	472	0.8	.	.	oyster toadfish	2	0	.	.
northern searobin	448	0.7	.	.	goosefish	1	0	.	.
fourbeard rockling	303	0.5	.	.	northern sennet	1	0	.	.
weakfish	260	0.4	.	.	northern puffer	1	0	.	.
hogchoker	252	0.4	.	.	red goatfish	1	0	.	.
cunner	220	0.4	.	.	Total	60,230			
summer flounder	150	0.2	.	.					
alewife	108	0.2	.	.	<u>Invertebrates</u>				
hickory shad	71	0.1	.	.	American lobster	2865	100	.	.
Atlantic menhaden	67	0.1	.	.	Total	2,865			

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1985.
Finfish species are in order of descending count. Number of tows (sample size)=126.

species	count	%	weight	%	species	count	%	weight	%
butterfish	34,512	41.4	.	.	spot	26	0	.	.
scup	12,155	14.6	.	.	round herring	15	0	.	.
windowpane flounder	11,194	13.4	.	.	rough scad	14	0	.	.
winter flounder	7,980	9.6	.	.	Atlantic mackerel	13	0	.	.
bluefish	5,302	6.4	.	.	spiny dogfish	13	0	.	.
weakfish	2,650	3.2	.	.	winter skate	13	0	.	.
northern searobin	2,098	2.5	.	.	alewife	9	0	.	.
little skate	1,705	2.0	.	.	planehead filefish	7	0	.	.
fourspot flounder	1,289	1.5	.	.	rock gunnel	4	0	.	.
striped searobin	1,078	1.3	.	.	oyster toadfish	4	0	.	.
red hake	573	0.7	.	.	goosefish	3	0	.	.
Atlantic herring	504	0.6	.	.	ocean pout	3	0	.	.
smooth dogfish	405	0.5	.	.	Atlantic bonito	2	0	.	.
tautog	323	0.4	.	.	crevalle jack	1	0	.	.
American shad	280	0.3	.	.	grubby	1	0	.	.
silver hake	250	0.3	.	.	gray triggerfish	1	0	.	.
summer flounder	175	0.2	.	.	hickory shad	1	0	.	.
hogchoker	163	0.2	.	.	orange filefish	1	0	.	.
moonfish	142	0.2	.	.	northern puffer	1	0	.	.
blueback herring	100	0.1	.	.	Atlantic sturgeon	1	0	.	.
longhorn sculpin	80	0.1	.	.	Atlantic tomcod	1	0	.	.
cunner	51	0.1	.	.	Total	83,395		-	
sea raven	50	0.1	.	.					
fourbeard rockling	44	0.1	.	.					
Atlantic menhaden	38	0	.	.	<u>Invertebrates</u>				
black sea bass	35	0	.	.	American lobster	1589	100	.	.
spotted hake	27	0	.	.	Total	1,589		-	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1986.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight. Number of tows (sample size)=196.

species	count	%	weight	%	species	count	%	weight	%
butterfish	25,192	28.0	.	.	winter skate	32	0	.	.
windowpane flounder	18,848	20.9	.	.	spotted hake	30	0	.	.
winter flounder	15,341	17.0	.	.	black sea bass	28	0	.	.
scup	7,910	8.8	.	.	spot	25	0	.	.
weakfish	5,427	6.0	.	.	Atlantic mackerel	19	0	.	.
little skate	3,210	3.6	.	.	moonfish	14	0	.	.
bluefish	2,789	3.1	.	.	ocean pout	14	0	.	.
red hake	2,657	3.0	.	.	oyster toadfish	9	0	.	.
Atlantic herring	1,999	2.2	.	.	hickory shad	6	0	.	.
fourspot flounder	1,487	1.7	.	.	rough scad	5	0	.	.
striped searobin	886	1.0	.	.	Atlantic sturgeon	4	0	.	.
silver hake	723	0.8	.	.	clearnose skate	2	0	.	.
tautog	566	0.6	.	.	American eel	1	0	.	.
smooth dogfish	430	0.5	.	.	goosefish	1	0	.	.
summer flounder	414	0.5	.	.	grubby	1	0	.	.
northern searobin	396	0.4	.	.	northern pipefish	1	0	.	.
American shad	344	0.4	.	.	northern puffer	1	0	.	.
Atlantic menhaden	318	0.4	.	.	smallmouth flounder	1	0	.	.
blueback herring	256	0.3	.	.	striped bass	1	0	.	.
alewife	216	0.2	.	.	Total	90,031		-	
fourbeard rockling	123	0.1	.	.					
cunner	76	0.1	.	.					
sea raven	70	0.1	.	.	<u>Invertebrates</u>				
hogchoker	60	0.1	.	.	American lobster	2,553	28.1	.	.
longhorn sculpin	51	0.1	.	.	long-finned squid	6,537	71.9	.	.
spiny dogfish	47	0.1	.	.	Total	9,090		-	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1987.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
winter flounder	15,600	25.6	.	.	longhorn sculpin	32	0.1	.	.
butterfish	14,674	24.1	.	.	spotted hake	22	0	.	.
windowpane flounder	11,031	18.1	.	.	spiny dogfish	19	0	.	.
scup	5,029	8.3	.	.	ocean pout	14	0	.	.
bluefish	2,611	4.3	.	.	black sea bass	13	0	.	.
little skate	2,140	3.5	.	.	winter skate	13	0	.	.
red hake	1,729	2.8	.	.	striped bass	10	0	.	.
Atlantic herring	1,628	2.7	.	.	Atlantic tomcod	8	0	.	.
fourspot flounder	1,298	2.1	.	.	smallmouth flounder	7	0	.	.
silver hake	906	1.5	.	.	moonfish	6	0	.	.
alewife	754	1.2	.	.	rock gunnel	4	0	.	.
striped searobin	543	0.9	.	.	Atlantic sturgeon	4	0	.	.
summer flounder	374	0.6	.	.	spot	3	0	.	.
American shad	371	0.6	.	.	clearnose skate	2	0	.	.
tautog	363	0.6	.	.	hickory shad	2	0	.	.
Atlantic menhaden	329	0.5	.	.	Atlantic bonito	1	0	.	.
smooth dogfish	257	0.4	.	.	Atlantic mackerel	1	0	.	.
weakfish	248	0.4	.	.	round herring	1	0	.	.
fourbeard rockling	241	0.4	.	.	sea lamprey	1	0	.	.
northern searobin	220	0.4	.	.	Total	60,862		-	
sea raven	86	0.1	.	.					
blueback herring	79	0.1	.	.	Invertebrates				
cunner	79	0.1	.	.	American lobster	3,544	25.1	.	.
hogchoker	61	0.1	.	.	long-finned squid	10,552	74.9	.	.
rough scad	48	0.1	.	.	Total	14,096		-	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1988.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	45,983	36.7	.	.	ocean pout	30	0	.	.
winter flounder	25,695	20.5	.	.	Atlantic mackerel	24	0	.	.
windowpane flounder	19,497	15.6	.	.	spot	18	0	.	.
scup	10,184	8.1	.	.	black sea bass	17	0	.	.
little skate	6,539	5.2	.	.	striped bass	17	0	.	.
bluefish	3,688	2.9	.	.	yellowtail flounder	6	0	.	.
fourspot flounder	2,478	2.0	.	.	grubby	5	0	.	.
red hake	1,933	1.5	.	.	rock gunnel	5	0	.	.
weakfish	1,287	1.0	.	.	rainbow smelt	5	0	.	.
silver hake	1,210	1.0	.	.	crevalle jack	4	0	.	.
striped searobin	1,194	1.0	.	.	bigeye scad	2	0	.	.
Atlantic herring	1,193	1.0	.	.	bigeye	2	0	.	.
American shad	1,187	0.9	.	.	planehead filefish	2	0	.	.
northern searobin	474	0.4	.	.	hickory shad	2	0	.	.
tautog	455	0.4	.	.	northern puffer	2	0	.	.
smooth dogfish	385	0.3	.	.	Atlantic sturgeon	2	0	.	.
summer flounder	320	0.3	.	.	Atlantic tomcod	2	0	.	.
fourbeard rockling	302	0.2	.	.	Atlantic bonito	1	0	.	.
blueback herring	164	0.1	.	.	dwarf goatfish	1	0	.	.
alewife	153	0.1	.	.	goosefish	1	0	.	.
moonfish	137	0.1	.	.	northern pipefish	1	0	.	.
rough scad	128	0.1	.	.	short bigeye	1	0	.	.
longhorn sculpin	103	0.1	.	.	striped cusk-eel	1	0	.	.
winter skate	101	0.1	.	.	sea lamprey	1	0	.	.
spotted hake	87	0.1	.	.	Total	125,344			-
hogchoker	75	0.1	.	.					
Atlantic menhaden	69	0.1	.	.					
sea raven	50	0	.	.	<u>Invertebrates</u>				
cunner	48	0	.	.	American lobster	2,114	8.5	.	.
spiny dogfish	39	0	.	.	long-finned squid	22,769	91.5	.	.
smallmouth flounder	34	0	.	.	Total	24,883			-

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1989.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	47,089	29.3	.	.	sea raven	34	0	.	.
winter flounder	32,361	20.2	.	.	black sea bass	15	0	.	.
windowpane flounder	25,109	15.6	.	.	rough scad	11	0	.	.
scup	17,391	10.8	.	.	striped bass	11	0	.	.
bluefish	8,649	5.4	.	.	yellow jack	11	0	.	.
little skate	7,079	4.4	.	.	goosefish	9	0	.	.
red hake	5,689	3.5	.	.	smallmouth flounder	9	0	.	.
weakfish	5,496	3.4	.	.	rock gunnel	8	0	.	.
American shad	1,977	1.2	.	.	grubby	7	0	.	.
fourspot flounder	1,877	1.2	.	.	spotted hake	7	0	.	.
striped searobin	1,763	1.1	.	.	rainbow smelt	4	0	.	.
silver hake	1,697	1.1	.	.	planehead filefish	3	0	.	.
Atlantic herring	1,154	0.7	.	.	Atlantic sturgeon	3	0	.	.
tautog	600	0.4	.	.	Atlantic tomcod	3	0	.	.
fourbeard rockling	397	0.2	.	.	bigeye	2	0	.	.
blueback herring	307	0.2	.	.	American eel	2	0	.	.
northern searobin	297	0.2	.	.	short bigeye	2	0	.	.
Atlantic mackerel	237	0.1	.	.	oyster toadfish	2	0	.	.
Atlantic menhaden	230	0.1	.	.	white perch	2	0	.	.
smooth dogfish	202	0.1	.	.	northern sennet	1	0	.	.
alewife	190	0.1	.	.	northern puffer	1	0	.	.
longhorn sculpin	107	0.1	.	.	banded rudderfish	1	0	.	.
cunner	106	0.1	.	.	Spanish mackerel	1	0	.	.
hogchoker	91	0.1	.	.	Total	160,581		-	
winter skate	91	0.1	.	.					
spiny dogfish	66	0	.	.					
ocean pout	58	0	.	.	Invertebrates				
bigeye scad	45	0	.	.	American lobster	3,447	19.9	.	.
moonfish	42	0	.	.	long-finned squid	13,883	80.1	.	.
summer flounder	35	0	.	.	Total	17,330		-	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1990.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
winter flounder	47,184	31.1	.	.	seasnail	8	0	.	.
butterfish	45,373	29.9	.	.	planehead filefish	7	0	.	.
scup	15,393	10.2	.	.	moonfish	7	0	.	.
windowpane flounder	9,825	6.5	.	.	rock gunnel	7	0	.	.
Atlantic herring	8,779	5.8	.	.	yellow jack	7	0	.	.
little skate	6,456	4.3	.	.	grubby	4	0	.	.
bluefish	4,688	3.1	.	.	spot	4	0	.	.
fourspot flounder	3,270	2.2	.	.	Atlantic sturgeon	4	0	.	.
silver hake	2,334	1.5	.	.	oyster toadfish	4	0	.	.
red hake	2,237	1.5	.	.	goosefish	3	0	.	.
weakfish	1,921	1.3	.	.	smallmouth flounder	3	0	.	.
striped searobin	866	0.6	.	.	Atlantic tomcod	3	0	.	.
tautog	554	0.4	.	.	clearnose skate	2	0	.	.
American shad	406	0.3	.	.	lookdown	2	0	.	.
fourbeard rockling	299	0.2	.	.	red goatfish	2	0	.	.
longhorn sculpin	243	0.2	.	.	rainbow smelt	2	0	.	.
northern searobin	232	0.2	.	.	bigeye scad	1	0	.	.
Atlantic menhaden	219	0.1	.	.	bigeye	1	0	.	.
smooth dogfish	209	0.1	.	.	hickory shad	1	0	.	.
summer flounder	170	0.1	.	.	mackerel scad	1	0	.	.
cunner	168	0.1	.	.	northern kingfish	1	0	.	.
alewife	160	0.1	.	.	northern puffer	1	0	.	.
spiny dogfish	150	0.1	.	.	red cornetfish	1	0	.	.
hogchoker	84	0.1	.	.	sandbar shark	1	0	.	.
winter skate	61	0	.	.	sea lamprey	1	0	.	.
blueback herring	46	0	.	.	yellowtail flounder	1	0	.	.
striped bass	45	0	.	.	Total	151,600			-
sea raven	42	0	.	.					
ocean pout	39	0	.	.					
black sea bass	27	0	.	.	<u>Invertebrates</u>				
spotted hake	21	0	.	.	American lobster	5,369	27.0.	.	.
Atlantic mackerel	10	0	.	.	long-finned squid	14,538	73.0.	.	.
rough scad	10	0	.	.	Total	19,907			-

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1991.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
scup	45,790	29.9	.	.	moonfish	24	0	.	.
butterfish	40,537	26.4	.	.	smallmouth flounder	20	0	.	.
winter flounder	26,623	17.4	.	.	sea raven	19	0	.	.
windowpane flounder	8,482	5.5	.	.	spiny dogfish	14	0	.	.
little skate	6,479	4.2	.	.	yellow jack	11	0	.	.
bluefish	5,845	3.8	.	.	goosefish	8	0	.	.
weakfish	4,320	2.8	.	.	northern puffer	5	0	.	.
Atlantic herring	4,003	2.6	.	.	northern kingfish	4	0	.	.
fourspot flounder	3,553	2.3	.	.	Atlantic tomcod	4	0	.	.
red hake	2,085	1.4	.	.	Atlantic sturgeon	3	0	.	.
silver hake	1,537	1.0	.	.	clearnose skate	2	0	.	.
striped searobin	865	0.6	.	.	Atlantic mackerel	2	0	.	.
northern searobin	609	0.4	.	.	mackerel scad	2	0	.	.
tautog	501	0.3	.	.	rainbow smelt	2	0	.	.
American shad	361	0.2	.	.	Spanish mackerel	2	0	.	.
Atlantic menhaden	348	0.2	.	.	spot	2	0	.	.
summer flounder	263	0.2	.	.	bigeye scad	1	0	.	.
smooth dogfish	193	0.1	.	.	planehead filefish	1	0	.	.
fourbeard rockling	163	0.1	.	.	hickory shad	1	0	.	.
longhorn sculpin	139	0.1	.	.	red goatfish	1	0	.	.
hogchoker	104	0.1	.	.	rough scad	1	0	.	.
alewife	103	0.1	.	.	sea lamprey	1	0	.	.
cunner	75	0	.	.	oyster toadfish	1	0	.	.
spotted hake	73	0	.	.	Total	153,389			-
winter skate	50	0	.	.					
ocean pout	42	0	.	.	Invertebrates				
black sea bass	39	0	.	.	American lobster	8,524	40.9	.	.
blueback herring	38	0	.	.	long-finned squid	12,322	59.1	.	.
striped bass	38	0	.	.	Total	20,846			-

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1992.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Number of tows (sample size)=160.

species	count	%	weight	%	species	count	%	weight	%
butterfish	95,961	65.7	1,357.3	11.7	black sea bass	5	0	1.8	0
scup	13,646	9.3	837.7	7.2	northern pipefish	5	0	0.4	0
winter flounder	9,548	6.5	1,344.8	11.5	Atlantic mackerel	4	0	1.0	0
bluefish	5,269	3.6	2,462.9	21.1	sea raven	4	0	3.9	0
Atlantic herring	4,565	3.1	797.5	6.8	northern kingfish	2	0	0.2	0
little skate	3,495	2.4	1,389.0	11.9	round herring	2	0	0.2	0
windowpane flounder	2,980	2.0	286.1	2.5	yellow jack	2	0	0.2	0
fourspot flounder	2,774	1.9	382.4	3.3	Atlantic silverside	1	0	0.1	0
red hake	1,606	1.1	127.7	1.1	conger eel	1	0	0.1	0
weakfish	1,317	0.9	94.8	0.8	northern puffer	1	0	0.1	0
Atlantic menhaden	1,115	0.8	60.6	0.5	Spanish mackerel	1	0	1.5	0
striped searobin	857	0.6	305.1	2.6	Total	146,035		11,648.2	
silver hake	544	0.4	22.0	0.2					
American shad	380	0.3	63.3	0.5	Invertebrates				
northern searobin	313	0.2	35.6	0.3	American lobster	8,160	19.9	1,537.9	28.6
smooth dogfish	304	0.2	863.2	7.4	blue mussel	nc	nc	1,157.1	21.5
tautog	265	0.2	508.3	4.4	long-finned squid	32,780	80.1	844.9	15.7
summer flounder	186	0.1	142.1	1.2	horseshoe crab	nc	nc	514.1	9.6
blueback herring	175	0.1	8.5	0.1	lady crab	nc	nc	375.4	7.0
fourbeard rockling	150	0.1	12.8	0.1	rock crab	nc	nc	239.1	4.5
alewife	122	0.1	9.2	0.1	boring sponge	nc	nc	225.5	4.2
spotted hake	68	0	10.3	0.1	spider crab	nc	nc	186.0	3.5
moonfish	62	0	1.5	0	starfish spp.	nc	nc	148.6	2.8
hogchoker	61	0	5.6	0	whelks	nc	nc	57.5	1.1
striped bass	42	0	89.4	0.8	flat claw hermit crab	nc	nc	34.7	0.6
longhorn sculpin	31	0	9.0	0.1	bluecrab	nc	nc	18.1	0.3
winter skate	31	0	105.3	0.9	mantis shrimp	nc	nc	10.3	0.2
cunner	30	0	3.7	0	northern moon snail	nc	nc	8.6	0.2
Atlantic sturgeon	30	0	244.8	2.1	common oyster	nc	nc	7.3	0.1
ocean pout	18	0	7.7	0.1	lion's mane jellyfish	nc	nc	2.4	0
hickory shad	12	0	4.9	0	surf clam	nc	nc	1.7	0
smallmouth flounder	12	0	0.6	0	hard clams	nc	nc	1.2	0
goosefish	10	0	2.5	0	bushy bryozoan	nc	nc	1.0	0
clearnose skate	8	0	10.3	0.1	purple sea urchin	nc	nc	0.4	0
Atlantic tomcod	8	0	1.3	0	mud crabs	nc	nc	0.3	0
mackerel scad	6	0	0.2	0	star coral	nc	nc	0.1	0
spiny dogfish	6	0	30.7	0.3	Total	40,940		5,372	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1993.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	35,361	33.0	847.8	7.1	goosefish	3	0	0.3	0
scup	18,785	17.6	581.4	4.8	American sand lance	3	0	0.3	0
winter flounder	16,090	15.0	1,855.7	15.4	Atlantic bonito	2	0	6.4	0.1
windowpane flounder	7,953	7.4	547.6	4.6	lumpfish	2	0	0.2	0
Atlantic herring	6,269	5.9	1,119.8	9.3	moonfish	2	0	0.2	0
little skate	5,186	4.8	2,172.3	18.1	sea lamprey	2	0	1.0	0
bluefish	4,402	4.1	1,343.2	11.2	Atlantic salmon	1	0	0.1	0
red hake	3,963	3.7	232.0	1.9	American eel	1	0	1.6	0
fourspot flounder	1,262	1.2	182.3	1.5	northern sennet	1	0	0.1	0
weakfish	1,142	1.1	60.3	0.5	orange filefish	1	0	0.1	0
striped searobin	1,079	1.0	165.4	1.4	round herring	1	0	0.1	0
northern searobin	935	0.9	96.8	0.8	red cornetfish	1	0	0.1	0
American shad	791	0.7	101.1	0.8	red goatfish	1	0	0.1	0
alewife	788	0.7	48.2	0.4	short bigeye	1	0	0.1	0
silver hake	500	0.5	21.1	0.2	sea raven	1	0	0.6	0
spotted hake	331	0.3	36.7	0.3	yellow jack	1	0	0.1	0
smooth dogfish	283	0.3	857.6	7.1	Total	107,035		12,012.4	
Atlantic menhaden	271	0.3	94.1	0.8					
fourbeard rockling	241	0.2	15.6	0.1	Invertebrates				
summer flounder	224	0.2	137.9	1.1	American lobster	10,306	20.6	2,173.5	34.4
tautog	157	0.1	308.2	2.6	long-finned squid	39,723	79.4	1,176.5	18.6
Spanish mackerel	136	0.1	2.2	0	blue mussel	nc	nc	945.1	15.0
blueback herring	96	0.1	4.3	0	horseshoe crab	nc	nc	673.8	10.7
rough scad	92	0.1	3.8	0	spider crab	nc	nc	511.2	8.1
striped bass	78	0.1	198.7	1.7	lady crab	nc	nc	428.0	6.8
ocean pout	66	0.1	16.4	0.1	rock crab	nc	nc	155.9	2.5
cunner	64	0.1	6.1	0.1	flat claw hermit crab	nc	nc	45.7	0.7
Atlantic sturgeon	60	0.1	633.6	5.3	starfish spp.	nc	nc	37.4	0.6
winter skate	59	0.1	213.2	1.8	boring sponge	nc	nc	36.6	0.6
spot	57	0.1	4.5	0	whelks	nc	nc	34.0	0.5
hogchoker	56	0.1	5.2	0	mantis shrimp	nc	nc	31.6	0.5
Atlantic silverside	54	0.1	1.0	0	lion's mane jellyfish	nc	nc	27.6	0.4
northern puffer	23	0	0.4	0	bluecrab	nc	nc	20.0	0.3
smallmouth flounder	23	0	2.1	0	northern moon snail	nc	nc	8.9	0.1
Atlantic croaker	20	0	1.1	0	common oyster	nc	nc	2.0	0
black sea bass	16	0	5.0	0	surf clam	nc	nc	1.0	0
spiny dogfish	14	0	58.4	0.5	hard clams	nc	nc	0.9	0
Atlantic mackerel	11	0	0.9	0	purple sea urchin	nc	nc	0.7	0
longhorn sculpin	11	0	3.2	0	arks	nc	nc	0.7	0
planehead filefish	9	0	0.7	0	mud crabs	nc	nc	0.4	0
hickory shad	9	0	4.1	0	star coral	nc	nc	0.3	0
northern pipefish	9	0	0.4	0	blood star	nc	nc	0.2	0
rainbow smelt	9	0	0.6	0	common slipper shell	nc	nc	0.2	0
crevalle jack	5	0	0.4	0	sand shrimp	nc	nc	0.1	0
northern kingfish	5	0	0.6	0	sand dollar	nc	nc	0.1	0
Atlantic tomcod	5	0	0.8	0	northern red shrimp	nc	nc	0.1	0
clearnose skate	4	0	7.7	0.1	polychaetes	nc	nc	0.1	0
white perch	4	0	0.3	0	Total	50,029		6,313	
conger eel	3	0	0.2	0					

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1994.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	33,538	28.7	776.8	6.3	longhorn sculpin	7	0	1.6	0
scup	25,451	21.8	660.8	5.4	grubby	5	0	0.3	0
winter flounder	20,615	17.6	1,992.2	16.2	mackerel scad	4	0	0.4	0
bluefish	7,703	6.6	1,159.8	9.4	Atlantic silverside	3	0	0.3	0
windowpane flounder	6,062	5.2	574.5	4.7	bigeye scad	2	0	0.2	0
little skate	5,604	4.8	2,565.3	20.9	lookdown	2	0	0.2	0
Atlantic herring	3,836	3.3	768.6	6.3	northern puffer	2	0	0.2	0
weakfish	3,320	2.8	160.0	1.3	Atlantic tomcod	2	0	0.3	0
silver hake	1,703	1.5	112.9	0.9	bigeye	1	0	0.1	0
fourspot flounder	1,494	1.3	195.6	1.6	clearnose skate	1	0	1.8	0
American shad	1,289	1.1	133.2	1.1	inshore lizardfish	1	0	0.1	0
alewife	1,211	1.0	75.0	0.6	northern pipefish	1	0	0.1	0
blueback herring	1,052	0.9	26.6	0.2	rock gunnel	1	0	0.1	0
striped searobin	927	0.8	183.6	1.5	sea raven	1	0	0.2	0
northern searobin	800	0.7	63.7	0.5	white perch	1	0	0.3	0
red hake	490	0.4	54.0	0.4	yellow jack	1	0	0.1	0
smooth dogfish	310	0.3	816.3	6.6	Total	117,002		12,284.5	
Atlantic menhaden	276	0.2	61.4	0.5	Invertebrates				
summer flounder	242	0.2	141.6	1.2	American lobster	7,057	31.6	1,533.9	38.6
tautog	207	0.2	346.5	2.8	long-finned squid	15,299	68.4	594.8	15.0
spotted hake	148	0.1	25.7	0.2	horseshoe crab	nc	nc	386.7	9.7
moonfish	93	0.1	2.6	0	blue mussel	nc	nc	377.5	9.5
fourbeard rockling	92	0.1	8.4	0.1	lady crab	nc	nc	338.5	8.5
striped bass	81	0.1	198.6	1.6	spider crab	nc	nc	335.0	8.4
Atlantic sturgeon	60	0.1	848.6	6.9	rock crab	nc	nc	136.8	3.4
spiny dogfish	55	0	186.2	1.5	starfish spp.	nc	nc	124.6	3.1
ocean pout	42	0	9.1	0.1	flat claw hermit crab	nc	nc	51.4	1.3
hogchoker	36	0	3.8	0	northern moon snail	nc	nc	34.6	0.9
black sea bass	33	0	10.9	0.1	common oyster	nc	nc	18.4	0.5
winter skate	33	0	101.5	0.8	whelks	nc	nc	14.1	0.4
American sand lance	25	0	0.6	0	mantis shrimp	nc	nc	9.8	0.2
Spanish mackerel	25	0	1.7	0	lion's mane jellyfish	nc	nc	4.2	0.1
cunner	18	0	1.3	0	bluecrab	nc	nc	3.7	0.1
smallmouth flounder	15	0	1.3	0	arks	nc	nc	3.0	0.1
hickory shad	14	0	3.7	0	boring sponge	nc	nc	1.9	0
rough scad	13	0	0.2	0	hard clams	nc	nc	1.3	0
Atlantic mackerel	11	0	0.9	0	bushy bryozoan	nc	nc	0.6	0
spot	11	0	1.1	0	mud crabs	nc	nc	0.3	0
rainbow smelt	9	0	0.6	0	surf clam	nc	nc	0.3	0
crevalle jack	8	0	0.5	0	purple sea urchin	nc	nc	0.1	0
goosefish	8	0	2.0	0	Total	22,356		3,972	
northern kingfish	7	0	0.5	0					

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1995.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	64,930	50.1	1,664.5	15.2	spot	3	0	0.3	0
winter flounder	15,558	12.0	1,614.7	14.7	Atlantic cod	2	0	0.1	0
scup	13,985	10.8	770.5	7.0	conger eel	2	0	1.2	0
Atlantic herring	9,135	7.0	1,631.7	14.9	haddock	2	0	0.2	0
bluefish	5,524	4.3	1,156.1	10.5	northern pipefish	2	0	0.1	0
windowpane flounder	3,815	2.9	356.2	3.2	sea raven	2	0	0.7	0
weakfish	2,881	2.2	275.7	2.5	African pompano	1	0	0.1	0
fourspot flounder	2,584	2.0	402.9	3.7	crevalle jack	1	0	0.1	0
little skate	2,372	1.8	1,055.3	9.6	grubby	1	0	0.1	0
red hake	1,977	1.5	145.6	1.3	Atlantic mackerel	1	0	0.1	0
silver hake	1,941	1.5	61.6	0.6	mackerel scad	1	0	0.1	0
northern searobin	1,317	1.0	166.9	1.5	northern puffer	1	0	0.1	0
American shad	755	0.6	81.4	0.7	oyster toadfish	1	0	0.5	0
striped searobin	682	0.5	277.5	2.5	yellowtail flounder	1	0	0.1	0
alewife	386	0.3	24.6	0.2	Total	129,609		10,966.8	
Atlantic menhaden	318	0.2	41.9	0.4					
blueback herring	255	0.2	7.5	0.1	<u>Invertebrates</u>				
fourbeard rockling	169	0.1	14.7	0.1	American lobster	9,944	29.3	2,141.9	55.1
smooth dogfish	168	0.1	566.8	5.2	long-finned squid	23,974	70.7	796.4	20.5
striped bass	165	0.1	185.3	1.7	lady crab	nc	nc	535.0	13.8
summer flounder	121	0.1	79.6	0.7	horseshoe crab	nc	nc	116.8	3
American sand lance	95	0.1	0.4	0	spider crab	nc	nc	95.4	2.5
spotted hake	72	0.1	6.5	0.1	lion's mane jellyfish	nc	nc	78.3	2
tautog	61	0	95.1	0.9	rock crab	nc	nc	47.0	1.2
cunner	41	0	4.4	0	blue mussel	nc	nc	14.0	0.4
winter skate	41	0	89.2	0.8	flat claw hermit crab	nc	nc	12.8	0.3
Atlantic silverside	39	0	0.9	0	boring sponge	nc	nc	11.2	0.3
moonfish	33	0	2.1	0	whelks	nc	nc	10.8	0.3
yellow jack	32	0	2.1	0	mantis shrimp	nc	nc	8.1	0.2
ocean pout	30	0	6.5	0.1	bluecrab	nc	nc	6.0	0.2
northern kingfish	25	0	2.5	0	northern moon snail	nc	nc	5.8	0.1
smallmouth flounder	19	0	1.2	0	starfish spp.	nc	nc	4.7	0.1
hogchoker	17	0	1.7	0	arks	nc	nc	1.4	0
black sea bass	12	0	4.7	0	hard clams	nc	nc	0.7	0
hickory shad	6	0	2.5	0	purple sea urchin	nc	nc	0.7	0
Atlantic sturgeon	6	0	145.5	1.3	sand shrimp	nc	nc	0.4	0
longhorn sculpin	5	0	1.3	0	ghost shrimp	nc	nc	0.3	0
clearnose skate	4	0	11.0	0.1	mud crabs	nc	nc	0.2	0
goosefish	4	0	3.3	0	common razor clam	nc	nc	0.1	0
rainbow smelt	4	0	0.3	0	shore shrimp	nc	nc	0.1	0
Atlantic tomcod	4	0	0.8	0	Total	33,918		3,888	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1996.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	49,360	37.0	1,844.7	12.4	northern puffer	3	0	0.3	0
winter flounder	22,722	17.0	3,335.0	22.5	rock gunnel	3	0	0.2	0
scup	16,087	12.0	739.4	5.0	short bigeye	3	0	0.3	0
windowpane flounder	14,116	10.6	1,223.6	8.2	Atlantic sturgeon	3	0	19.9	0.1
bluefish	6,705	5.0	1,118.2	7.5	bigeye scad	2	0	0.1	0
weakfish	6,375	4.8	414.9	2.8	grubby	2	0	0.2	0
little skate	6,203	4.6	2,801.8	18.9	sea raven	2	0	1.5	0
fourspot flounder	2,815	2.1	407.2	2.7	Atlantic tomcod	2	0	0.3	0
alewife	1,402	1.0	134.6	0.9	clearnose skate	1	0	1.7	0
striped searobin	1,008	0.8	278.7	1.9	conger eel	1	0	0.1	0
Atlantic herring	972	0.7	189.8	1.3	gizzard shad	1	0	0.1	0
moonfish	921	0.7	11.6	0.1	goosefish	1	0	0.1	0
red hake	872	0.7	95.5	0.6	sea lamprey	1	0	0.7	0
northern searobin	672	0.5	57.4	0.4	spiny dogfish	1	0	2.1	0
American shad	501	0.4	36.2	0.2	white perch	1	0	0.1	0
silver hake	489	0.4	20.0	0.1	Total	133,546		14,835.2	
summer flounder	434	0.3	266.4	1.8					
spotted hake	384	0.3	42.6	0.3	Invertebrates				
smooth dogfish	275	0.2	862.8	5.8	American lobster	9,490	29.5	2,113.5	39.1
striped bass	232	0.2	373.5	2.5	lady crab	nc	nc	1,160.4	21.5
spot	195	0.1	14.1	0.1	long-finned squid	22,720	70.5	720.4	13.3
tautog	136	0.1	225.9	1.5	horseshoe crab	nc	nc	717.0	13.3
fourbeard rockling	109	0.1	8.6	0.1	spider crab	nc	nc	293.9	5.4
blueback herring	97	0.1	6.2	0	rock crab	nc	nc	162.7	3.0
Atlantic menhaden	88	0.1	40.5	0.3	lion's mane jellyfish	nc	nc	42.7	0.8
winter skate	88	0.1	212.7	1.4	blue mussel	nc	nc	42.5	0.8
hogchoker	45	0	5.4	0	flat claw hermit crab	nc	nc	39.4	0.7
smallmouth flounder	41	0	2.3	0	whelks	nc	nc	33.0	0.6
rough scad	35	0	1.5	0	mantis shrimp	nc	nc	20.9	0.4
hickory shad	29	0	10.2	0.1	boring sponge	nc	nc	19.2	0.4
black sea bass	27	0	12.1	0.1	bushy bryozoan	nc	nc	15.2	0.3
ocean pout	26	0	7.2	0	starfish spp.	nc	nc	6.2	0.1
cunner	17	0	2.6	0	arks	nc	nc	4.3	0.1
striped anchovy	11	0	0.2	0	northern moon snail	nc	nc	4.3	0.1
longhorn sculpin	7	0	2.1	0	bluecrab	nc	nc	4.0	0.1
northern kingfish	6	0	0.6	0	hard clams	nc	nc	3.2	0.1
yellow jack	6	0	0.5	0	surf clam	nc	nc	1.4	0
Atlantic mackerel	5	0	0.5	0	mud crabs	nc	nc	0.3	0
planehead filefish	3	0	0.3	0	purple sea urchin	nc	nc	0.1	0
mackerel scad	3	0	0.1	0	Total	32,210		5,405	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1997.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	70,985	50.3	2,017.2	15.5	American sand lance	2	0	0.1	0
winter flounder	14,701	10.4	2,439.4	18.8	short bigeye	2	0	0.2	0
bluefish	10,815	7.7	977.6	7.5	yellow jack	2	0	0.2	0
windowpane flounder	10,324	7.3	986.1	7.6	bigeye scad	1	0	0.1	0
scup	9,582	6.8	530.5	4.1	Atlantic cod	1	0	0.3	0
fourspot flounder	4,122	2.9	615.3	4.7	haddock	1	0	0.1	0
little skate	4,068	2.9	1,945.8	15.0	northern pipefish	1	0	0.1	0
weakfish	3,904	2.8	362.0	2.8	northern puffer	1	0	0.1	0
Atlantic herring	3,455	2.4	515.1	4.0	rougthead stingray	1	0	50.6	0.4
silver hake	1,973	1.4	70.8	0.5	sea lamprey	1	0	0.1	0
alewife	1,194	0.8	81.3	0.6	Atlantic tomcod	1	0	0.1	0
American shad	922	0.7	66.8	0.5	yellowtail flounder	1	0	0.3	0
striped searobin	819	0.6	230.5	1.8	Total	141,040		12,974.6	
red hake	748	0.5	80.5	0.6					
blueback herring	630	0.4	16.5	0.1	<u>Invertebrates</u>				
northern searobin	579	0.4	60.4	0.5	American lobster	16,467	55.3	3,800.9	64.6
summer flounder	486	0.3	326.0	2.5	lady crab	nc	nc	592.5	10.1
striped bass	319	0.2	509.9	3.9	long-finned squid	13,048	43.8	515.2	8.8
moonfish	287	0.2	4.6	0	horseshoe crab	204	0.7	472.4	8.0
fourbeard rockling	199	0.1	17.3	0.1	spider crab	nc	nc	188.3	3.2
tautog	190	0.1	271.8	2.1	rock crab	nc	nc	94.1	1.6
smooth dogfish	167	0.1	527.3	4.1	lion's mane jellyfish	nc	nc	88.0	1.5
Atlantic menhaden	116	0.1	38.5	0.3	bushy bryozoan	nc	nc	28.0	0.5
spotted hake	77	0.1	19.0	0.1	flat claw hermit crab	nc	nc	21.7	0.4
rough scad	65	0	2.0	0	boring sponge	nc	nc	16.5	0.3
smallmouth flounder	58	0	2.4	0	whelks	22	0.1	14.8	0.3
winter skate	48	0	109.7	0.8	bluecrab	33	0.1	13.6	0.2
cunner	43	0	4.1	0	mantis shrimp	nc	nc	9.3	0.2
hickory shad	25	0	9.1	0.1	starfish spp.	nc	nc	7.3	0.1
black sea bass	22	0	10.5	0.1	hard clams	nc	nc	3.8	0.1
hogchoker	15	0	1.8	0	blue mussel	nc	nc	3.5	0.1
ocean pout	15	0	4.8	0	northern moon snail	nc	nc	3.3	0.1
grubby	11	0	0.7	0	northern comb jelly	nc	nc	2.0	0
spot	10	0	1.1	0	arks	nc	nc	1.8	0
Atlantic mackerel	8	0	1.7	0	common oyster	nc	nc	1.8	0
northern kingfish	7	0	0.9	0	surf clam	nc	nc	0.9	0
spiny dogfish	7	0	13.7	0.1	common slipper shell	nc	nc	0.7	0
Atlantic sturgeon	5	0	37.8	0.3	mud crabs	nc	nc	0.6	0
clearnose skate	4	0	7.4	0.1	sand shrimp	nc	nc	0.2	0
longhorn sculpin	4	0	0.8	0	common razor clam	nc	nc	0.2	0
white perch	4	0	0.9	0	blood star	nc	nc	0.1	0
crevalle jack	3	0	0.6	0	star coral	nc	nc	0.1	0
sea raven	3	0	0.4	0	northern red shrimp	nc	nc	0.1	0
Atlantic silverside	2	0	0.1	0	shore shrimp	nc	nc	0.1	0
goosefish	2	0	1.6	0	purple sea urchin	nc	nc	0.1	0
inshore lizardfish	2	0	0.2	0	Total	29,774		5,882	
round scad	2	0	0.2	0					

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1998.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	136,926	64.0	3,661.1	24.4	goosefish	3	0	3.2	0
scup	23,742	11.1	740.5	4.9	oyster toadfish	3	0	0.9	0
winter flounder	15,697	7.3	2,450.3	16.3	gray triggerfish	2	0	2.3	0
bluefish	8,814	4.1	899.0	6.0	longhorn sculpin	2	0	1.0	0
windowpane flounder	6,483	3.0	741.1	4.9	bigeye scad	1	0	0.1	0
little skate	4,305	2.0	2,085.5	13.9	inshore lizardfish	1	0	0.1	0
weakfish	3,495	1.6	268.2	1.8	mackerel scad	1	0	0.1	0
red hake	3,015	1.4	217.5	1.4	rougtail stingray	1	0	3.4	0
fourspot flounder	1,908	0.9	306.0	2.0	Total	214,025		15,005.7	
silver hake	1,870	0.9	88.3	0.6					
striped searobin	1,321	0.6	509.7	3.4	Invertebrates				
moonfish	1,188	0.6	13.4	0.1	American lobster	16,211	36.7	3,873.9	60.2
American shad	901	0.4	60.2	0.4	long-finned squid	27,443	62.1	767.0	11.9
Atlantic herring	893	0.4	74.6	0.5	horseshoe crab	303	0.7	489.4	7.6
alewife	456	0.2	35.1	0.2	blue mussel	nc	nc	309.0	4.8
summer flounder	436	0.2	431.3	2.9	lady crab	nc	nc	291.2	4.5
striped bass	400	0.2	484.2	3.2	rock crab	nc	nc	241.4	3.8
northern searobin	360	0.2	39.4	0.3	spider crab	nc	nc	157.2	2.4
smooth dogfish	310	0.1	989.8	6.6	lion's mane jellyfish	nc	nc	63.1	1.0
Atlantic menhaden	306	0.1	9.2	0.1	flat claw hermit crab	nc	nc	56.0	0.9
blueback herring	211	0.1	5.1	0	bushy bryozoan	nc	nc	55.6	0.9
tautog	194	0.1	347.1	2.3	boring sponge	nc	nc	24.9	0.4
spotted hake	142	0.1	12.2	0.1	knobbed whelk	51	0.1	22.5	0.3
fourbeard rockling	133	0.1	11.6	0.1	starfish spp.	nc	nc	18.2	0.3
smallmouth flounder	97	0	6.4	0	bluecrab	49	0.1	12.8	0.2
cunner	65	0	8.1	0.1	channeled whelk	40	0.1	10.1	0.2
winter skate	62	0	180.7	1.2	whelks	52	0.1	9.8	0.2
hickory shad	40	0	15.9	0.1	northern moon snail	nc	nc	8.6	0.1
round herring	31	0	0.6	0	mantis shrimp	nc	nc	5.6	0.1
sea raven	30	0	11.3	0.1	common oyster	nc	nc	5.4	0.1
northern puffer	28	0	0.5	0	hard clams	nc	nc	3.7	0.1
clearnose skate	20	0	36.8	0.2	arks	nc	nc	2.0	0
black sea bass	18	0	10.6	0.1	red bearded sponge	nc	nc	1.4	0
spiny dogfish	18	0	44.5	0.3	surf clam	nc	nc	1.1	0
Atlantic sturgeon	17	0	189.7	1.3	sea grape	nc	nc	0.8	0
northern kingfish	15	0	1.3	0	mud crabs	nc	nc	0.7	0
Atlantic mackerel	13	0	1.1	0	boreal squid	18	0	0.7	0
ocean pout	13	0	2.7	0	purple sea urchin	nc	nc	0.6	0
hogchoker	12	0	1.9	0	common slipper shell	nc	nc	0.5	0
haddock	7	0	0.5	0	star coral	nc	nc	0.4	0
yellow jack	6	0	0.7	0	moon jelly	nc	nc	0.2	0
grubby	5	0	0.3	0	ghost shrimp	nc	nc	0.1	0
round scad	4	0	0.3	0	Total	44,167		6,434	
American sand lance	4	0	0.3	0					

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 1999.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	191,100	54.1	4,171.6	21.9	goosefish	2	0	0.3	0
scup	101,095	28.6	3,641.3	19.1	grubby	2	0	0.2	0
weakfish	12,416	3.5	771.3	4.0	northern pipefish	2	0	0.1	0
winter flounder	10,288	2.9	2,011.7	10.6	longhorn sculpin	2	0	0.3	0
bluefish	7,843	2.2	1,218.0	6.4	oyster toadfish	2	0	1.8	0
silver hake	5,126	1.5	99.6	0.5	Atlantic silverside	1	0	0.1	0
windowpane flounder	4,643	1.3	594.2	3.1	gizzard shad	1	0	0.1	0
little skate	3,686	1.0	1,829.6	9.6	haddock	1	0	0.1	0
red hake	2,973	0.8	226.5	1.2	round scad	1	0	0.1	0
Atlantic herring	2,511	0.7	45.4	0.2	striped cusk-eel	1	0	0.1	0
striped searobin	1,690	0.5	497.0	2.6	sharksucker	1	0	0.3	0
alewife	1,393	0.4	107.6	0.6	Spanish mackerel	1	0	0.2	0
fourspot flounder	1,393	0.4	203.9	1.1	Atlantic tomcod	1	0	0.7	0
Atlantic menhaden	1,187	0.3	90.9	0.5	white perch	1	0	0.4	0
American shad	987	0.3	117.3	0.6	Total	353,203		19,054.7	
moonfish	645	0.2	9.6	0.1					
summer flounder	582	0.2	459.8	2.4					
bay anchovy	548	0.2	5.6	0	Invertebrates				
northern searobin	547	0.2	52.0	0.3	American lobster	13,922	38.1	3,397.9	61.6
striped bass	397	0.1	815.4	4.3	long-finned squid	21,580	59.0	826.4	15.0
spotted hake	381	0.1	38.8	0.2	horseshoe crab	384	1.1	634.1	11.5
smooth dogfish	305	0.1	923.0	4.8	lady crab	nc	nc	159.7	2.9
fourbeard rockling	233	0.1	28.8	0.2	rock crab	nc	nc	118.6	2.2
tautog	217	0.1	326.6	1.7	spider crab	nc	nc	95.4	1.7
striped anchovy	216	0.1	6.1	0	bushy bryozoan	nc	nc	78.0	1.4
American sand lance	178	0.1	0.3	0	flat claw hermit crab	nc	nc	32.5	0.6
smallmouth flounder	96	0	5.2	0	knobbed whelk	61	0.2	24.8	0.4
hickory shad	56	0	19.4	0.1	bluecrab	89	0.2	21.3	0.4
cunner	51	0	5.9	0	channeled whelk	81	0.2	21.1	0.4
black sea bass	50	0	17.2	0.1	mantis shrimp	376	1.0	19.3	0.4
spot	45	0	5.7	0	boring sponge	nc	nc	19.3	0.4
winter skate	41	0	89.8	0.5	lion's mane jellyfish	61	0.2	16.7	0.3
hogchoker	39	0	5.0	0	blue mussel	nc	nc	14.1	0.3
Atlantic sturgeon	39	0	498.6	2.6	northern moon snail	nc	nc	9.1	0.2
clearnose skate	22	0	39.4	0.2	starfish spp.	nc	nc	8.8	0.2
bigeye scad	21	0	1.4	0	common oyster	nc	nc	4.7	0.1
Atlantic mackerel	21	0	3.1	0	arks	nc	nc	2.8	0.1
yellow jack	20	0	1.9	0	common slipper shell	nc	nc	1.8	0
blueback herring	19	0	1.1	0	mud crabs	nc	nc	1.7	0
ocean pout	17	0	3.9	0	hard clams	nc	nc	1.5	0
northern puffer	14	0	1.1	0	sand shrimp	nc	nc	1.0	0
spiny dogfish	10	0	51.1	0.3	purple sea urchin	nc	nc	1.0	0
sea raven	9	0	4.9	0	northern red shrimp	nc	nc	0.9	0
crevalle jack	8	0	0.7	0	surf clam	nc	nc	0.4	0
inshore lizardfish	7	0	0.5	0	sea grape	nc	nc	0.2	0
northern kingfish	6	0	0.6	0	star coral	nc	nc	0.1	0
northern sennet	6	0	0.5	0	common razor clam	nc	nc	0.1	0
planehead filefish	3	0	0.3	0	moon jelly	nc	nc	0.1	0
bigeye	2	0	0.2	0	nemerteans	nc	nc	0.1	0
conger eel	2	0	0.5	0	Total	36,554		5,514	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 2000.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
scup	101,464	44.4	6,679.0	34.9	northern kingfish	2	0	0.3	0
butterfish	60,490	26.5	1,458.3	7.6	round scad	2	0	0.2	0
weakfish	23,595	10.3	554.5	2.9	bigeye	1	0	0.1	0
winter flounder	8,867	3.9	1,921.4	10.0	Atlantic cod	1	0	0.1	0
bluefish	6,135	2.7	1,408.0	7.3	goosefish	1	0	0.2	0
little skate	3,340	1.5	1,604.7	8.4	inshore lizardfish	1	0	0.1	0
striped searobin	3,129	1.4	1,036.1	5.4	lined seahorse	1	0	0.1	0
fourspot flounder	2,590	1.1	398.6	2.1	white perch	1	0	0.2	0
windowpane flounder	2,488	1.1	368.8	1.9	yellowtail flounder	1	0	0.1	0
red hake	2,393	1.0	162.6	0.8	Total	228,425		19,156.5	
bay anchovy	2,303	1.0	12.2	0.1					
northern searobin	2,014	0.9	251.2	1.3	Invertebrates				
moonfish	1,817	0.8	15.0	0.1	American lobster	10,481	36.0	2,184.5	49.9
alewife	1,572	0.7	96.0	0.5	horseshoe crab	420	1.4	689.4	15.8
spotted hake	1,425	0.6	92.3	0.5	long-finned squid	16,585	57.0	582.3	13.3
Atlantic herring	770	0.3	124.1	0.6	lady crab	nc	nc	308.4	7.1
silver hake	679	0.3	28.8	0.2	spider crab	nc	nc	99.4	2.3
summer flounder	555	0.2	471.3	2.5	bushy bryozoan	nc	nc	95.2	2.2
Atlantic menhaden	492	0.2	31.8	0.2	rock crab	nc	nc	60.4	1.4
smooth dogfish	467	0.2	1,038.5	5.4	boring sponge	nc	nc	58.6	1.3
American shad	316	0.1	25.8	0.1	mantis shrimp	1,086	3.7	49.0	1.1
striped bass	293	0.1	602.6	3.1	blue mussel	nc	nc	36.8	0.8
tautog	287	0.1	463.5	2.4	lion's mane jellyfish	223	0.8	36.4	0.8
spot	204	0.1	17.8	0.1	channeled whelk	138	0.5	32.0	0.7
fourbeard rockling	185	0.1	14.7	0.1	knobbed whelk	76	0.3	29.9	0.7
blueback herring	143	0.1	6.8	0	starfish spp.	nc	nc	29.0	0.7
black sea bass	69	0	22.6	0.1	flat claw hermit crab	nc	nc	26.0	0.6
smallmouth flounder	61	0	2.7	0	bluecrab	104	0.4	19.3	0.4
cunner	50	0	5.3	0	northern moon snail	nc	nc	9.7	0.2
hickory shad	42	0	17.1	0.1	hydroid spp.	nc	nc	4.8	0.1
hogchoker	40	0	5.9	0	fan worm tubes	nc	nc	3.4	0.1
winter skate	31	0	66.5	0.3	hard clams	nc	nc	3.3	0.1
sea raven	19	0	9.2	0	arks	nc	nc	3.1	0.1
clearnose skate	18	0	37.9	0.2	mud crabs	nc	nc	2.8	0.1
ocean pout	18	0	4.9	0	sand shrimp	nc	nc	2.7	0.1
longhorn sculpin	14	0	5.0	0	common slipper shell	nc	nc	2.4	0.1
Atlantic sturgeon	7	0	79.0	0.4	purple sea urchin	nc	nc	2.3	0.1
oyster toadfish	6	0	2.5	0	common oyster	nc	nc	1.4	0
northern pipefish	4	0	0.2	0	sea grape	nc	nc	1.1	0
northern puffer	4	0	0.4	0	blood star	nc	nc	0.2	0
American sand lance	4	0	0.3	0	northern comb jelly	nc	nc	0.1	0
spiny dogfish	4	0	9.9	0.1	common razor clam	nc	nc	0.1	0
rock gunnel	3	0	0.2	0	northern cyclocardia	nc	nc	0.1	0
yellow jack	3	0	0.2	0	northern red shrimp	nc	nc	0.1	0
Atlantic silverside	2	0	0.1	0	surf clam	nc	nc	0.1	0
Atlantic mackerel	2	0	0.8	0	Total	29,113		4,374	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 2001.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay anchovy, striped anchovy, and American sand lance and Atlantic herring are not quantified. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
scup	58,325	37.7	5,828.4	30.7	American eel	1	0	0.6	0
butterfish	45,264	29.3	1,834.0	9.7	planehead filefish	1	0	0.1	0
weakfish	12,739	8.2	415.0	2.2	goosefish	1	0	0.4	0
winter flounder	9,826	6.4	1,993.6	10.5	naked goby	1	0	0.1	0
little skate	4,311	2.8	2,022.6	10.6	northern sennet	1	0	0.1	0
bluefish	3,986	2.6	751.2	4.0	rock gunnel	1	0	0.1	0
silver hake	3,945	2.6	152.2	0.8	red goatfish	1	0	0.1	0
windowpane flounder	3,065	2.0	475.5	2.5	rougtail stingray	1	0	2.5	0
fourspot flounder	2,167	1.4	362.7	1.9	short bigeye	1	0	0.1	0
striped searobin	2,061	1.3	861.0	4.5	yellowtail flounder	1	0	0.2	0
northern searobin	1,594	1.0	222.7	1.2	Total	154,514		18,997.8	
red hake	1,382	0.9	109.7	0.6					
summer flounder	875	0.6	628.1	3.3	<u>Finfish not ranked</u>				
alewife	638	0.4	41.7	0.2	American sand lance, yoy				
spotted hake	606	0.4	34.9	0.2	anchovy spp, yoy				
smooth dogfish	598	0.4	1,407.6	7.4	Atlantic herring, yoy				
Atlantic herring	497	0.3	72.6	0.4					
bay anchovy	443	0.3	3.6	0	<u>Invertebrates</u>				
tautog	319	0.2	491.2	2.6	American lobster	5,626	35.1	1,531.2	39.2
blueback herring	279	0.2	11.1	0.1	horseshoe crab	503	3.1	870.7	22.3
fourbeard rockling	251	0.2	21.5	0.1	long-finned squid	9,080	56.6	346.2	8.9
moonfish	225	0.1	3.8	0	spider crab	nc	nc	302.5	7.7
striped bass	214	0.1	472.5	2.5	bushy bryozoan	nc	nc	162.9	4.2
black sea bass	134	0.1	74.8	0.4	starfish spp.	nc	nc	154.7	4.0
American shad	109	0.1	9.6	0.1	rock crab	nc	nc	86.3	2.2
smallmouth flounder	98	0.1	3.8	0	blue mussel	nc	nc	84.7	2.2
Atlantic menhaden	86	0.1	4.7	0	lady crab	nc	nc	79.0	2.0
hogchoker	85	0.1	10.5	0.1	flat claw hermit crab	nc	nc	57.6	1.5
clearnose skate	65	0	132.4	0.7	knobbed whelk	118	0.7	53.3	1.4
cunner	51	0	5.9	0	channeled whelk	190	1.2	48.0	1.2
spiny dogfish	48	0	128.6	0.7	boring sponge	nc	nc	30.0	0.8
striped anchovy	47	0	1.2	0	lion's mane jellyfish	182	1.1	25.9	0.7
winter skate	38	0	112.2	0.6	northern moon snail	nc	nc	17.5	0.4
inshore lizardfish	21	0	2.2	0	mantis shrimp	304	1.9	16.5	0.4
Atlantic sturgeon	18	0	270.6	1.4	bluecrab	38	0.2	6.2	0.2
hickory shad	14	0	6.7	0	sea grape	nc	nc	6.1	0.2
spot	13	0	1.3	0	common slipper shell	nc	nc	5.3	0.1
rough scad	10	0	0.7	0	hydroid spp.	nc	nc	5.0	0.1
northern puffer	8	0	0.7	0	arks	nc	nc	4.0	0.1
sea raven	7	0	4.1	0	mud crabs	nc	nc	3.6	0.1
ocean pout	6	0	2.3	0	hard clams	nc	nc	3.0	0.1
round herring	5	0	0.1	0	sand shrimp	nc	nc	2.8	0.1
longhorn sculpin	5	0	1.5	0	common oyster	1	0	1.2	0
fawn cusk-eel	4	0	0.2	0	fan worm tubes	nc	nc	1.0	0
northern pipefish	4	0	0.3	0	purple sea urchin	nc	nc	0.8	0
American sand lance	4	0	0.3	0	moon jelly	nc	nc	0.4	0
seasnail	4	0	0.3	0	ghost shrimp	nc	nc	0.3	0
yellow jack	3	0	0.3	0	bobtail squid	1	0	0.1	0
conger eel	2	0	0.3	0	common razor clam	nc	nc	0.1	0
northern kingfish	2	0	0.2	0	northern red shrimp	nc	nc	0.1	0
oyster toadfish	2	0	0.4	0	surf clam	nc	nc	0.1	0
Atlantic silverside	1	0	0.1	0	Total	16,043		3,907	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 2002.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring are not quantified. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
scup	100,481	47.0	13,814.1	46.0	inshore lizardfish	1	0	0.1	0
butterfish	66,550	31.1	1,924.2	6.4	northern kingfish	1	0	0.2	0
weakfish	10,713	5.0	442.0	1.5	rock gunnel	1	0	0.1	0
winter flounder	6,884	3.2	1,584.1	5.3	rainbow smelt	1	0	0.1	0
little skate	4,242	2.0	2,121.9	7.1	rougtail stingray	1	0	24.4	0.1
bluefish	3,450	1.6	1,099.7	3.7	Total	213,796		30,062.0	
striped searobin	2,394	1.1	1,065.0	3.5					
northern searobin	2,123	1.0	267.3	0.9	Finfish not ranked				
red hake	2,103	1.0	206.6	0.7	anchovy spp, yoy				
silver hake	2,013	0.9	89.6	0.3	Atlantic herring, yoy				
windowpane flounder	1,991	0.9	343.3	1.1					
fourspot flounder	1,859	0.9	326.9	1.1	Invertebrates				
summer flounder	1,356	0.6	989.3	3.3	blue mussel	nc	nc	2,497.8	43.9
smooth dogfish	1,019	0.5	2,814.3	9.4	American lobster	3,880	29.7	1,005.7	17.7
bay anchovy	992	0.5	6.6	0	horseshoe crab	517	4.0	862.9	15.2
alewife	855	0.4	70.2	0.2	spider crab	nc	nc	348.4	6.1
spotted hake	798	0.4	48.2	0.2	long-finned squid	8,034	61.5	279.9	4.9
American shad	593	0.3	40.3	0.1	lady crab	nc	nc	117.0	2.1
tautog	565	0.3	921.1	3.1	starfish spp.	nc	nc	91.8	1.6
striped bass	469	0.2	855.2	2.8	bushy bryozoan	nc	nc	85.0	1.5
moonfish	424	0.2	7.4	0	boring sponge	nc	nc	83.9	1.5
black sea bass	394	0.2	188.3	0.6	rock crab	nc	nc	74.6	1.3
Atlantic menhaden	366	0.2	96.3	0.3	flat claw hermit crab	36	0.3	55.8	1.0
Atlantic herring	365	0.2	63.9	0.2	channeled whelk	174	1.3	43.6	0.8
smallmouth flounder	139	0.1	4.9	0	northern moon snail	nc	nc	40.3	0.7
fourbeard rockling	106	0	9.7	0	knobbed whelk	40	0.3	19.1	0.3
hogchoker	100	0	13.3	0	bluecrab	84	0.6	16.1	0.3
blueback herring	68	0	2.4	0	lion's mane jellyfish	71	0.5	12.3	0.2
clearnose skate	59	0	107.3	0.4	mantis shrimp	226	1.7	11.2	0.2
cunner	55	0	7.2	0	arks	nc	nc	7.8	0.1
spot	52	0	7.2	0	common slipper shell	nc	nc	7.3	0.1
hickory shad	45	0	19.6	0.1	hydroid spp.	nc	nc	7.3	0.1
winter skate	45	0	133.5	0.4	sea grape	nc	nc	5.3	0.1
Atlantic sturgeon	18	0	275.3	0.9	hard clams	3	0	5.2	0.1
spiny dogfish	17	0	48.0	0.2	mud crabs	nc	nc	4.7	0.1
ocean pout	13	0	4.3	0	purple sea urchin	nc	nc	2.3	0
yellow jack	13	0	1.4	0	sand shrimp	nc	nc	1.6	0
sea raven	11	0	4.1	0	rubbery bryozoan	nc	nc	1.0	0
rough scad	10	0	0.7	0	surf clam	nc	nc	1.0	0
oyster toadfish	8	0	4.7	0	deadman's fingers sponge	nc	nc	0.5	0
northern puffer	6	0	0.3	0	blood star	nc	nc	0.4	0
Atlantic mackerel	5	0	2.5	0	common oyster	nc	nc	0.4	0
short bigeye	5	0	0.2	0	mixed sponge species	nc	nc	0.4	0
goosefish	3	0	0.6	0	northern red shrimp	nc	nc	0.3	0
American sand lance	3	0	0.1	0	anemones	nc	nc	0.1	0
longhorn sculpin	3	0	0.9	0	bobtail squid	1	0	0.1	0
northern sennet	2	0	0.2	0	ghost shrimp	nc	nc	0.1	0
northern pipefish	2	0	0.2	0	ribbed mussel	nc	nc	0.1	0
Atlantic bonito	1	0	2.4	0	sea cucumber	1	0	0.1	0
crevalle jack	1	0	0.1	0	Total	13,067		5,691	
gizzard shad	1	0	0.1	0					
grubby	1	0	0.1	0					

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 2003.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring are not quantified. Number of tows (sample size)=160.

species	count	%	weight	%	Species	count	%	weight	%
butterfish	25,483	34.4	524.6	3.7	barndoor skate	1	0	0.4	0
scup	17,552	23.7	4,389.3	30.6	Planehead filefish	1	0	0.1	0
weakfish	5,596	7.6	131.9	0.9	rainbow smelt	1	0	0.1	0
winter flounder	4,245	5.7	1,276.5	8.9	sea lamprey	1	0	1.3	0
bluefish	3,717	5.0	655.0	4.6	Spanish mackerel	1	0	2.1	0
little skate	2,867	3.9	1,554.1	10.8	Total	74,107		14,323.6	
bay anchovy	2,254	3.0	12.5	0.1					
windowpane flounder	1,858	2.5	333.9	2.3	Finfish not ranked				
fourspot flounder	1,658	2.2	327.7	2.3	anchovy spp, yoy				
striped searobin	1,529	2.1	687.0	4.8	Atlantic herring, yoy				
northern searobin	1,468	2.0	240.7	1.7					
summer flounder	1,151	1.6	825.0	5.8	Invertebrates				
red hake	681	0.9	31.1	0.2	Horseshoe crab	399	1.7	670.5	23.2
alewife	608	0.8	49.4	0.3	spider crab	nc	nc	640.6	22.2
smooth dogfish	552	0.7	1,508.8	10.5	American lobster	1,958	8.3	479.7	16.6
spotted hake	527	0.7	41.6	0.3	long-finned squid	19,231	81.9	421.3	14.6
Atlantic herring	448	0.6	87.8	0.6	boring sponge	nc	nc	107.5	3.7
American shad	305	0.4	23.5	0.2	rock crab	nc	nc	80.9	2.8
silver hake	217	0.3	8.3	0.1	starfish spp.	nc	nc	73.7	2.6
striped bass	215	0.3	542.1	3.8	flat claw hermit crab	nc	nc	61.3	2.1
tautog	210	0.3	325.4	2.3	channeled whelk	334	1.4	58.8	2.0
Atlantic menhaden	121	0.2	16.1	0.1	bushy bryozoan	nc	nc	54.3	1.9
fourbeard rockling	111	0.1	9.0	0.1	lion's mane jellyfish	1,307	5.6	40.6	1.4
blueback herring	98	0.1	3.4	0	knobbed whelk	96	0.4	35.1	1.2
moonfish	97	0.1	1.3	0	sea grape	nc	nc	31.1	1.1
hogchoker	89	0.1	8.3	0.1	northern moon snail	nc	nc	20.9	0.7
black sea bass	57	0.1	45.7	0.3	blue mussel	nc	nc	19.7	0.7
Atlantic cod	57	0.1	2.7	0	common slipper shell	nc	nc	16.8	0.6
clearnose skate	55	0.1	105.9	0.7	lady crab	nc	nc	12.0	0.4
smallmouth flounder	38	0.1	2.4	0	hydroid spp.	nc	nc	9.6	0.3
winter skate	38	0.1	90.6	0.6	ribbed mussel	nc	nc	8.8	0.3
cunner	36	0	5.9	0	sand shrimp	nc	nc	6.8	0.2
haddock	26	0	1.3	0	arks	nc	nc	6.5	0.2
Atlantic sturgeon	23	0	391.9	2.7	mud crabs	nc	nc	6.5	0.2
hickory shad	22	0	10.3	0.1	rubbery bryozoan	nc	nc	6.0	0.2
American sand lance	19	0	0.2	0	mantis shrimp	110	0.5	4.9	0.2
ocean pout	14	0	2.9	0	bluecrab	24	0.1	4.3	0.1
rough scad	12	0	0.5	0	hard clams	nc	nc	3.9	0.1
oyster toadfish	9	0	5.0	0	star coral	nc	nc	1.9	0.1
spiny dogfish	7	0	34.8	0.2	coastal mud shrimp	4	0	0.7	0
rock gunnel	6	0	0.4	0	purple sea urchin	nc	nc	0.6	0
round scad	4	0	0.3	0	blood star	nc	nc	0.4	0
glasseye snapper	3	0	0.1	0	northern red shrimp	2	0	0.4	0
conger eel	3	0	1.1	0	Japanese shore crab	4	0	0.3	0
Atlantic mackerel	3	0	0.3	0	anemones	nc	nc	0.1	0
crevalle jack	2	0	0.2	0	sand dollar	1	0	0.1	0
northern pipefish	2	0	0.2	0	common razor clam	1	0	0.1	0
northern puffer	2	0	0.2	0	moon jelly	nc	nc	0.1	0
longhorn sculpin	2	0	0.9	0	northern cyclocardia	nc	nc	0.1	0
sea raven	2	0	1.3	0	mixed sponge species	nc	nc	0.1	0
striped anchovy	2	0	0.1	0	Total	23,471		2,887	
Atlantic silverside	1	0	0.1	0					

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 2004.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring are not quantified. Number of tows (sample size)=199.

species	count	%	weight	%	species	count	%	weight	%
butterfish	94,735	46.7	1,842.7	9.7	American plaice	1	0	0.1	0
scup	61,521	30.3	6,801.1	35.7	conger eel	1	0	0.1	0
weakfish	17,505	8.6	426.9	2.2	gizzard shad	1	0	0.1	0
bluefish	6,504	3.2	2,140.6	11.2	goosefish	1	0	0.1	0
winter flounder	4,021	2.0	839.9	4.4	pollock	1	0	0.1	0
little skate	3,044	1.5	1,689.8	8.9	rougtail stingray	1	0	4.1	0
windowpane flounder	2,275	1.1	333.7	1.8	oyster toadfish	1	0	0.8	0
bay anchovy	1,523	0.8	10.3	0.1	yellow jack	1	0	0.1	0
silver hake	1,417	0.7	27.3	0.1	Total	202,887		19,056.6	
fourspot flounder	1,406	0.7	309.3	1.6					
striped searobin	1,308	0.6	465.4	2.4	<u>Finfish not ranked</u>				
alewife	859	0.4	56.1	0.3	anchovy spp, yoy				
Atlantic herring	851	0.4	58.3	0.3	Atlantic herring, yoy				
red hake	829	0.4	51.6	0.3					
northern searobin	784	0.4	112.0	0.6	<u>Invertebrates</u>				
Atlantic menhaden	746	0.4	110.7	0.6	long-finned squid	23,022	86.5	953.4	28.8
summer flounder	644	0.3	627.2	3.3	horseshoe crab	534	2.0	873.4	26.4
smooth dogfish	503	0.2	1,435.3	7.5	American lobster	1,843	6.9	481.5	14.5
striped bass	378	0.2	811.8	4.3	spider crab	nc	nc	355.5	10.7
American shad	356	0.2	24.2	0.1	blue mussel	nc	nc	250.2	7.6
tautog	232	0.1	353.7	1.9	bushy bryozoan	nc	nc	50.9	1.5
spotted hake	230	0.1	37.8	0.2	flat claw hermit crab	nc	nc	42.4	1.3
blueback herring	218	0.1	6.5	0	channeled whelk	199	0.7	42.3	1.3
moonfish	182	0.1	3.4	0	starfish spp.	nc	nc	41.7	1.3
fourbeard rockling	173	0.1	13.0	0.1	boring sponge	nc	nc	41.7	1.3
black sea bass	124	0.1	40.5	0.2	rock crab	1	0.0	35.2	1.1
hogchoker	83	0	9.5	0	lion's mane jellyfish	803	3.0	34.0	1.0
American sand lance	70	0	0.2	0	common slipper shell	nc	nc	22.9	0.7
winter skate	53	0	100.3	0.5	sea grape	nc	nc	16.4	0.5
smallmouth flounder	50	0	2.8	0	lady crab	nc	nc	14.5	0.4
hickory shad	39	0	14.2	0.1	northern moon snail	nc	nc	11.5	0.3
spiny dogfish	38	0	104.7	0.5	knobbed whelk	21	0.1	7.7	0.2
Atlantic cod	33	0	4.7	0	mantis shrimp	159	0.6	7.0	0.2
clearnose skate	22	0	48.2	0.3	arks	nc	nc	7.0	0.2
cunner	21	0	3.7	0	mud crabs	nc	nc	5.4	0.2
ocean pout	18	0	5.4	0	sand shrimp	nc	nc	4.7	0.1
rough scad	14	0	0.7	0	bluecrab	13	0	2.8	0.1
round scad	11	0	0.3	0	hard clams	nc	nc	2.3	0.1
spot	8	0	0.9	0	surf clam	5	0	1.0	0
Atlantic sturgeon	8	0	117.6	0.6	purple sea urchin	nc	nc	0.8	0
haddock	7	0	0.6	0	mixed sponge species	nc	nc	0.6	0
sea raven	7	0	2.4	0	hydroid spp.	nc	nc	0.6	0
northern kingfish	5	0	0.5	0	deadman's fingers sponge	nc	nc	0.5	0
northern puffer	5	0	0.4	0	rubbery bryozoan	nc	nc	0.4	0
longhorn sculpin	5	0	3.4	0	star coral	nc	nc	0.3	0
seasnail	4	0	0.2	0	northern red shrimp	nc	nc	0.3	0
crevalle jack	2	0	0.2	0	northern cyclocardia	nc	nc	0.2	0
northern pipefish	2	0	0.2	0	blood star	nc	nc	0.1	0
rock gunnel	2	0	0.2	0	coastal mud shrimp	1	0	0.1	0
Atlantic tomcod	2	0	0.2	0	sea cucumber	2	0	0.1	0
white perch	2	0	0.5	0	Total	26,603		3,309.4	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 2005.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring are not quantified. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	92,996	52.2	2,097.3	16.8	haddock	2	0	0.2	0
scup	52,642	29.6	3,080.7	24.7	seasnail	2	0	0.2	0
weakfish	9,191	5.2	449.9	3.6	glasseye snapper	1	0	0.1	0
bluefish	6,532	3.7	1,333.8	10.7	inshore lizardfish	1	0	0.1	0
winter flounder	4,692	2.6	566.1	4.5	lookdown	1	0	0.1	0
windowpane flounder	1,982	1.1	177.5	1.4	pollock	1	0	0.1	0
little skate	1,317	0.7	682.5	5.5	Total	178,073		12,474.3	
Atlantic herring	1,168	0.7	131.1	1.1					
bay anchovy	814	0.5	5.8	0	Finfish not ranked				
striped searobin	757	0.4	183.7	1.5	anchovy spp, yoy				
alewife	742	0.4	47.6	0.4	Atlantic herring, yoy				
fourspot flounder	688	0.4	125.9	1					
red hake	585	0.3	56.0	0.4	Invertebrates				
summer flounder	506	0.3	406.1	3.3	blue mussel	nc	nc	971.0	32.6
striped bass	469	0.3	675.1	5.4	long-finned squid	17,542	83.2	683.5	22.9
smooth dogfish	467	0.3	1,421.7	11.4	American lobster	1,389	6.6	364.3	12.2
moonfish	356	0.2	6.0	0	horseshoe crab	161	0.8	304.2	10.2
northern searobin	265	0.1	21.3	0.2	starfish spp.	nc	nc	198.4	6.7
Atlantic menhaden	235	0.1	77.9	0.6	lion's mane jellyfish	1,806	8.6	97.3	3.3
spotted hake	234	0.1	17.4	0.1	spider crab	nc	nc	92.0	3.1
tautog	179	0.1	269.2	2.2	bushy bryozoan	nc	nc	64.6	2.2
American shad	177	0.1	18.2	0.1	lady crab	nc	nc	48.8	1.6
silver hake	165	0.1	7.1	0.1	boring sponge	nc	nc	26.1	0.9
hickory shad	136	0.1	43.1	0.3	flat claw hermit crab	nc	nc	23.1	0.8
blueback herring	111	0.1	5.4	0	channeled whelk	101	0.5	23.0	0.8
fourbeard rockling	106	0.1	6.8	0.1	common slipper shell	nc	nc	12.2	0.4
clearnose skate	102	0.1	187.1	1.5	rubbery bryozoan	nc	nc	11.0	0.4
rough scad	62	0	1.9	0	knobbed whelk	23	0.1	9.7	0.3
hogchoker	61	0	8.7	0.1	rock crab	nc	nc	9.3	0.3
smallmouth flounder	44	0	2.4	0	ribbed mussel	nc	nc	7.6	0.3
black sea bass	42	0	26.4	0.2	hard clams	nc	nc	7.2	0.2
spiny dogfish	41	0	102.0	0.8	northern moon snail	nc	nc	4.7	0.2
Atlantic mackerel	37	0	5.7	0	sea grape	nc	nc	4.5	0.2
winter skate	31	0	59.9	0.5	mantis shrimp	64	0.3	3.8	0.1
yellow jack	28	0	3.0	0	arks	nc	nc	3.5	0.1
cunner	24	0	4.1	0	hydroid spp.	nc	nc	3.4	0.1
round scad	12	0	0.3	0	mud crabs	nc	nc	2.5	0.1
Atlantic cod	10	0	0.9	0	sand shrimp	nc	nc	2.1	0.1
rock gunnel	9	0	0.6	0	deadman's fingers sponge	nc	nc	1.1	0
Atlantic sturgeon	9	0	152.7	1.2	purple sea urchin	nc	nc	0.7	0
northern sennet	8	0	0.7	0	bluecrab	3	0	0.6	0
American sand lance	6	0	0.2	0	mixed sponge species	nc	nc	0.4	0
northern puffer	5	0	0.3	0	surf clam	nc	nc	0.4	0
northern kingfish	4	0	0.6	0	star coral	nc	nc	0.3	0
northern pipefish	4	0	0.3	0	sand dollar	1	0	0.2	0
ocean pout	3	0	0.7	0	northern red shrimp	nc	nc	0.2	0
sea raven	3	0	0.5	0	boreal squid	1	0	0.1	0
crevalle jack	2	0	0.2	0	Japanese shore crab	5	0	0.1	0
gizzard shad	2	0	0.2	0	northern cyclocardia	nc	nc	0.1	0
goosefish	2	0	0.7	0	common oyster	nc	nc	0.1	0
grubby	2	0	0.2	0	Total	21,096		2,982.1	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in LISTS in 2006.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring and American sand lance are not quantified. Number of tows (sample size)=120.

species	count	%	weight	%	species	count	%	weight	%
butterfish	50,022	54.3	1,631.4	15.5					
scup	28,829	31.3	4,636.1	44.2					
bluefish	2,100	2.3	358.6	3.4	<u>Finfish not ranked</u>				
winter flounder	1,699	1.8	271.2	2.6	anchovy spp. yoy				
bay anchovy	1,492	1.6	8.3	0.1	Atlantic herring, yoy				
silver hake	1,267	1.4	37.7	0.4	American sand lance (yoy)				
windowpane flounder	1,077	1.2	128.9	1.2					
northern searobin	630	0.7	74.5	0.7					
red hake	625	0.7	37.4	0.4					
little skate	593	0.6	310.6	3	<u>Invertebrates</u>				
alewife	573	0.6	49.5	0.5	long-finned squid	7,802	83.4	326	32.5
fourspot flounder	466	0.5	88.1	0.8	horseshoe crab	109	1.2	205.8	20.5
striped searobin	366	0.4	113.5	1.1	American lobster	748	8	197.9	19.7
moonfish	361	0.4	3.5	0	boring sponge	nc	nc	51.3	5.1
smooth dogfish	332	0.4	1,176.6	11.2	spider crab	nc	nc	50.6	5
spotted hake	321	0.3	24.3	0.2	lion's mane jellyfish	558	6	45.4	4.5
weakfish	241	0.3	52.2	0.5	rock crab	nc	nc	40.4	4
summer flounder	203	0.2	180.5	1.7	bushy bryozoan	nc	nc	17.8	1.8
tautog	186	0.2	301.4	2.9	blue mussel	nc	nc	7.6	0.8
striped bass	144	0.2	418.7	4	channeled whelk	41	0.4	7.6	0.8
hickory shad	75	0.1	19.1	0.2	lady crab	nc	nc	7.5	0.7
American shad	68	0.1	6.1	0.1	deadman's fingers sponge	nc	nc	6.8	0.7
Atlantic herring	66	0.1	10.3	0.1	hydroid spp.	nc	nc	5.9	0.6
blueback herring	63	0.1	2.5	0	flat claw hermit crab	nc	nc	5.7	0.6
clearnose skate	36	0	52.4	0.5	starfish spp.	nc	nc	4.8	0.5
Atlantic menhaden	28	0	5.5	0.1	rubbery bryozoan	nc	nc	4	0.4
winter skate	23	0	60	0.6	common slipper shell	nc	nc	3.9	0.4
hogchoker	22	0	3.2	0	mantis shrimp	70	0.7	3.4	0.3
Atlantic sturgeon	21	0	368.7	3.5	mud crabs	nc	nc	2.1	0.2
black sea bass	19	0	9.3	0.1	blue crab	11	0.1	1.8	0.2
fourbeard rockling	14	0	1.5	0	knobbed whelk	5	0.1	1.2	0.1
rough scad	14	0	0.5	0	sand shrimp	nc	nc	0.6	0.1
spot	14	0	1.2	0	mixed sponge species	nc	nc	0.6	0.1
spiny dogfish	11	0	47	0.4	moon jelly	2	0	0.5	0
cunner	8	0	1.3	0	sea grape	nc	nc	0.5	0
smallmouth flounder	7	0	0.6	0	arks	nc	nc	0.4	0
ocean pout	5	0	0.9	0	purple sea urchin	2	0	0.4	0
glasseye snapper	4	0	0.1	0	star coral	nc	nc	0.3	0
inshore lizardfish	4	0	0.4	0	hard clams	1	0	0.3	0
northern pipefish	3	0	0.2	0	northern red shrimp	1	0	0.3	0
rock gunnel	2	0	0.1	0	red bearded sponge	nc	nc	0.2	0
yellow jack	2	0	0.1	0	fan worm tubes	nc	nc	0.2	0
Atlantic bonito	1	0	3.2	0	northern moon snail	nc	nc	0.2	0
planehead filefish	1	0	0.1	0	surf clam	1	0	0.2	0
goosefish	1	0	1.2	0	brown shrimp	1	0	0.1	0
pollock	1	0	0.1	0	ghost shrimp	nc	nc	0.1	0
oyster toadfish	1	0	1.2	0	Japanese shore crab	nc	nc	0.1	0
yellowtail flounder	1	0	0.4	0	northern cyclocardia	nc	nc	0.1	0
Total	92,042		10,500.2		Total	9,352		1,002.6	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in 2007.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring and American sand lance are not quantified. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
scup	75,681	42.6	5,333.5	30.4	grubby	1	0	0.1	0
butterfish	49,137	27.6	1,446.2	8.2	pollock	1	0	0.1	0
weakfish	17,386	9.8	584.8	3.3	rock gunnel	1	0	0.1	0
bluefish	9,378	5.3	1,801.3	10.3	striped burrfish	1	0	0.5	0
winter flounder	4,550	2.6	951.3	5.4	sea lamprey	1	0	0.1	0
windowpane flounder	4,051	2.3	510.8	2.9	yellowtail flounder	1	0	1.0	0
red hake	2,788	1.6	200.4	1.1					
bay anchovy	2,440	1.4	14.5	0.1	Finfish not ranked				
Atlantic herring	1,932	1.1	234.2	1.3	anchovy spp. yoy				
alewife	1,537	0.9	101.3	0.6	Atlantic herring, yoy				
little skate	1,277	0.7	697.0	4.0	American sand lance (yoy)				
fourspot flounder	1,094	0.6	224.9	1.3					
moonfish	979	0.6	12.0	0.1	Invertebrates				
striped searobin	755	0.4	217.0	1.2	long-finned squid	24,212	88.2	773.6	30.8
summer flounder	733	0.4	590.9	3.4	horseshoe crab	333	1.2	596.4	23.7
northern searobin	691	0.4	74.2	0.4	American lobster	1,648	6.0	396.5	15.8
smooth dogfish	580	0.3	2,110.2	12.0	spider crab	nc	nc	165.5	6.6
Atlantic menhaden	426	0.2	63.9	0.4	lion's mane jellyfish	660	2.4	129.8	5.2
striped bass	422	0.2	888.0	5.1	bushy bryozoan	nc	nc	107.4	4.3
spotted hake	340	0.2	23.9	0.1	mixed sponge species	nc	nc	84.5	3.4
silver hake	290	0.2	14.6	0.1	rock crab	nc	nc	41.4	1.6
tautog	280	0.2	551.4	3.1	channeled whelk	196	0.7	33.4	1.3
American shad	236	0.1	15.8	0.1	flat claw hermit crab	nc	nc	27.5	1.1
blueback herring	156	0.1	9.1	0.1	blue mussel	nc	nc	20.4	0.8
black sea bass	116	0.1	46.8	0.3	starfish spp.	nc	nc	20.3	0.8
clearnose skate	97	0.1	193.3	1.1	boring sponge	nc	nc	17.7	0.7
fourbeard rockling	87	0	7.6	0	blue crab	68	0.2	13.0	0.5
hogchoker	78	0	11.4	0.1	mantis shrimp	264	1.0	12.1	0.5
smallmouth flounder	48	0	2.6	0	deadman's fingers sponge	nc	nc	11.5	0.5
winter skate	44	0	117.8	0.7	lady crab	nc	nc	11.5	0.5
hickory shad	37	0	10.4	0.1	knobbed whelk	23	0.1	11.1	0.4
spiny dogfish	32	0	122.3	0.7	common slipper shell	nc	nc	9.3	0.4
American sand lance	30	0	0.3	0	mud crabs	nc	nc	4.3	0.2
Atlantic sturgeon	18	0	336.4	1.9	northern moon snail	nc	nc	4.3	0.2
cunner	16	0	3.0	0	sand shrimp	nc	nc	3.5	0.1
rough scad	13	0	0.7	0	sea grape	nc	nc	3.5	0.1
ocean pout	12	0	3.2	0	arks	2	0	2.7	0.1
Atlantic mackerel	9	0	0.8	0	hydroid spp.	nc	nc	2.5	0.1
glasseye snapper	8	0	0.7	0	hard clams	1	0	2.2	0.1
northern puffer	8	0	0.5	0	rubbery bryozoan	nc	nc	1.4	0.1
striped anchovy	6	0	0.1	0	common oyster	nc	nc	1.1	0
sea raven	5	0	3.6	0	surf clam	10	0	1.0	0
oyster toadfish	5	0	2.0	0	anemones	16	0.1	0.6	0
yellow jack	5	0	0.4	0	purple sea urchin	2	0	0.6	0
northern kingfish	4	0	0.4	0	red bearded sponge	nc	nc	0.5	0
round scad	3	0	0.3	0	star coral	nc	nc	0.4	0
longhorn sculpin	3	0	0.8	0	water jelly	1	0	0.3	0
American eel	2	0	0.9	0	jonah crab	1	0	0.2	0
inshore lizardfish	2	0	0.2	0	northern red shrimp	1	0	0.2	0
mackerel scad	2	0	0.1	0	blood star	nc	nc	0.1	0
northern sennet	2	0	0.2	0	coastal mud shrimp	1	0	0.1	0
northern pipefish	2	0	0.2	0	green sea urchin	1	0	0.1	0
Atlantic silverside	1	0	0.1	0	Japanese shore crab	nc	nc	0.1	0
gizzard shad	1	0	0.1	0	tunicates, misc	1	0	0.1	0
Total	177,841		17,540.3		Total	27,441		2,512.7	

Note: nc= not counted

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in 2008.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring and American sand lance are not quantified. Number of tows (sample size)=120.

species	count	%	weight	%	species	count	%	weight	%
scup	53,560	38	6,509.9	45.7	sea lamprey	1	0	0.8	0
butterfish	48,766	34.6	1,442.0	10.1	striped anchovy	1	0	0.1	0
American sand lance	7,495	5.3	7.2	0.1	Total	140,777		14,239.8	
silver hake	6,587	4.7	208.5	1.5					
winter flounder	4,973	3.5	751.9	5.3	Finfish not ranked				
windowpane flounder	3,511	2.5	524.0	3.7	anchovy spp. yoy				
weakfish	2,531	1.8	116.1	0.8	Atlantic herring, yoy				
red hake	1,723	1.2	141.3	1.0	American sand lance (yoy)				
bluefish	1,699	1.2	641.4	4.5					
spotted hake	1,267	0.9	65.8	0.5	Invertebrates				
bay anchovy	1,128	0.8	7.7	0.1	horseshoe crab	289	2.2	496.8	29.2
alewife	931	0.7	51.1	0.4	long-finned squid	10,490	80.5	330.1	19.4
fourspot flounder	902	0.6	186.3	1.3	American lobster	1,096	8.4	314.1	18.5
northern searobin	809	0.6	58.8	0.4	spider crab	nc	nc	145.8	8.6
moonfish	689	0.5	13.4	0.1	rock crab	nc	nc	64.0	3.8
little skate	682	0.5	327.4	2.3	bushy bryozoan	nc	nc	54.2	3.2
striped searobin	612	0.4	263.0	1.8	lady crab	nc	nc	36.3	2.1
summer flounder	477	0.3	398.0	2.8	starfish spp.	nc	nc	32.1	1.9
American shad	405	0.3	20.2	0.1	boring sponge	nc	nc	30.1	1.8
Atlantic herring	356	0.3	52.1	0.4	channeled whelk	177	1.4	29.3	1.7
smooth dogfish	328	0.2	1,134.2	8.0	mixed sponge species	nc	nc	27.8	1.6
spot	308	0.2	21.3	0.1	hydroid spp.	nc	nc	24.6	1.4
striped bass	199	0.1	456.3	3.2	flat claw hermit crab	nc	nc	22.8	1.3
tautog	179	0.1	309.4	2.2	common slipper shell	nc	nc	15.7	0.9
black sea bass	122	0.1	29.8	0.2	lion's mane jellyfish	520	4	14.3	0.8
smallmouth flounder	89	0.1	3.2	0	mantis shrimp	244	1.9	9.1	0.5
fourbeard rockling	81	0.1	7.1	0	sea grape	nc	nc	6.6	0.4
blueback herring	74	0.1	3.2	0	arks	124	1	6.1	0.4
winter skate	51	0	140.8	1.0	knobbed whelk	17	0.1	5.9	0.3
Atlantic menhaden	47	0	10.4	0.1	blue mussel	nc	nc	5.8	0.3
hogchoker	38	0	5.6	0	northern moon snail	1	0	5.6	0.3
clearnose skate	37	0	78.1	0.5	sand shrimp	nc	nc	4.0	0.2
spiny dogfish	35	0	127.7	0.9	blue crab	16	0.1	3.8	0.2
cunner	26	0	3.6	0	mud crabs	nc	nc	3.5	0.2
inshore lizardfish	10	0	0.5	0	rubbery bryozoan	nc	nc	3.1	0.2
ocean pout	9	0	2.1	0	common oyster	1	0	2.1	0.1
Atlantic sturgeon	7	0	111.3	0.8	hard clams	8	0.1	1.4	0.1
hickory shad	5	0	1.1	0	purple sea urchin	15	0.1	0.9	0.1
feather blenny	4	0	0.2	0	northern red shrimp	21	0.2	0.7	0
white perch	4	0	0.1	0	deadman's fingers sponge	nc	nc	0.6	0
northern kingfish	3	0	0.4	0	surf clam	9	0.1	0.6	0
oyster toadfish	3	0	1.9	0	red bearded sponge	nc	nc	0.4	0
Atlantic silverside	2	0	0.2	0	Jonah crab	2	0	0.4	0
rock gunnel	2	0	0.2	0	star coral	nc	nc	0.3	0
longhorn sculpin	2	0	0.3	0	sea cucumber	2	0	0.3	0
yellowtail flounder	2	0	0.4	0	tunicates, misc	nc	nc	0.3	0
Atlantic croaker	1	0	0.1	0	anemones	nc	nc	0.2	0
planehead filefish	1	0	0.1	0	coastal mud shrimp	1	0	0.1	0
glasseye snapper	1	0	0.1	0	green crab	1	0	0.1	0
pollock	1	0	0.1	0	moon jelly	1	0	0.1	0
rougtail stingray	1	0	3.0	0	northern cyclocardia	1	0	0.1	0
					Total	13,036		1,700.1	

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in 2009.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring and American sand lance are not quantified. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	108,087	53.6	3,186.9	17	striped cusk-eel	1	0	0.1	0
scup	46,991	23.3	6,332.1	33.8	spot	1	0	0.2	0
bay anchovy	11,128	5.5	35.3	0.2	northern stargazer	1	0	0.1	0
Atlantic herring	6,330	3.1	239.2	1.3	Atlantic tomcod	1	0	0.1	0
winter flounder	4,068	2	524.0	2.8	white perch	1	0	0.1	0
bluefish	3,657	1.8	1,157.4	6.2	yellow jack	1	0	0.1	0
weakfish	2,604	1.3	108.7	0.6	yellowtail flounder	1	0	0.2	0
moonfish	2,575	1.3	19.5	0.1	Total	201,476		18,750	
windowpane flounder	2,496	1.2	342.8	1.8					
northern searobin	2,012	1	194.3	1	Finfish not ranked				
striped searobin	1,507	0.7	471.8	2.5	anchovy spp, yoy				
American sand lance	1,227	0.6	2.0	0	Atlantic herring, yoy				
alewife	1,175	0.6	96.0	0.5	American sand lance (yoy)				
fourspot flounder	1,036	0.5	169.8	0.9					
silver hake	947	0.5	50.0	0.3	Invertebrates				
red hake	897	0.4	59.5	0.3	long-finned squid	24,130	91.4	648.4	30.2
summer flounder	881	0.4	694.4	3.7	horseshoe crab	340	1.3	645.8	30
little skate	709	0.4	390.0	2.1	American lobster	853	3.2	244	11.3
smooth dogfish	588	0.3	2,213.3	11.8	spider crab	.	.	144.1	6.7
striped bass	466	0.2	897.4	4.8	lion's mane jellyfish	641	2.4	89.3	4.2
American shad	422	0.2	28.9	0.2	lady crab	.	.	63.6	3
spotted hake	327	0.2	32.1	0.2	rock crab	.	.	42.4	2
blueback herring	291	0.1	14.6	0.1	common slipper shell	.	.	37	1.7
tautog	163	0.1	285.4	1.5	flat claw hermit crab	.	.	33.8	1.6
spiny dogfish	148	0.1	545.7	2.9	bushy bryozoan	.	.	33.3	1.5
black sea bass	121	0.1	59.5	0.3	starfish spp.	.	.	26.6	1.2
smallmouth flounder	96	0	4.7	0	channeled whelk	127	0.5	26	1.2
clearnose skate	69	0	148.5	0.8	hydroid spp.	.	.	25.7	1.2
Atlantic menhaden	69	0	18.0	0.1	knobbed whelk	39	0.1	11.6	0.5
rough scad	59	0	2.8	0	mantis shrimp	215	0.8	10.7	0.5
fourbeard rockling	47	0	3.9	0	Tubularia, spp.	.	.	9	0.4
winter skate	44	0	108.5	0.6	northern moon snail	.	.	7.2	0.3
hogchoker	39	0	4.5	0	anemones	.	.	5.6	0.3
blue runner	34	0	2.3	0	mixed sponge species	.	.	5.4	0.3
ocean pout	22	0	4.8	0	sea grape	.	.	5.0	0.2
Atlantic sturgeon	18	0	286.6	1.5	boring sponge	.	.	4.2	0.2
cunner	18	0	1.8	0	blue crab	19	0.1	4.1	0.2
pollock	18	0	0.8	0	sand shrimp	.	.	3.8	0.2
Atlantic cod	15	0	1.0	0	deadman's fingers sponge	.	.	3.5	0.2
hickory shad	13	0	3.6	0	blue mussel	8	0	3.5	0.2
northern kingfish	7	0	0.4	0	mud crabs	.	.	3.1	0.1
glasseye snapper	6	0	0.6	0	common oyster	1	0	3.1	0.1
Atlantic mackerel	5	0	0.4	0	arks	2	0	2.5	0.1
northern sennet	5	0	0.4	0	surf clam	18	0.1	1.7	0.1
northern puffer	5	0	0.4	0	hard clams	4	0	1.1	0.1
sea raven	5	0	1.7	0	red bearded sponge	.	.	0.8	0
striped anchovy	5	0	0.4	0	purple sea urchin	4	0	0.8	0
Atlantic silverside	3	0	0.3	0	rubbery bryozoan	.	.	0.6	0
oyster toadfish	3	0	0.8	0	star coral	.	.	0.2	0
inshore lizardfish	2	0	0.2	0	ghost shrimp	2	0	0.2	0
northern pipefish	2	0	0.2	0	coastal mud shrimp	2	0	0.1	0
rock gunnel	2	0	0.2	0	northern cyclocardia	1	0	0.1	0
longhorn sculpin	2	0	0.3	0	northern red shrimp	1	0	0.1	0
crevalle jack	1	0	0.1	0	sea cucumber	1	0	0.1	0
planehead filefish	1	0	0.1	0	tunicates, misc	1	0	0.1	0
round scad	1	0	0.1	0	Total	26,409		2,148.2	

Note: nc= not counted

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in 2010.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring and American sand lance are not quantified. Number of tows (sample size)=78.

species	count	%	weight	%	species	count	%	weight	%
American sand lance	13,061	35.3	5.2	0.1	<u>Invertebrates</u>				
scup	7,157	19.3	1,971.6	44.3	long-finned squid	1,906	62.9	161.4	28.4
butterfish	2,894	7.8	166.9	3.7	horseshoe crab	58	1.9	112.2	19.8
windowpane flounder	2,850	7.7	449.3	10.1	American lobster	293	9.7	83.6	14.7
winter flounder	2,579	7.0	450.5	10.1	spider crab	.	.	81.6	14.4
silver hake	1,747	4.7	35.4	0.8	bushy bryozoan	.	.	23.1	4.1
Atlantic herring	1,318	3.6	179.0	4	rock crab	.	.	16.7	2.9
northern searobin	1,128	3	149.5	3.4	starfish spp.	.	.	15.1	2.7
red hake	990	2.7	64.3	1.4	common slipper shell	.	.	11.2	2
spotted hake	665	1.8	15.8	0.4	lion's mane jellyfish	401	13.2	7.8	1.4
summer flounder	517	1.4	229.6	5.2	lady crab	.	.	7.7	1.4
bay anchovy	475	1.3	2.8	0.1	flat claw hermit crab	.	.	6.8	1.2
fourspot flounder	402	1.1	92.0	2.1	hydroid spp.	.	.	6.7	1.2
little skate	281	0.8	148.3	3.3	channeled whelk	33	1.1	4.5	0.8
alewife	172	0.5	14.3	0.3	northern moon snail	.	.	4.1	0.7
American shad	165	0.4	8.6	0.2	blue mussel	.	.	3.1	0.5
striped searobin	141	0.4	66.4	1.5	common oyster	.	.	2.9	0.5
blueback herring	101	0.3	3.4	0.1	sea grape	.	.	2.7	0.5
striped bass	71	0.2	173.2	3.9	sand shrimp	.	.	2.3	0.4
tautog	53	0.1	83.1	1.9	deadman's fingers sponge	.	.	2.3	0.4
black sea bass	37	0.1	20.1	0.5	blue crab	10	0.3	2.0	0.4
fourbeard rockling	35	0.1	2.9	0.1	arks	.	.	1.6	0.3
hogchoker	34	0.1	4.4	0.1	mud crabs	.	.	1.6	0.3
smallmouth flounder	31	0.1	1.4	0	rubbery bryozoan	.	.	1.2	0.2
rock gunnel	29	0.1	0.5	0	mantis shrimp	19	0.6	1.1	0.2
Atlantic cod	21	0.1	2.1	0	Unknown Jellyfish	300	9.9	0.8	0.1
winter skate	16	0	37.7	0.8	Tubularia, spp.	.	.	0.5	0.1
cunner	11	0	1.3	0	anemones	5	0.1	0.4	0.1
smooth dogfish	10	0	34.4	0.8	surf clam	2	0.1	0.4	0.1
Atlantic menhaden	7	0	2.7	0.1	knobbed whelk	1	0	0.3	0.1
ocean pout	6	0	1.4	0	mixed sponge species	.	.	0.3	0.1
sea raven	6	0	1.6	0	northern comb jelly	1	0	0.2	0
northern pipefish	4	0	0.3	0	purple sea urchin	4	0.1	0.2	0
spiny dogfish	3	0	16.2	0.4	boring sponge	.	.	0.1	0
bluefish	2	0	6.1	0.1	red bearded sponge	.	.	0.1	0
hickory shad	2	0	0.4	0	coastal mud shrimp	.	.	0.1	0
pollock	2	0	0.1	0	star coral	.	.	0.1	0
American plaice	1	0	0.1	0	hard clams	.	.	0.1	0
Atlantic silverside	1	0	0.1	0	sea cucumber	.	.	0.1	0
Atlantic sturgeon	1	0	5.6	0.1	Total	3,033		567.0	
clearnose skate	1	0	4.5	0.1	Note: nc= not counted				
longhorn sculpin	1	0	0.4	0					
weakfish	1	0	1.0	0					
Total	37,029		4,455						

Finfish not ranked

anchovy spp, yoy
 Atlantic herring, yoy
 American sand lance (yoy)

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in 2011.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring and American sand lance are not quantified. Number of tows (sample size)=172.

species	count	%	weight	%	species	count	%	weight	%
butterfish	42,141	36.7	1,600.8	9.9	striped burrfish	1	0	0.5	0
scup	34,458	30.0	6,759.0	41.7	striped anchovy	1	0	0.1	0
American sand lance	9,535	8.3	7.5	0.0	silver perch	1	0	0.1	0
bay anchovy	4,693	4.1	10.5	0.1	oyster toadfish	1	0	0.2	0
winter flounder	3,092	2.7	613.8	3.8	white perch	1	0	0.1	0
windowpane flounder	2,831	2.5	395.9	2.4	white mullet	1	0	0.1	0
bluefish	2,765	2.4	584.7	3.6	yellowtail flounder	1	0	0.3	0
weakfish	2,583	2.3	192.6	1.2	Total	114,706		16,210.3	
striped searobin	1,630	1.4	558.7	3.4	<u>Finfish not ranked</u>				
Atlantic herring	1,482	1.3	199.4	1.2	anchovy spp, yoy				
fourspot flounder	1,400	1.2	224.2	1.4	Atlantic herring, yoy				
summer flounder	1,051	0.9	713.0	4.4	American sand lance (yoy)				
silver hake	948	0.8	40.3	0.2	<u>Invertebrates</u>				
northern searobin	803	0.7	85.5	0.5	horseshoe crab				
spotted hake	725	0.6	76.8	0.5	long-finned squid				
little skate	674	0.6	359.4	2.2	spider crab				
moonfish	640	0.6	6.3	0	lady crab				
smooth dogfish	613	0.5	2,031.7	12.5	American lobster				
alewife	512	0.4	29.8	0.2	rock crab				
red hake	278	0.2	25.1	0.2	hydroid spp.				
American shad	271	0.2	17.5	0.1	mantis shrimp				
striped bass	243	0.2	721.9	4.5	bushy bryozoan				
Atlantic menhaden	181	0.2	69.8	0.4	knobbed whelk				
rough scad	150	0.1	6.8	0	flat claw hermit crab				
hogchoker	147	0.1	16.8	0.1	channeled whelk				
Atlantic cod	109	0.1	9.2	0.1	starfish spp.				
tautog	106	0.1	151.7	0.9	blue crab				
black sea bass	91	0.1	54.2	0.3	lion's mane jellyfish				
blueback herring	72	0.1	3.2	0	mixed sponge species				
smallmouth flounder	67	0.1	3.5	0	blue mussel				
spiny dogfish	58	0.1	203.5	1.3	northern moon snail				
clearnose skate	56	0	109.8	0.7	boring sponge				
inshore lizardfish	43	0	4.6	0	hard clams				
fourbeard rockling	43	0	4.0	0	common slipper shell				
winter skate	37	0	101.2	0.6	sand shrimp				
northern kingfish	34	0	3.7	0	Tubularia, spp.				
ocean pout	27	0	4.5	0	mud crabs				
blue runner	24	0	1.7	0	rubbery bryozoan				
cunner	14	0	1.9	0	common oyster				
northern puffer	9	0	0.9	0	sea grape				
longhorn sculpin	9	0	2.0	0	arks				
hickory shad	8	0	1.5	0	surf clam				
Atlantic sturgeon	5	0	181.9	1.1	purple sea urchin				
pollock	5	0	0.5	0	red bearded sponge				
spot	5	0	0.7	0	northern comb jelly				
crevalle jack	4	0	0.4	0	anemones				
grubby	4	0	0.1	0	star coral				
northern pipefish	4	0	0.3	0	coastal mud shrimp				
rock gunnel	4	0	0.2	0	common razor clam				
conger eel	3	0	1.1	0	ghost shrimp				
sea raven	3	0	0.9	0	northern red shrimp				
striped cusk-eel	2	0	0.2	0	polychaetes				
Atlantic tomcod	2	0	0.2	0	tunicates, misc				
American plaice	1	0	0.1	0	water jelly				
Atlantic croaker	1	0	0.2	0	Total				
northern sennet	1	0	0.1	0	15,076				
round scad	1	0	0.1	0	1,505.0				
rougtail stingray	1	0	13.0	0.1	Note: nc= not counted				

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in 2012.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring and American sand lance are not quantified. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	60,539	37.9	1,891.3	10.8	longhorn sculpin	1	0	0.2	0
scup	53,119	33.2	6,170.2	35.1	white perch	1	0	0.2	0
silver hake	7,519	4.7	171.0	1.0	white mullet	1	0	0.1	0
weakfish	6,785	4.2	409.2	2.3	Total	159,770		17,570.3	
bluefish	3,851	2.4	532.7	3.0					
northern searobin	3,642	2.3	405.2	2.3	Finfish not ranked				
windowpane flounder	3,536	2.2	501.1	2.9	anchovy spp. yoy				
winter flounder	3,365	2.1	604.9	3.4	Atlantic herring, yoy				
striped searobin	2,973	1.9	1,086.4	6.2	American sand lance (yoy)				
fourspot flounder	2,597	1.6	454.5	2.6					
red hake	1,720	1.1	148.6	0.8	Invertebrates				
little skate	1,406	0.9	657.9	3.7	horseshoe crab	199	1.7	385.8	30.6
bay anchovy	1,296	0.8	8.6	0.0	long-finned squid	9,767	84.5	333.9	26.5
summer flounder	980	0.6	718.5	4.1	spider crab	.	.	162.4	12.9
spot	858	0.5	107.5	0.6	American lobster	349	3.0	70.0	5.6
alewife	708	0.4	47.0	0.3	boring sponge	.	.	47.9	3.8
spotted hake	626	0.4	64.2	0	lady crab	.	.	45.3	3.6
smooth dogfish	610	0.4	1,833.3	10.4	rock crab	.	.	40.7	3.2
Atlantic herring	571	0.4	61.5	0.4	mantis shrimp	846	7.3	26.6	2.1
Atlantic menhaden	426	0.3	144.6	0.8	bushy bryozoan	.	.	20.4	1.6
black sea bass	410	0.3	141.0	0.8	flat claw hermit crab	.	.	18.3	1.5
hogchoker	340	0.2	30.7	0.2	blue crab	72	0.6	14.5	1.2
American shad	321	0.2	25.3	0.1	knobbed whelk	36	0.3	13.8	1.1
clearnose skate	280	0.2	491.7	3	channeled whelk	76	0.7	13.7	1.1
moonfish	262	0.2	3.6	0.0	blue mussel	1	0.0	9.4	0.7
smallmouth flounder	258	0.2	7.5	0.0	common slipper shell	.	.	9.4	0.7
striped bass	170	0.1	278.0	1.6	mixed sponge species	.	.	7.4	0.6
tautog	135	0.1	128.9	0.7	Tubularia, spp.	.	.	5.0	0.4
winter skate	97	0.1	179.8	1	hydroid spp.	.	.	4.8	0.4
northern kingfish	59	0.0	8.4	0	lion's mane jellyfish	50	0.4	4.4	0.3
northern puffer	47	0.0	3.1	0.0	mud crabs	.	.	3.9	0.3
blueback herring	46	0	1.6	0.0	starfish spp.	.	.	3.3	0.3
fourbeard rockling	43	0	3.5	0	northern red shrimp	118	1.0	3.0	0.2
hickory shad	42	0	14.1	0	northern moon snail	.	.	1.8	0.1
blue runner	27	0	2.7	0.0	sand shrimp	.	.	1.7	0.1
cunner	20	0	2.8	0	arks	.	.	1.4	0.1
rough scad	19	0	1.1	0	hard clams	3	0	1.3	0.1
spiny dogfish	16	0	62.8	0	red bearded sponge	.	.	1.2	0.1
ocean pout	14	0	2.0	0	sea grape	.	.	1.1	0.1
Atlantic sturgeon	7	0	154.2	1	deadman's fingers sponge	.	.	0.8	0.1
sea raven	5	0	1.1	0	purple sea urchin	7	0	0.8	0
northern sennet	3	0	0.3	0	common oyster	.	.	0.8	0
striped anchovy	3	0	0.2	0.0	surf clam	10	0.1	0.8	0
crevalle jack	2	0	0.2	0	star coral	.	.	0.4	0
goosefish	2	0	0.8	0	rubbery bryozoan	.	.	0.4	0
pinfish	2	0	0.2	0	sea cucumber	3	0	0.4	0
round herring	2	0	0.1	0	tunicates, misc	16	0	0.4	0
American sand lance	2	0	0.2	0	water jelly	4	0	0.3	0
African pompano	1	0	0.1	0	coastal mud shrimp	1	0	0.2	0
conger eel	1	0	0.3	0	northern comb jelly	.	.	0.1	0
gizzard shad	1	0	0.1	0	moon jelly	.	.	0.1	0
northern pipefish	1	0	0.1	0	Total	11,558		1,257.9	
rock gunnel	1	0	0.1	0					
rougtail stingray	1	0	5.0	0					

Note: nc= not counted

Appendix 5.4. cont. Total number and weight (kg) of finfish and invertebrates caught in 2013.

Finfish species are in order of descending count. Invertebrate species are in order of descending weight (nc = not counted). Young-of-year bay and striped anchovy are neither separated by species or quantified; young-of-year Atlantic herring and American sand lance are not quantified. Number of tows (sample size)=200.

species	count	%	weight	%	species	count	%	weight	%
butterfish	29,569	35.4	1,252.5	7.9					
scup	24,961	29.9	5,945.6	37.5					
Atlantic herring	3,566	4.3	321.2	2.0	Finfish not ranked				
striped searobin	2,724	3.3	1,112.5	7.0	anchovy spp. (yoy)				
windowpane flounder	2,096	2.5	326.6	2.1	Atlantic herring. (yoy)				
weakfish	1,964	2.4	203.7	1.3	American sand lance (yoy)				
northern searobin	1,934	2.3	161.7	1.0	gadid spp. (yoy)				
spot	1,917	2.3	195.4	1.2	Invertebrates				
winter flounder	1,912	2.3	576.8	3.6	blue mussel	3	0.0	622.1	31.9
bluefish	1,829	2.2	517.7	3.3	horseshoe crab	265	3.4	531.8	27.3
bay anchovy	1,350	1.6	6.8	0.0	long-finned squid	5,393	69.6	170.8	8.8
fourspot flounder	1,144	1.4	203.4	1.3	spider crab	nc		156.5	8.0
summer flounder	1,071	1.3	726.6	4.6	lion's mane jellyfish	1,067	13.8	150.0	7.7
smooth dogfish	1,051	1.3	2,162.3	13.6	common slipper shell	nc		61.0	3.1
spotted hake	927	1.1	66.8	0.4	American lobster	144	1.9	37.3	1.9
moonfish	868	1.0	10.0	0.1	bushy bryozoan	nc		26.8	1.4
red hake	849	1.0	61.1	0.4	boring sponge	nc		26.1	1.3
little skate	583	0.7	317.8	2.0	mantis shrimp	646	8.3	21.6	1.1
silver hake	519	0.6	23.6	0.1	flat claw hermit crab	nc		21.4	1.1
black sea bass	449	0.5	181.2	1.1	knobbed whelk	51	0.7	18.7	1.0
alewife	376	0.5	34.1	0.2	channeled whelk	95	1.2	18.6	1.0
hogchoker	250	0.3	27.2	0.2	hydroid spp.	nc		13.2	0.7
Atlantic menhaden	234	0.3	87.5	0.6	lady crab	nc		13.2	0.7
American shad	222	0.3	15.3	0.1	rock crab	nc		13.0	0.7
clearnose skate	218	0.3	387.0	2.4	blue crab	52	0.7	10.4	0.5
striped bass	200	0.2	421.0	2.7	Tubularia, spp.	nc		6.7	0.3
tautog	161	0.2	160.8	1.0	common oyster	nc		5.3	0.3
smallmouth flounder	128	0.2	5.2	0.0	mud crabs	nc		3.5	0.2
winter skate	91	0.1	111.2	0.7	sand shrimp	nc		2.9	0.1
blueback herring	68	0.1	4.3	0.0	northern moon snail	nc		2.9	0.1
hickory shad	33	0.0	10.8	0.1	surf clam	8	0.1	2.4	0.1
rough scad	28	0.0	1.3	0.0	starfish spp.	1	0.0	2.1	0.1
red goatfish	21	0.0	0.5	0.0	sea grape	nc		2.1	0.1
spiny dogfish	21	0.0	91.5	0.6	arks	nc		1.9	0.1
cunner	20	0.0	1.8	0.0	hard clams	6	0.1	0.9	0.0
northern kingfish	14	0.0	2.3	0.0	comb jelly spp	nc		0.8	0.0
American sand lance	7	0.0	0.1	0.0	red bearded sponge	nc		0.6	0.0
haddock	5	0.0	0.4	0.0	rubbery bryozoan	nc		0.5	0.0
oyster toadfish	5	0.0	0.9	0.0	purple sea urchin	10	0.1	0.5	0.0
Atlantic sturgeon	4	0.0	98.0	0.6	coastal mud shrimp	4	0.1	0.3	0.0
Atlantic silverside	3	0.0	0.3	0.0	deadman's fingers sponge	nc		0.3	0.0
northern puffer	3	0.0	0.3	0.0	mixed sponge species	nc		0.3	0.0
fourbeard rockling	3	0.0	0.2	0.0	star coral	nc		0.2	0.0
bullnose ray	2	0.0	5.7	0.0	sea cucumber	2	0.0	0.2	0.0
harvestfish	2	0.0	0.2	0.0	fan worm tubes	nc		0.1	0.0
northern pipefish	2	0.0	0.2	0.0	ghost shrimp	1	0.0	0.1	0.0
conger eel	1	0.0	1.2	0.0	Japanese shore crab	1	0.0	0.1	0.0
Atlantic croaker	1	0.0	0.1	0.0	northern red shrimp	1	0.0	0.1	0.0
glasseye snapper	1	0.0	0.1	0.0	ribbed mussel	nc		0.1	0.0
pollock	1	0.0	0.1	0.0	Total	7,750		1,947.4	
round scad	1	0.0	0.1	0.0	Note: nc= not counted				
red cornetfish	1	0.0	0.1	0.0					
longhorn sculpin	1	0.0	0.4	0.0					
striped anchovy	1	0.0	0.1	0.0					
northern stargazer	1	0.0	0.1	0.0					
Total	83,413		15,843.7						

Appendix 5.5: Endangered Species Interactions: Thirteen (13) Atlantic sturgeon (ATS) were captured on two of the 199 tows completed in 2014. This yields a lower encounter rate (1.0%) than the average for the LISTS time series of tows (2.4%). One tow occurred over transition bottom type in the 18.3-27.3m depth interval, while the other occurred over sand bottom type in the 5-9m depth interval. All individuals were released alive and uninjured. Each sturgeon was scanned for a passive integrated transponder (PIT). Since no PITs were detected, a PIT was inserted near the base of each dorsal fin. T-bar tags were inserted into sturgeon until the supply ran out (USFWS no longer supplies these tags). All captures were reported to NMFS within 24 hours. Details for each fish are provided below:

Sample	Date	Site	Tow Start	Duration (min)	Species	Total Length (mm)	Fork Length (mm)	Weight (kg)	Left Pec T-bar	Dorsal T-bar	PIT	Photo	Tissue Sample	Release time	Release lat (N)	Release lon (W)
FA2014015	9/15/2014	1228	8:11	30	ATS	1,205	1,065	8.9	NONE	NONE	ADDED	YES	YES	10:03	41.2025	72.5630
FA2014015	9/15/2014	1228	8:11	30	ATS	1,265	1,095	9.6	NONE	NONE	ADDED	YES	YES	10:10	41.2023	72.5622
FA2014015	9/15/2014	1228	8:11	30	ATS	1,357	1,188	13.3	NONE	NONE	ADDED	YES	YES	10:15	41.2017	72.2272
FA2014015	9/15/2014	1228	8:11	30	ATS	1,458	1,280	16.7	ADDED	NONE	ADDED	YES	YES	9:59	41.2025	72.5643
FA2014015	9/15/2014	1228	8:11	30	ATS	1,470	1,282	13.8	ADDED	NONE	ADDED	YES	YES	9:52	41.2027	72.5655
FA2014015	9/15/2014	1228	8:11	30	ATS	1,610	1,425	23.8	ADDED	NONE	ADDED	YES	YES	9:42	41.2027	72.5678
FA2014015	9/15/2014	1228	8:11	30	ATS	1,630	1,443	22.9	ADDED	NONE	ADDED	YES	YES	9:48	41.2027	72.5665
FA2014015	9/15/2014	1228	8:11	30	ATS	1,705	1,508	30.5	ADDED	NONE	ADDED	YES	YES	9:30	41.2027	72.5713
FA2014015	9/15/2014	1228	8:11	30	ATS	1,880	1,625	40.7	ADDED	NONE	ADDED	YES	YES	9:23	41.2028	72.5732
FA2014015	9/15/2014	1228	8:11	30	ATS	1,905	1,655	35.4	ADDED	NONE	ADDED	YES	YES	9:36	42.2027	72.5698
FA2014015	9/15/2014	1228	8:11	30	ATS	2,010	1,780	42.3 (est)	ADDED	NONE	ADDED	YES	YES	9:12	41.2030	72.5765
FA2014067	10/29/2014	1533	9:46	30	ATS	1,020	880	5.4	NONE	NONE	ADDED	YES	YES	10:40	41.2456	72.3479
FA2014067	10/29/2014	1533	9:46	30	ATS	1,190	1,035	9.1	NONE	NONE	ADDED	YES	YES	10:35	41.2467	72.3453

Appendix 5.6: Cold and warm temperate species captured in LISTS. Thirty-three (33) species are included in the cold temperate group, while thirty-four (34) species are included in the warm temperate group. Cold temperate species are defined as being more abundant north of Cape Cod, MA than south of New York, behaviorally adapted to cold temperatures including subfreezing but prefers ~3-15⁰C, and spawns at lower end of temperature tolerance. Warm temperate species are defined as being more abundant south of New York than north of Cape Cod, MA, behaviorally avoids temperatures < 7-10⁰C; prefers ~11-22⁰C, and spawns at higher end of temperature tolerance.

Cold Temperate Group		Warm Temperate Group	
Common Name	Scientific Name	Common Name	Scientific Name
alewife	<i>Alosa pseudoharengus</i>	American eel	<i>Anguilla rostrata</i>
American plaice	<i>Hippoglossoides platessoides</i>	American shad	<i>Alosa sapidissima</i>
Atlantic herring	<i>Clupea harengus</i>	Atlantic bonito	<i>Sarda sarda</i>
Atlantic cod	<i>Gadus morhua</i>	Atlantic croaker	<i>Micropogonias undulatus</i>
Atlantic mackerel	<i>Scomber scombrus</i>	Atlantic silversides	<i>Menidia menidia</i>
Atlantic salmon	<i>Salmo salar</i>	black seabass	<i>Centropristis striata</i>
Atlantic seasnail	<i>Liparis atlanticus</i>	blueback herring	<i>Alosa aestivalis</i>
Atlantic sturgeon	<i>Acipenser oxyrinchus</i>	bluefish	<i>Pomatomus saltatrix</i>
Atlantic tomcod	<i>Microgadus tomcod</i>	butterfish	<i>Peprilus triacanthus</i>
barndoor skate	<i>Dipturus laevis</i>	clearnose skate	<i>Raja eglanteria</i>
cunner	<i>Tautoglabrus adspersus</i>	conger eel	<i>Conger oceanicus</i>
fawn cusk-eel	<i>Lepophidium profundorum</i>	gizzard shad	<i>Dorosoma cepedianum</i>
fourspot flounder	<i>Hippoglossina oblonga</i>	hickory shad	<i>Alosa mediocris</i>
grubby	<i>Myoxocephalus aeneus</i>	hogchoker	<i>Trinectes maculatus</i>
haddock	<i>Melanogrammus aeglefinus</i>	lined seahorse	<i>Hippocampus erectus</i>
little skate	<i>Leucoraja erinacea</i>	menhaden	<i>Brevoortia tyrannus</i>
longhorn sculpin	<i>Myoxocephalus octodecemspinosus</i>	naked goby	<i>Gobiosoma boscii</i>
lumpfish	<i>Cyclopterus lumpus</i>	northern kingfish	<i>Menticirrhus saxatilis</i>
monkfish (goosefish)	<i>Lophius americanus</i>	northern puffer	<i>Sphoeroides maculatus</i>
northern pipefish	<i>Syngnathus fuscus</i>	northern searobin	<i>Prionotus carolinus</i>
ocean pout	<i>Zoarces americanus</i>	oyster toadfish	<i>Opsanus tau</i>
pollock	<i>Pollachius virens</i>	scup (porgy)	<i>Stenotomus chrysops</i>
rainbow smelt	<i>Osmerus mordax</i>	sea lamprey	<i>Petromyzon marinus</i>
red hake	<i>Urophycis chuss</i>	smallmouth flounder	<i>Etropus microstomus</i>
rock gunnel	<i>Pholis gunnellus</i>	smooth dogfish	<i>Mustelus canis</i>
rockling	<i>Enchelyopus cimbrius</i>	spot	<i>Leiostomus xanthurus</i>
searaven	<i>Hemitripterus americanus</i>	spotted hake	<i>Urophycis regia</i>
spiny dogfish	<i>Squalus acanthias</i>	striped bass	<i>Morone saxatilis</i>
whiting (silver hake)	<i>Merluccius bilinearis</i>	striped cusk-eel	<i>Ophidion marginatum</i>
windowpane	<i>Scophthalmus aquosus</i>	striped searobin	<i>Prionotus evolans</i>
winter flounder	<i>Pseudopleuronectes americanus</i>	summer flounder	<i>Paralichthys dentatus</i>
winter skate	<i>Leucoraja ocellata</i>	tautog (blackfish)	<i>Tautoga onitis</i>
yellowtail flounder	<i>Limanda ferruginea</i>	white perch	<i>Morone Americana</i>
		weakfish	<i>Cynoscion regalis</i>