

Number and percentage of persons receiving HIV care^e between 01/01/2016 through 12/31/2016 among persons who were >=13 years old on 12/31/2015, resided in Connecticut (based on most recent residence), diagnosed with HIV infection through 12/31/2015 and living with HIV on 12/31/2016, by selected characteristics - based on HIV surveillance data reported through 12/21/2017.

Characteristics	No. of persons diagnosed with HIV infection through 12/31/2015 and living with HIV on 12/31/2016 (overall population) ^h	No. of persons who have >=1 care visit ^f between 01/01/2016 through 12/31/2016	No. of persons who have >=2 care visits between 01/01/2016 through 12/31/2016, at least 3 months apart ^g	% of persons who have >=2 care visits between 01/01/2016 through 12/31/2016, at least 3 months apart among the overall population	% of persons who have >=2 care visits between 01/01/2016 through 12/31/2016, at least 3 months apart among persons have >=1 care visit between 01/01/2016 through 12/31/2016
Sex					
Male	6722	5141	3937	58.57	76.58
Female	3895	3085	2340	60.08	75.85
Age on 12/31/2015					
13-24	280	215	156	55.71	72.56
25-34	1046	809	541	51.72	66.87
35-44	1851	1414	1034	55.86	73.13
45-54	3615	2812	2165	59.89	76.99
55-64	2937	2299	1836	62.51	79.86
>=65	888	677	545	61.37	80.5
Race/ethnicity					
Black/African American	3589	2847	2167	60.38	76.12
Hispanic/Latino ^a	3642	2682	2075	56.97	77.37
White	3148	2507	1885	59.88	75.19
Other races ^b	238	190	150	63.03	78.95
Transmission category					
Male-to-male sexual contact (MSM)	2839	2288	1693	59.63	73.99
Injection drug use (IDU)	3119	2357	1903	61.01	80.74
MSM and IDU	235	186	149	63.4	80.11
Heterosexual contact ^c	3052	2475	1853	60.71	74.87
NIR/NRR (unknown)	1176	778	571	48.55	73.39
Other ^d	196	142	108	55.1	76.06
MSM					
Black/African American	594	478	331	55.72	69.25
Hispanic/Latino ^a	719	557	422	58.69	75.76
White	1451	1187	894	61.61	75.32
Other races ^b	75	66	46	61.33	69.70
Subtotal	2839	2288	1693	59.63	73.99
Injection drug use (male)					
Black/African American	684	538	440	64.33	81.78
Hispanic/Latino ^a	861	572	457	53.08	79.9
White	375	296	231	61.6	78.04
Other races ^b	25	20	19	76.00	95.00
Subtotal	1945	1426	1147	58.97	80.43
Injection drug use (female)					
Black/African American	395	325	262	66.33	80.62
Hispanic/Latino ^a	441	338	272	61.68	80.47
White	313	246	201	64.22	81.71
Other races ^b	25	22	21	84.00	95.45
Subtotal	1174	931	756	64.4	81.2
Heterosexual contact (male)					
Black/African American	429	337	264	61.54	78.34
Hispanic/Latino ^a	304	229	173	56.91	75.55
White	186	150	113	60.75	75.33
Other races ^b	23	20	17	73.91	85.00
Subtotal	942	736	567	60.19	77.04
Heterosexual contact (female)					
Black/African American	894	736	536	59.96	72.83
Hispanic/Latino ^a	817	672	514	62.91	76.49
White	357	298	213	59.66	71.48
Other races ^b	42	33	23	54.76	69.70
Subtotal	2110	1739	1286	60.95	73.95
Total	10617	8226	6277	59.12	76.31

^aHispanics/Latinos can be of any race.

^bIncludes American Indian/Alaska Native, Asian, Native Hawaiian/Other Pacific Islander, and Multiple races

^cHeterosexual contact with person known to have, or to be at high risk for, HIV infection.

^dIncludes hemophilia, blood transfusion, and persons (169 cases) who were exposed perinatally but were aged >13 at most recent year.

^ePersons who have at least one CD4 or viral load or HIV-1 genotype test are considered as receiving HIV care.

^fTests that have been done in the same month are considered as one care visit, even two tests that one is CD4 test and the other one is viral load (or genotype) test are still considered as one care visit.

^gTwo care visits should be 3 months apart.

^hThe overall population is overestimated because cases are only followed up for 11 months after 12/31/2016. CDC suggests that every case should be followed up at least 18 months to collect death certificate information.

