

Number and percentage of persons with HIV viral suppression* among persons who were >=13 years old on 12/31/2017, resided in Connecticut (based on most recent residence), diagnosed with HIV infection through 12/31/2017 and living with HIV on 12/31/2018, by selected characteristics
 - based on HIV surveillance data reported through 12/24/2019.

Characteristics	No. of persons diagnosed with HIV infection through 12/31/2017 and living with HIV on 12/31/2018 (overall population) ^f	No. of persons with >=1 VL test between 01/01/2018 through 12/31/2018	No. of persons with HIV viral suppression	% of persons with HIV viral suppression among the overall population	% of persons with HIV viral suppression among persons with >=1 VL test between 01/01/2018 through 12/31/2018
Sex					
Male	6841	5428	5013	73.28	92.35
Female	3503	2839	2557	72.99	90.07
Age on 12/31/2017					
13-24	243	187	147	60.49	78.61
25-34	1031	790	680	65.96	86.08
35-44	1558	1235	1100	70.6	89.07
45-54	3206	2566	2353	73.39	91.7
55-64	3150	2567	2411	76.54	93.92
>=65	1156	922	879	76.04	95.34
Race/ethnicity					
Black/African American	3507	2826	2519	71.83	89.14
Hispanic/Latino ^b	3514	2713	2474	70.4	91.19
White	3078	2529	2396	77.84	94.74
Other races	245	199	181	73.88	90.95
Transmission category					
Male-to-male sexual contact (MSM)	3137	2617	2456	78.29	93.85
Injection drug use (IDU)	2795	2169	1964	70.27	90.55
MSM and IDU	235	197	177	75.32	89.85
Heterosexual contact ^c	2876	2378	2153	74.86	90.54
NIR/NRR (unknown)	1121	770	719	64.14	93.38
Other ^d	180	136	101	56.11	74.26
MSM					
Black/African American	668	558	491	73.5	87.99
Hispanic/Latino ^b	847	689	649	76.62	94.19
White	1533	1298	1250	81.54	96.3
Other races	89	72	66	74.16	91.67
Subtotal	3137	2617	2456	78.29	93.85
Injection drug use (male)					
Black/African American	625	493	454	72.64	92.09
Hispanic/Latino ^b	802	570	518	64.59	90.88
White	334	269	249	74.55	92.57
Other races	23	17	16	69.57	94.12
Subtotal	1784	1349	1237	69.34	91.7
Injection drug use (female)					
Black/African American	343	288	258	75.22	89.58
Hispanic/Latino ^b	376	294	257	68.35	87.41
White	272	218	195	71.69	89.45
Other races	20	20	17	85	85
Subtotal	1011	820	727	71.91	88.66
Heterosexual contact (male)					
Black/African American	439	348	301	68.56	86.49
Hispanic/Latino ^b	308	240	221	71.75	92.08
White	184	152	143	77.72	94.08
Other races	26	23	21	80.77	91.3
Subtotal	957	763	686	71.68	89.91
Heterosexual contact (female)					
Black/African American	844	715	642	76.07	89.79
Hispanic/Latino ^b	714	596	543	76.05	91.11
White	321	270	253	78.82	93.7
Other races	40	34	29	72.5	85.29
Subtotal	1919	1615	1467	76.45	90.84
Total	10344	8267	7570	73.18	91.57

*Includes Asian/Pacific Islander legacy cases.

^bHispanics/Latinos can be of any race.

^cHeterosexual contact with person known to have, or to be at high risk for, HIV infection.

^dIncludes hemophilia, blood transfusion, and persons (152 cases) who were exposed perinatally but were aged >13 at most recent year.

^ePersons who have most recent viral load test result <200 copies/ml between 01/01/2018 through 12/31/2018 are considered as HIV viral suppression.

^fThe overall population is overestimated because cases are only followed up for 11 months after 12/31/2018. CDC suggests that every case should be followed up at least 18 months to collect death certificate information.