Connecticut Behavioral Risk Factor Survey

Prevalence Estimates for Risk Factors and Health Indicators

Selected Summary Tables 2018



October, 2019

State of Connecticut Department of Public Health





Acknowledgments

Diane Aye, MPH, PhD

Section Chief

Health Statistics and Surveillance Connecticut

Connecticut Department of Public Health

Celeste Jorge, MPH
CT BRFSS Coordinator
Health Statistics and Surveillance Section Connecticut
Department of Public Health

This report was prepared by:
Xi Zheng, MPH, MS
BRFSS Epidemiologist
Health Statistics and Surveillance Section
Connecticut Department of Public Health

The CT BRFSS team acknowledges with gratitude the time contributed by over 10,000 citizen volunteers within the State of Connecticut who responded anonymously to the survey during the 2018 calendar year. The results presented in this report would not be possible without their participation.

For questions or comments about this report, please contact: Xi Zheng, MPH, MS
Epidemiologist
Health Statistics and Surveillance Section
Connecticut Department of Public Health
Hartford, Connecticut, 06106
Xi.Zheng@ct.gov (860-509-7670)

Find more BRFSS factsheets, reports and publications at the Connecticut Department of Public Health BRFSS website: http://www.ct.gov/dph/BRFSS.

Introduction

The Connecticut Behavioral Risk Factor Surveillance System (CT BRFSS) is an ongoing statewide voluntary phone survey of Connecticut citizen volunteers aged 18 and over. It is funded by the Centers for Disease Control and Prevention (CDC) in all 50 states, and has been implemented in Connecticut since 1989.

For the 2018 CT BRFSS, the landline sample was a disproportionate stratified random digit dial (RDD) sample, stratified by geography and listed status. Within each contacted household, one adult was selected at random to be interviewed. The cell phone sample was an un-stratified RDD sample drawn from dedicated cellular telephone banks with equal probability. An adult contacted by cell phone was eligible to complete the survey if he or she lived in a private residence or college housing either without a landline present, or with a landline but with at least 90 percent of all calls received by cell phone. Each month, survey data from Connecticut are sent to CDC for editing and checking. At the end of each year, data are compiled and weighted to be representative of all adults in the state, and returned to states for analysis and use in planning and monitoring health programs. Summary data for all states are available on the CDC BRFSS website (http://www.cdc.gov/BRFSS).

In 2018, the total sample size of completed interviews for the core survey was 10,709 (land line=5,514; cell phone=5,195). Due to variability in response rates to individual questions and differences in specific sub-populations of interest for particular questions, not all estimates are based on a total sample size of 10,588. Prevalence estimates and 95% confidence intervals were computed using SAS PROC SURVEYFREQ, which can properly compute variances for complex sampling plans. Any responses of "Not Known/Not sure" or "Refused" were classified as missing.

In an effort to provide the most reliable information possible, coefficients of variation (CV) were used to assess the validity of each estimate. Prevalence estimates with a CV of between 15% and 20%, inclusive, are marked with a single asterisk (*). Prevalence estimates with a CV between 20.1% and 30%, inclusive, are rounded to the nearest five percent, and are marked with two asterisks (**) to indicate that caution should be exercised when interpreting these estimates. Prevalence estimates with a CV greater than 30% are suppressed due to poor validity. Only comparisons between estimates with non-overlapping 95% confidence intervals should be considered significantly different.

Due to the BRFSS methodology changes that took place in 2011, the estimates provided within the following tables should only be compared to CT BRFSS estimates from 2011-2018 and not to CT BRFSS estimates from years prior to 2011.

If you have any questions regarding any of the estimates within the following tables, please contact Xi Zheng at xi.zheng@ct.gov. To access to the 2018 CT BRFSS questionnaire, please visit the CT BRFSS web page at: www.ct.gov/dph/brfss.

Table of Contents

Health Indicator Summary	1
Adult Health	4
Health Status	4
Risk Behaviors	12
Clinical Preventive Practice	18
Chronic Diseases	28
Other Topics	33
Child Health	38

HEALTH INDICATOR SUMMARY

Adult Health Indicators	%	95%	6 CL
Health Status	1		
Fair or poor health	13.8	12.9	14.7
Poor physical health	10.5	9.7	11.2
Poor mental health	11.0	10.1	11.9
Disability	23.2	22.0	24.3
Overweight	37.0	35.6	38.4
Obese	27.4	26.2	28.6
Insurance coverage (all ages)	92.5	91.7	93.3
Private insurance (18-64 years old)	64.1	62.5	65.7
Medicare (18-64 years old)	5.6	4.9	6.4
Medicaid (18-64 years old)	11.8	10.7	12.9
No insurance (18-64 years old)	9.1	8.1	10.1
No dental insurance	26.0	24.4	27.7
At least one doctor	84.2	83.1	85.3
Not see doctor because of cost	9.2	8.4	10.0
Not get prescribed medicine because of cost	7.2	6.5	7.9
Food insecurity	7.7	6.5	9.0
Housing insecurity	13.1	11.5	14.6
Risk Behaviors	<u> </u>	L	L
No leisure time physical activities	22.4	21.2	23.5
Current smoker	12.2	11.2	13.1
Ever use E-cigar	19.6	18.4	20.8
Binge drinking	17.1	15.9	18.2
Heavy drinking	6.4	5.8	7.1
Excessive alcohol drinking	18.8	17.6	19.9
Always or nearly always wear a seatbelt	95.0	94.4	95.7
Drinking and driving	3.2	2.6	3.8
Have fallen in the past 12 months (45+)	21.6	20.4	22.7
 Injured among adults fell in the past 12 months (45+) 	39.9	36.9	43.0
Insufficient sleep (less than 8 hours)	66.0	64.7	67.3
Clinical Preventive Practice		•	
Routine Check-up in past year	81.1	80.0	82.2
Had flu shot in past year (all ages)	35.0	33.7	36.2
Had flu shot in past year (65+)	55.3	53.2	57.5
Ever had pneumonia shot (all ages)	32.0	30.8	33.3
Ever had pneumonia shot (65+)	72.4	70.4	74.4
Ever tested for HIV	39.1	37.7	40.4
Dentist visit in past year	76.5	75.4	77.7
Ever had permanent teeth extracted	37.3	36.1	38.6
Had a mammogram in the past two years (women 50-74 years old)	83.0	81.0	85.0
Ever had a pap test (21+ women)	93.8	92.6	94.9
Had Pap test within past 3 years (women 21-65 years old)	84.0	82.0	86.0
Women ever had HPV test	50.3	48.1	52.5
Had PSA test in past 2 years (Men 40+)	33.8	31.8	35.8
Ever talked about advantages of PSA test (Men 40+)	45.4	43.2	47.5
Ever talked about disadvantages of PSA test (Men 40+)	21.1	19.4	22.9
Ever received recommendation to get PSA test (Men 40+)	40.2	38.1	42.3
Had blood stool test within the past 3 years (50-75 years old)	14.8	13.6	16.0

HEALTH INDICATOR SUMMARY

Ever tested or Hepatitis C	Adult Health Indicators	%	95%	6 CL
Ever tack the HPV vaccination (18-49 years old) 21.4 19.1 23.5 Ever tested or Hepatitis C 28.6 26.6 30.6 Had blood sugar test in past 3 years 56.6 54.5 58.7 Chronic Diseases Current asthma 10.3 9.5 11.1 Former asthma 4.5 3.9 5.1 COPD 5.3 4.8 5.9 5.1 CARDIO Security 18 years 26.0 24.9 27.1 CARDIO Security 18 years 26.0 22.9 3.0 May 18 years 26.0 2.2 3.0 May 18 years 26.0 2.2 3.0 May 18 years 26.0 2.2	Clinical Preventive Practice (Cont.)			
Ever had the HPV vaccination (18-49 years old)	Ever had sigmoidoscopy/ colonoscopy (50-75 years old)	78.1	76.6	79.6
Current asthma		21.4	19.1	23.7
Current asthma	Ever tested or Hepatitis C	28.6	26.6	30.6
Current asthma	Had blood sugar test in past 3 years	56.6	54.5	58.7
Former asthma	Chronic Diseases			
COPD	Current asthma	10.3	9.5	11.1
Arthritis	Former asthma	4.5	3.9	5.1
Cardiovascular diseases 7.5 6.9 8.1	COPD	5.3	4.8	5.9
Cardiovascular diseases 7.5 6.9 8.1	Arthritis	26.0	24.9	27.1
Diabetes Serial taken class in diabetes management among adults diagnosed with diabetes Serial diabetes Carlo Carlo Serial diabetes Carlo Ca			6.9	
■ Ever taken class in diabetes management among adults diagnosed with diabetes 52.1 46.6 57.5 Pre-diabetes 10.6 9.4 11.8 Kidney disease 2.6 2.2 3.0 Depression 15.5 14.5 16.5 Periodontal disease 11.7 10.6 12.8 • Treatment for periodontal disease among adults diagnosed with periodontal disease 17.4 16.1 18.8 • Treatment for periodontal disease Other Topics Suicide Prevention Suicide Attempt among adults indicated ever thought of suicide 30.5 23.8 37.3 Awareness of suicide campaign "1 WORD, 1 VOICE, 1 LIFE" 35.4 30.5 23.8 37.3 Pages and Anxiety Felt down, depressed or hopeless for more than half the days or nearly everyday in past 2 weeks 8.6 7.7 9.5 Past 2 weeks Water Source Water Source Public water 69.8 68.0 71.6 Private well water in the pa				
Ridney disease 2.6 2.2 3.0 Depression 15.5 14.5 16.5 Periodontal disease 11.7 10.6 12.8 Treatment for periodontal disease among adults diagnosed with periodontal disease 11.7 10.6 12.8 Treatment for periodontal disease among adults diagnosed with periodontal disease 16.1 18.8 Periodontal disease 16.1 16.1 18.8 Depression 12.4 10.9 14.0 Suicide Prevention 12.4 10.9 14.0 Suicide Attempt among adults indicated ever thought of suicide 30.5 23.8 37.3 Awareness of suicide campaign "1 WORD, 1 VOICE, 1 LIFE" 35.4 33.4 33.4 33.4 Depression and Anxiety 5.5 Felt down, depressed or hopeless for more than half the days or nearly everyday in past 2 weeks 11.2 10.2 12.2 past 2 weeks 11.2 10.2 12.2 Public water 69.8 68.0 71.6 Private well water 69.8 68.0 71.6 Private well water 69.8 68.0 71.6 Private well water in the past year among adults have private well 28.0 24.2 31.8 Awareness of official advice for well water testing in CT 35.5 31.8 39.1 Respiratory Health 13.2 11.9 14.5 Cough up phlegm or mucus on most days during the past 3 months 13.2 11.9 14.5 Cough up phlegm or mucus on most days during the past 3 months 6.8 5.8 7.7 Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs 28.0 26.3 29.7 Hepatitis C Testing 28.6 26.6 30.6 Had last hepatitis C testing at private doctor or HMO office among 57.2 53.0 61.3 Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6 Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6	 Ever taken class in diabetes management among adults diagnosed with 	1		57.5
Depression 15.5 14.5 16.5 Periodontal disease 11.7 10.6 12.8 Periodontal disease 11.7 10.6 12.8 Periodontal disease 11.7 10.6 12.8 Periodontal disease 11.7 16.1 16.1 18.8 Periodontal disease 11.7 16.1 16.1 16.1 18.8 Periodontal disease 11.7 16.1 16.1 18.8 Periodontal disease 11.7 16.1 16.1 16.1 18.8 Periodontal disease 11.7 16.1 16.1 18.8 Periodontal disease 11.7 16.1 16.1 16.1 16.1 16.1 16.1 16.1	Pre-diabetes	10.6	9.4	11.8
Periodontal disease Treatment for periodontal disease among adults diagnosed with periodontal disease Other Topics Suicide Prevention Suicide ideation Suicide Attempt among adults indicated ever thought of suicide attempt as 3.4 3.4 3.4 3.4 3.4 3.4 3.4 3.4 3.4 3.4	Kidney disease	2.6	2.2	3.0
Treatment for periodontal disease among adults diagnosed with periodontal disease Other Topics Suicide Prevention Suicide ideation	Depression	15.5	14.5	16.5
Deteriod Prevention 12.4 10.9 14.0	Periodontal disease	11.7	10.6	12.8
Suicide Prevention Suicide ideation 12.4 10.9 14.0 Suicide Attempt among adults indicated ever thought of suicide 30.5 23.8 37.3 Awareness of suicide campaign "1 WORD, 1 VOICE, 1 LIFE" 35.4 33.4 37.4 Depression and Anxiety Felt down, depressed or hopeless for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks 2 11.2 11.2 11.2 12.2 12.2 12.2 12.2 12.2 12.2 12.2 12.2 12.2 12.2 12.2 28.6 12.5 12.5 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.6 12.7 12.7 12.0 12.2 12.	· · · · · · · · · · · · · · · · · · ·	17.4	16.1	18.8
Suicide ideation	Other Topics			
Suicide Attempt among adults indicated ever thought of suicide 30.5 23.8 37.3 Awareness of suicide campaign "1 WORD, 1 VOICE, 1 LIFE" 35.4 33.4 37.4 Depression and Anxiety Felt down, depressed or hopeless for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Water Source Public water 69.8 68.0 71.6 Private well water 69.8 68.0 71.6 Private well water 26.7 25.0 28.4 * Tested well water in the past year among adults have private well 28.0 24.2 31.8 Awareness of official advice for well water testing in CT 35.5 31.8 39.1 Respiratory Health Had a cough on most days during the past 3 months 13.2 11.9 14.5 Cough up phlegm or mucus on most days during the past 3 months 6.8 5.8 7.7 Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems 28.0 26.3 29.7 Hepatitis C Testing Ever tested for hepatitis C * Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6	Suicide Prevention			
Awareness of suicide campaign "1 WORD, 1 VOICE, 1 LIFE" Peptression and Anxiety Felt down, depressed or hopeless for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks	Suicide ideation	12.4	10.9	14.0
Awareness of suicide campaign "1 WORD, 1 VOICE, 1 LIFE" Peptression and Anxiety Felt down, depressed or hopeless for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks Felt nervous, anxious or nearly everyday in past 2 weeks	 Suicide Attempt among adults indicated ever thought of suicide 	30.5	23.8	37.3
Felt down, depressed or hopeless for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Water Source Public water Public water 169.8 68.0 71.6 Private well water 26.7 25.0 28.4 1 Tested well water in the past year among adults have private well 28.0 24.2 31.8 Awareness of official advice for well water testing in CT 35.5 31.8 39.1 Respiratory Health Had a cough on most days during the past 3 months 13.2 11.9 14.5 Cough up phlegm or mucus on most days during the past 3 months 6.8 5.8 7.7 Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems 28.0 26.3 29.7 Hepatitis C Testing Ever tested for hepatitis C 1 Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6		35.4	33.4	37.4
Felt down, depressed or hopeless for more than half the days or nearly everyday in past 2 weeks Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Water Source Public water 69.8 68.0 71.6 Private well water 26.7 25.0 28.4 Tested well water in the past year among adults have private well 28.0 24.2 31.8 Awareness of official advice for well water testing in CT 35.5 31.8 39.1 Respiratory Health Had a cough on most days during the past 3 months 13.2 11.9 14.5 Cough up phlegm or mucus on most days during the past 3 months 6.8 5.8 7.7 Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems 28.0 26.3 29.7 Hepatitis C Testing Ever tested for hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C 1.5* 0.9 2.1 Biagnosed with hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6			l	l
Felt nervous, anxious or on edge for more than half the days or nearly everyday in past 2 weeks Water Source Public water 69.8 68.0 71.6 Private well water 26.7 25.0 28.4 Tested well water in the past year among adults have private well 28.0 24.2 31.8 Awareness of official advice for well water testing in CT 35.5 31.8 39.1 Respiratory Health Had a cough on most days during the past 3 months 13.2 11.9 14.5 Cough up phlegm or mucus on most days during the past 3 months 6.8 5.8 7.7 Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems 28.0 26.3 29.7 Hepatitis C Testing Ever tested for hepatitis C testing at private doctor or HMO office among adults diagnosed with hepatitis C 1.5* 0.9 2.1 Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6		8.6	7.7	9.5
Water SourcePublic water69.868.071.6Private well water26.725.028.4■ Tested well water in the past year among adults have private well28.024.231.8Awareness of official advice for well water testing in CT35.531.839.1Respiratory Health35.531.839.1Had a cough on most days during the past 3 months13.211.914.5Cough up phlegm or mucus on most days during the past 3 months6.85.87.7Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs17.516.019.0Ever been given a breathing test to diagnose breathing problems28.026.329.7Hepatitis C TestingEver tested for hepatitis C28.626.630.6■ Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C57.253.061.3Diagnosed with hepatitis C1.5*0.92.1■ Ever treatment for hepatitis C among adults diagnosed with hepatitis C61.2*40.781.6	Felt nervous, anxious or on edge for more than half the days or nearly everyday in	11.2	10.2	12.2
Public water 69.8 68.0 71.6 Private well water 26.7 25.0 28.4 Tested well water in the past year among adults have private well 28.0 24.2 31.8 Awareness of official advice for well water testing in CT 35.5 31.8 39.1 Respiratory Health Had a cough on most days during the past 3 months 13.2 11.9 14.5 Cough up phlegm or mucus on most days during the past 3 months 6.8 5.8 7.7 Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems 28.0 26.3 29.7 Hepatitis C Testing Ever tested for hepatitis C 28.6 26.6 30.6 Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C 3.1.5* 0.9 2.1 Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6	past 2 weeks			
Private well water Tested well water in the past year among adults have private well Awareness of official advice for well water testing in CT Respiratory Health Had a cough on most days during the past 3 months Cough up phlegm or mucus on most days during the past 3 months Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems Ever tested for hepatitis C Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 1.5* 28.0 25.0 28.4 25.0 24.2 31.8 39.1 14.5 15.0 16.0 17.5 16.0 19.0 19.0 19.0 19.0 19.0 19.0 19.0 19.0 19.0 19.0 10.0	Water Source			
Tested well water in the past year among adults have private well 28.0 24.2 31.8 Awareness of official advice for well water testing in CT 35.5 31.8 39.1 Respiratory Health Had a cough on most days during the past 3 months 13.2 11.9 14.5 Cough up phlegm or mucus on most days during the past 3 months 6.8 5.8 7.7 Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems 28.0 26.3 29.7 Hepatitis C Testing Ever tested for hepatitis C Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6	Public water	69.8	68.0	71.6
Awareness of official advice for well water testing in CT Respiratory Health Had a cough on most days during the past 3 months Cough up phlegm or mucus on most days during the past 3 months Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems Ever tested for hepatitis C Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 35.5 31.8 39.1 39.1 35.5 31.8 39.1 35.5 31.8 39.1 Augustation 14.5 5.8 7.7 16.0 19.0 28.0 26.3 29.7 40.7 61.3	Private well water	26.7	25.0	28.4
Respiratory Health Had a cough on most days during the past 3 months Cough up phlegm or mucus on most days during the past 3 months Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems Ever tested for hepatitis C Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 1.5* 0.9 2.1	 Tested well water in the past year among adults have private well 	28.0	24.2	31.8
Had a cough on most days during the past 3 months Cough up phlegm or mucus on most days during the past 3 months Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems Ever tested for hepatitis C Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 11.5* 12.0 13.2 11.9 14.5 6.8 5.8 7.7 16.0 19.0 26.3 29.7 26.3 29.7 28.6 26.6 30.6 30.6 30.6 1.5* 0.9 2.1	Awareness of official advice for well water testing in CT	35.5	31.8	39.1
Cough up phlegm or mucus on most days during the past 3 months Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems Ever tested for hepatitis C Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 68.8 7.7 16.0 19.0 28.0 26.3 29.7 28.6 26.6 30.6 30.6 1.5* 0.9 2.1	Respiratory Health			
Have shortness of breath either when hurrying on level ground or when walking up a slight hill or stairs Ever been given a breathing test to diagnose breathing problems Ever tested for hepatitis C Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 1.5* 16.0 19.0	Had a cough on most days during the past 3 months	13.2	11.9	14.5
slight hill or stairs Ever been given a breathing test to diagnose breathing problems 28.0 26.3 29.7 Hepatitis C Testing Ever tested for hepatitis C Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 81.6 26.6 30.6 1.5* 0.9 2.1	Cough up phlegm or mucus on most days during the past 3 months	6.8	5.8	7.7
Ever been given a breathing test to diagnose breathing problems 28.0 26.3 29.7 **Hepatitis C Testing** Ever tested for hepatitis C 28.6 26.6 30.6 **Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C 1.5* 0.9 2.1 **Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6	Have shortness of breath either when hurrying on level ground or when walking up a	17.5	16.0	19.0
Ever been given a breathing test to diagnose breathing problems 28.0 26.3 29.7 **Hepatitis C Testing** Ever tested for hepatitis C 28.6 26.6 30.6 **Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C 1.5* 0.9 2.1 **Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6	the state of the s			
Hepatitis C TestingEver tested for hepatitis C28.626.630.6Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C57.253.061.3Diagnosed with hepatitis C1.5*0.92.1Ever treatment for hepatitis C among adults diagnosed with hepatitis C61.2*40.781.6		28.0	26.3	29.7
Ever tested for hepatitis C Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 28.6 57.2 53.0 61.3 61.3* 0.9 2.1 81.6				
 Had last hepatitis C testing at private doctor or HMO office among adults ever tested for hepatitis C Diagnosed with hepatitis C Ever treatment for hepatitis C among adults diagnosed with hepatitis C 57.2 53.0 61.3 	Ever tested for hepatitis C	28.6	26.6	30.6
Diagnosed with hepatitis C 1.5* 0.9 2.1 Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6	 Had last hepatitis C testing at private doctor or HMO office among 	57.2		61.3
■ Ever treatment for hepatitis C among adults diagnosed with hepatitis C 61.2* 40.7 81.6	·	1.5*	0.9	2.1
 Received hepatitis C treatment from Specialist (ID or GI medical doc) among adults diagnosed with hepatitis C 	 Received hepatitis C treatment from Specialist (ID or GI medical doc) 	40.0**	-	-

HEALTH INDICATOR SUMMARY

Adult Health Indicators	%	95%	6 CL	
Other Topics (Cont.)				
Prescription Drug Access				
Safe storage of prescription opioid-based pain relievers	44.7	42.5	46.8	
Aware of safe disposal of prescription opioid-based pain relievers	78.2	76.4	80.0	
Safe Storage of Firearms				
Has at least one firearm kept in or around home	17.9	16.3	19.5	
 The firearm secured in a location where youth, other at-risk or 	90.7	87.8	93.6	
unauthorized persons cannot have access to it, among adults kept				
firearm in their home				
 The ammunition secured in a separate location from the firearms, 	81.7	77.9	85.5	
among adults kept firearm in their home				
Air Quality				
Heard or read about the air quality index or air quality alerts where you live	44.8	42.8	46.8	
 At least once reduced or changed outdoor activity level based on the air 	32.3	29.8	34.8	
quality index or air quality alerts, among adults heard or read about the				
air quality index or air quality alerts where they live				
Sexual Violence				
Had ever been forced to have sexual intercourse, since 18 years old	8.7	7.4	10.0	
PrEP				
Heard of PrEP but not sure what it is used for	2.9	2.4	3.4	
Heard of PrEP and know what it is used for		11.7	13.6	
Considered PrEP as a way of reducing chances of getting HIV		3.8	7.7	
Special Population				
Prevalence of LGBT population	6.1	5.4	6.9	

	%	95%	6 CL
Child Health Indicators			
Overweight (5-17 years old)	16.2	13.2	19.3
Obese (5-17 years old)	18.4	15.0	21.7
Ever breastfed	81.6	79.3	83.9
Excessive screen time (2-17 years old)	56.5	53.1	59.9
Drank soda or sugary drinks at least once daily (2-17 years old)	28.3	25.1	31.5
Ate fast food at least twice weekly (2-17 years old)	41.3	37.9	44.7
Visited dentist in past year	86.9	84.8	89.0
Ever received dental sealant on permanent teeth (5-17 years old)	47.2	43.4	51.0
Had dental decay in the past 12 months	14.3	12.0	16.7
Current asthma	9.7	8.0	11.4
Former asthma	4.6	3.3	5.8

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Fair or Poor Health

<u>Indicator Description: Among all adults, the proportion who reported that their health, in general, was either fair or poor.</u>

Demograph	Demographics		95% CI	
Total		13.8	12.9	14.7
Age	18-34 years old	8.8	7.0	10.7
	35-54 years old	12.5	10.9	14.0
	55 and over	18.9	17.5	20.3
Sex	Male	13.0	11.8	14.3
	Female	14.5	13.2	15.8
Race/	Non-Hispanic White	11.6	10.7	12.6
Ethnicity	Non-Hispanic Black	18.0	14.7	21.3
	Hispanic	21.4	18.0	24.7
	Non-Hispanic Other	10.7	7.6	13.8
Income	Less than \$35,000	27.1	24.5	29.7
	\$35,000-\$74,999	12.2	10.4	14.1
	\$75,000 and more	5.6	4.7	6.5
Insurance	Insured	13.3	12.4	14.2
	Not Insured	19.8	15.6	24.1
Disability	Disabled	37.7	35.0	40.4
	Non-disabled	6.7	5.9	7.5
Education	High School Graduate or Less	21.2	19.3	23.1
	More than High School	9.2	8.4	10.1

Poor Physical Health

<u>Indicator Description: Among all adults, the proportion who reported 14 or more days of poor physical health, which includes physical illness and injury, during the past 30 days.</u>

Demograph	Demographics		95% CI	
Total		10.5	9.7	11.2
Age	18-34 years old	6.1	4.6	7.7
	35-54 years old	9.7	8.4	11.0
	55 and over	14.4	13.2	15.6
Sex	Male	9.5	8.4	10.6
	Female	11.3	10.2	12.4
Race/	Non-Hispanic White	10.4	9.5	11.3
Ethnicity	Non-Hispanic Black	11.5	8.8	14.1
	Hispanic	11.1	8.7	13.4
	Non-Hispanic Other*	6.9	4.8	9.0
Income	Less than \$35,000	18.0	16.0	20.1
	\$35,000-\$74,999	10.4	8.7	12.0
	\$75,000 and more	5.4	4.5	6.2
Insurance	Insured	10.4	9.6	11.1
	Not Insured	11.4	8.1	14.7
Disability	Disabled	29.3	26.9	31.8
	Non-disabled	5.1	4.4	5.7
Education	High School Graduate or Less	14.0	12.4	15.5
	More than High School	8.3	7.5	9.1

Poor Mental Health

<u>Indicator Description: Among all adults, the proportion who reported 14 or more days of poor mental</u>

health, during the past 30 days.

Demographics		%	95% CI	
Total		11.0	10.1	11.9
Age	18-34 years old	13.9	11.7	16.1
	35-54 years old	11.5	10.0	13.1
	55 and over	8.7	7.6	9.8
Sex	Male	9.1	7.9	10.2
	Female	12.7	11.3	14.0
Race/	Non-Hispanic White	10.3	9.3	11.3
Ethnicity	Non-Hispanic Black	15.0	11.5	18.6
	Hispanic	11.6	8.8	14.3
	Non-Hispanic Other*	10.4	6.7	14.1
Income	Less than \$35,000	17.3	15.1	19.5
	\$35,000-\$74,999	9.9	8.2	11.7
	\$75,000 and more	6.4	5.4	7.5
Insurance	Insured	10.9	10.0	11.8
	Not Insured*	12.5	8.6	16.5
Disability	Disabled	26.0	23.4	28.5
	Non-disabled	6.5	5.7	7.3
Education	High School Graduate or Less	12.7	11.0	14.3
	More than High School	9.9	8.9	11.0

Disability

Indicator Description: Among all adults, the proportion who reported being blind, or deaf, or serious difficulty in walking or climbing stairs, difficulty in dressing or bathing, or limited in any activities because of physical, mental, or emotional problems, or reported that they required use of special equipment (such as a cane, a wheelchair, a special bed, or a special telephone) due to a health problem.

Demographics		%	95%	6 CI
Total	Total		22.0	24.3
Age	18-34 years old	17.8	15.2	20.5
	35-54 years old	17.3	15.4	19.2
	55 and over	31.8	30.2	33.4
Sex	Male	21.3	19.7	22.9
	Female	24.8	23.1	26.4
Race/	Non-Hispanic White	21.6	20.3	22.8
Ethnicity	Non-Hispanic Black	26.6	22.4	30.8
	Hispanic	29.5	25.6	33.5
	Non-Hispanic Other	19.2	14.6	23.8
Income	Less than \$35,000	39.8	36.9	42.7
	\$35,000-\$74,999	21.8	19.4	24.2
	\$75,000 and more	11.8	10.3	13.2
Insurance	Insured	22.7	21.6	23.9
	Not Insured	28.5	23.2	33.8
Education	High School Graduate or Less	32.7	30.3	35.0
	More than High School	17.4	16.2	18.5

Overweight

<u>Indicator Description: Among all adults, the proportion of respondents whose BMI was greater than or equal to 25.0, but less than 30.0.</u>

Demographics		%	95% CI	
Total		37.0	35.6	38.4
Age	18-34 years old	34.0	30.7	37.4
	35-54 years old	37.3	34.9	39.7
	55 and over	38.6	36.9	40.2
Sex	Male	43.5	41.5	45.6
	Female	30.3	28.5	32.1
Race/	Non-Hispanic White	36.4	34.9	37.9
Ethnicity	Non-Hispanic Black	36.5	31.9	41.1
	Hispanic	39.5	35.2	43.9
	Non-Hispanic Other	37.4	31.2	43.6
Income	Less than \$35,000	35.7	32.8	38.6
	\$35,000-\$74,999	35.6	32.7	38.5
	\$75,000 and more	40.1	38.0	42.2
Insurance	Insured	37.1	35.7	38.5
	Not Insured	36.4	30.3	42.5
Disability	Disabled	33.0	30.3	35.8
	Non-disabled	38.1	36.6	39.7
Education	High School Graduate or Less	37.4	34.9	39.9
	More than High School	36.8	35.2	38.3

Obese

<u>Indicator Description: Among all adults, the proportion of respondents whose BMI was greater than or equal to 30.0.</u>

Demograph	Demographics		95% CI	
Total		27.4	26.2	28.6
Age	18-34 years old	20.3	17.6	22.9
	35-54 years old	32.5	30.2	34.8
	55 and over	28.8	27.2	30.4
Sex	Male	28.4	26.6	30.2
	Female	26.4	24.7	28.1
Race/	Non-Hispanic White	26.1	24.7	27.5
Ethnicity	Non-Hispanic Black	36.5	31.9	41.1
	Hispanic	31.1	27.0	35.1
	Non-Hispanic Other	18.0	13.5	22.5
Income	Less than \$35,000	31.7	28.9	34.4
	\$35,000-\$74,999	30.5	27.8	33.2
	\$75,000 and more	24.9	23.1	26.7
Insurance	Insured	27.5	26.2	28.8
	Not Insured	26.2	21.1	31.3
Disability	Disabled	36.6	33.9	39.4
	Non-disabled	24.8	23.4	26.1
Education	High School Graduate or Less	30.7	28.4	33.0
	More than High School	25.4	24.0	26.8

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Insurance Coverage

Indicator Description: Among all adults, the proportion who reported having health care coverage, including health insurance, prepaid plans such as HMOs, or government plans, such as Medicare or Indian Health Services.

Demograph	Demographics		95%	6 CI
Total		92.5	91.7	93.3
Age	18-34 years old	87.1	84.9	89.4
	35-54 years old	92.0	90.6	93.4
	55 and over	96.6	95.9	97.4
Sex	Male	91.2	89.9	92.5
	Female	93.7	92.7	94.7
Race/	Non-Hispanic White	96.6	96.0	97.3
Ethnicity	Non-Hispanic Black	88.5	85.4	91.5
	Hispanic	75.8	72.2	79.4
	Non-Hispanic Other	91.8	88.3	95.4
Income	Less than \$35,000	84.6	82.3	86.9
	\$35,000-\$74,999	92.1	90.2	93.9
	\$75,000 and more	97.8	97.0	98.5
Disability	Disabled	91.0	89.2	92.9
	Non-disabled	93.2	92.3	94.1
Education	High School Graduate or Less	86.8	85.0	88.5
	More than High School	96.1	95.4	96.8

Private Insurance

<u>Indicator Description: Among adults aged 18-64 years old, the proportion who reported currently having private insurance.</u>

Demograph	ics	%	95%	6 CI
Total		64.1	62.5	65.7
Age	18-34 years old	54.6	51.3	57.9
	35-54 years old	69.0	66.8	71.2
	55 and over	70.5	68.1	72.9
Sex	Male	64.7	62.4	67.0
	Female	63.5	61.3	65.8
Race/	Non-Hispanic White	74.6	72.8	76.4
Ethnicity	Non-Hispanic Black	47.0	41.9	52.0
	Hispanic	38.6	34.3	43.0
	Non-Hispanic Other	58.5	51.8	65.1
Income	Less than \$35,000	20.8	17.8	23.8
	\$35,000-\$74,999	67.7	64.1	71.3
	\$75,000 and more	89.7	88.2	91.2
Disability	Disabled	38.3	34.5	42.1
	Non-disabled	70.7	69.0	72.4
Education	High School Graduate or Less	45.9	42.8	48.9
	More than High School	74.7	73.0	76.4

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Medicare

<u>Indicator Description: Among adults aged 18-64 years old, the proportion who reported currently having Medicare.</u>

Demographics		%	95% CI	
Total		5.6	4.9	6.4
Age	18-34 years old*	4.5	3.1	5.9
	35-54 years old	5.0	4.0	6.1
	55 and over	8.4	6.9	9.8
Sex	Male	4.6	3.7	5.5
	Female	6.6	5.4	7.8
Race/	Non-Hispanic White	4.0	3.3	4.7
Ethnicity	Non-Hispanic Black*	10.6	7.4	13.7
	Hispanic*	8.2	5.7	10.7
	Non-Hispanic Other**	5.0	-	-
Income	Less than \$35,000	15.5	13.0	18.1
	\$35,000-\$74,999*	4.2	2.8	5.6
	\$75,000 and more*	0.7	0.5	1.0
Disability	Disabled	15.5	12.8	18.1
	Non-disabled	3.1	2.4	3.7
Education	High School Graduate or Less	9.2	7.5	10.8
	More than High School	3.6	2.9	4.3

Medicaid

<u>Indicator Description: Among adults aged 18-64 years old, the proportion who reported currently having Medicaid.</u>

Demograph	ics	%	95%	6 CI
Total		11.8	10.7	12.9
Age	18-34 years old	15.1	12.7	17.5
	35-54 years old	10.8	9.3	12.4
	55 and over	8.4	6.9	9.8
Sex	Male	9.1	7.7	10.5
	Female	14.5	12.7	16.2
Race/	Non-Hispanic White	9.2	7.9	10.4
Ethnicity	Non-Hispanic Black	20.3	16.3	24.3
	Hispanic	16.4	13.2	19.7
	Non-Hispanic Other*	13.7	8.8	18.5
Income	Less than \$35,000	29.9	26.6	33.2
	\$35,000-\$74,999	10.9	8.3	13.5
	\$75,000 and more*	1.3	0.8	1.8
Disability	Disabled	21.2	17.8	24.6
	Non-disabled	9.7	8.6	10.9
Education	High School Graduate or Less	16.9	14.5	19.2
	More than High School	8.9	7.8	10.1

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

No Insurance

<u>Indicator Description:</u> Among adults aged 18-64 years old, the proportion who reported currently having no health insurance.

Demograph	Demographics		95%	6 CI
Total		9.1	8.1	10.1
Age	18-34 years old	12.9	10.6	15.1
	35-54 years old	8.0	6.6	9.4
	55 and over	5.0	3.7	6.4
Sex	Male	10.5	8.9	12.0
	Female	7.8	6.4	9.1
Race/	Non-Hispanic White	4.1	3.2	5.0
Ethnicity	Non-Hispanic Black	11.9	8.5	15.2
	Hispanic	25.7	21.9	29.5
	Non-Hispanic Other**	10.0	ı	-
Income	Less than \$35,000	19.9	16.9	22.9
	\$35,000-\$74,999	10.4	8.0	12.9
	\$75,000 and more*	2.2	1.4	3.0
Disability	Disabled	13.0	10.1	15.8
	Non-disabled	7.9	6.8	9.0
Education	High School Graduate or Less	16.9	14.5	19.2
	More than High School	4.5	3.7	5.4

At Least One Regular Doctor

<u>Indicator Description: Among all adults, the proportion who reported that they have at least one person they thought of as their personal doctor or health care provider.</u>

Demographics		%	95%	6 CI
Total		84.2	83.1	85.3
Age	18-34 years old	68.5	65.4	71.6
	35-54 years old	84.9	83.1	86.7
	55 and over	94.5	93.6	95.3
Sex	Male	80.4	78.6	82.1
	Female	87.7	86.3	89.1
Race/	Non-Hispanic White	89.2	88.1	90.3
Ethnicity	Non-Hispanic Black	77.2	72.9	81.4
	Hispanic	67.2	63.1	71.2
	Non-Hispanic Other	80.4	75.5	85.3
Income	Less than \$35,000	76.0	73.3	78.8
	\$35,000-\$74,999	82.5	80.0	85.0
	\$75,000 and more	91.1	89.8	92.4
Insurance	Insured	87.5	86.4	88.5
	Not Insured	46.0	40.4	51.7
Disability	Disabled	86.2	83.8	88.5
	Non-disabled	84.0	82.7	85.3
Education	High School Graduate or Less	80.0	77.8	82.1
	More than High School	86.8	85.6	88.1

Skip Doctor Visit Due to Cost

<u>Indicator Description: Among all adults, the proportion who that in the past 12 months, they could not see a doctor when needed due to the cost.</u>

Demograph	Demographics		95% CI	
Total		9.2	8.4	10.0
Age	18-34 years old	12.1	10.0	14.1
	35-54 years old	11.8	10.2	13.4
	55 and over	5.3	4.4	6.2
Sex	Male	8.1	6.9	9.2
	Female	10.2	9.1	11.4
Race/	Non-Hispanic White	6.9	6.1	7.7
Ethnicity	Non-Hispanic Black	13.2	9.8	16.6
	Hispanic	17.9	14.7	21.1
	Non-Hispanic Other*	8.5	5.8	11.3
Income	Less than \$35,000	15.1	13.0	17.3
	\$35,000-\$74,999	10.9	9.0	12.8
	\$75,000 and more	4.9	4.0	5.9
Insurance	Insured	7.4	6.6	8.1
	Not Insured	32.2	26.9	37.4
Disability	Disabled	14.8	12.6	17.0
	Non-disabled	7.3	6.5	8.1
Education	High School Graduate or Less	11.4	9.8	13.0
	More than High School	7.8	6.9	8.7

Abandon Prescription Drug Due to Cost

<u>Indicator Description: Among all adults, the proportion who that in the past 12 months, they did not take their medication as prescribed because of cost.</u>

Demograph	Demographics		95%	6 CI
Total		7.2	6.5	7.9
Age	18-34 years old	6.0	4.5	7.6
	35-54 years old	9.2	7.7	10.6
	55 and over	6.5	5.5	7.5
Sex	Male	5.7	4.8	6.7
	Female	8.5	7.4	9.6
Race/	Non-Hispanic White	6.1	5.3	6.9
Ethnicity	Non-Hispanic Black*	8.4	5.8	11.1
	Hispanic	12.6	9.8	15.5
	Non-Hispanic Other**	5.0	-	-
Income	Less than \$35,000	12.0	10.0	14.0
	\$35,000-\$74,999	8.4	6.8	10.1
	\$75,000 and more	4.2	3.3	5.1
Insurance	Insured	6.6	5.9	7.3
	Not Insured	14.7	10.6	18.8
Disability	Disabled	14.6	12.5	16.7
	Non-disabled	4.9	4.2	5.7
Education	High School Graduate or Less	8.9	7.5	10.3
	More than High School	6.1	5.3	6.9

Food Insecurity

<u>Indicator Description: Among all adults, the proportion who "always" or "usually" felt worried or stressed about having enough money to buy nutritious meals.</u>

Demograph	Demographics		95%	6 CI
Total		7.7	6.5	9.0
Age	18-34 years old*	9.9	6.7	13.1
	35-54 years old	10.9	8.4	13.4
	55 and over	4.1	3.1	5.1
Sex	Male	6.8	5.0	8.5
	Female	8.7	6.9	10.4
Race/	Non-Hispanic White	5.7	4.6	6.9
Ethnicity	Non-Hispanic Black**	10.0	-	-
	Hispanic*	17.3	11.8	22.8
	Non-Hispanic Other	-	-	-
Income	Less than \$35,000	17.6	14.0	21.3
	\$35,000-\$74,999*	6.6	4.3	9.0
	\$75,000 and more**	<0.1	-	-
Insurance	Insured	7.0	5.8	8.3
msurance	Not Insured**	20.0	ı	-
Disability	Disabled	14.5	11.4	17.5
	Non-disabled	5.5	4.3	6.8
Education	High School Graduate or Less	10.8	8.4	13.2
	More than High School	5.9	4.6	7.3

Housing Insecurity

<u>Indicator Description: Among all adults, the proportion who "always" or "usually" felt worried or</u> stressed about having enough money to pay for housing.

Demograph	ics	%	95%	6 CI
Total		13.1	11.5	14.6
Age	18-34 years old	15.3	11.0	19.7
	35-54 years old	16.2	13.3	19.2
	55 and over	9.6	7.9	11.3
Sex	Male	10.6	8.6	12.7
	Female	15.2	12.9	17.5
Race/	Non-Hispanic White	10.8	9.2	12.4
Ethnicity	Non-Hispanic Black*	21.9	14.9	28.9
	Hispanic*	19.5	13.7	25.3
	Non-Hispanic Other**	10.0	ı	-
Income	Less than \$35,000	25.3	21.1	29.4
	\$35,000-\$74,999	12.7	9.5	15.8
	\$75,000 and more*	4.8	3.4	6.3
Insurance	Insured	12.4	10.8	13.9
	Not Insured*	23.2	15.5	30.9
Disability	Disabled	23.8	19.9	27.7
	Non-disabled	9.6	8.1	11.2
Education	High School Graduate or Less	17.4	14.3	20.5
	More than High School	10.6	8.9	12.3

No Leisure Time Physical Activities

<u>Indicator Description: Among all adults, the proportion who reported not participating in any leisure</u> <u>time physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise during the past month.</u>

Demographics		%	95%	6 CI
Total	Total		21.2	23.5
Age	18-34 years old	18.2	15.5	20.8
	35-54 years old	20.9	19.0	22.9
	55 and over	26.9	25.3	28.5
Sex	Male	20.0	18.4	21.6
	Female	24.6	22.9	26.2
Race/	Non-Hispanic White	20.0	18.8	21.3
Ethnicity	Non-Hispanic Black	30.4	26.1	34.6
	Hispanic	30.4	26.5	34.3
	Non-Hispanic Other	17.3	12.6	22.1
Income	Less than \$35,000	35.4	32.6	38.2
	\$35,000-\$74,999	22.1	19.7	24.6
	\$75,000 and more	13.2	11.8	14.6
Insurance	Insured	21.5	20.4	22.7
	Not Insured	32.7	27.4	38.0
Disability	Disabled	38.7	35.9	41.4
	Non-disabled	17.5	16.2	18.7
Education	High School Graduate or Less	33.7	31.4	36.0
	More than High School	15.4	14.3	16.6

Current Smoker

<u>Indicator Description:</u> Among all adults, the proportion who reported that they had ever smoked at least 100 cigarettes in their life and that they smoke cigarettes now, either every day or on some days.

Demograph	Demographics		95%	6 CI
Total		12.2	11.2	13.1
Age	18-34 years old	14.1	11.7	16.5
	35-54 years old	13.7	12.0	15.3
	55 and over	10.0	8.9	11.1
Sex	Male	13.6	12.1	15.1
	Female	10.8	9.6	12.0
Race/	Non-Hispanic White	10.8	9.8	11.8
Ethnicity	Non-Hispanic Black	18.2	14.6	21.9
	Hispanic	16.5	13.2	19.9
	Non-Hispanic Other*	8.7	5.7	11.6
Income	Less than \$35,000	21.2	18.7	23.6
	\$35,000-\$74,999	12.2	10.1	14.2
	\$75,000 and more	7.4	6.3	8.5
Insurance	Insured	11.6	10.6	12.5
	Not Insured	19.7	14.6	24.8
Disability	Disabled	18.2	16.0	20.4
	Non-disabled	10.3	9.3	11.4
Education	High School Graduate or Less	19.0	17.0	20.9
	More than High School	8.0	7.1	8.9

Ever Use E-cigarette

<u>Indicator Description:</u> Among all adults, the proportion who reported that they ever used e-cigarettes or other electronic vaping products, either every day or on some days.

Demograph	Demographics		95%	6 CI
Total		19.6	18.4	20.8
Age	18-34 years old	38.1	34.7	41.5
	35-54 years old	18.9	17.0	20.8
	55 and over	8.4	7.5	9.3
Sex	Male	23.0	21.2	24.8
	Female	16.5	15.0	18.1
Race/	Non-Hispanic White	18.9	17.5	20.3
Ethnicity	Non-Hispanic Black	20.8	17.1	24.6
	Hispanic	23.0	19.2	26.8
	Non-Hispanic Other	19.6	14.8	24.5
Income	Less than \$35,000	23.0	20.4	25.6
	\$35,000-\$74,999	20.3	17.8	22.9
	\$75,000 and more	18.0	16.1	19.8
Insurance	Insured	19.5	18.3	20.7
	Not Insured	21.3	16.2	26.4
Disability	Disabled	22.2	19.7	24.6
	Non-disabled	18.8	17.4	20.2
Education	High School Graduate or Less	20.5	18.3	22.6
	More than High School	19.2	17.8	20.6

Binge Drinking

<u>Indicator Description:</u> Among all adults, the proportion who reported consuming five or more drinks per <u>occasion</u> (for men) or four or more (for women) at least once in the previous month.

Demographics		%	95% CI	
Total		17.1	15.9	18.2
Age	18-34 years old	27.4	24.3	30.4
	35-54 years old	18.9	16.9	20.8
	55 and over	9.2	8.1	10.3
Sex	Male	21.4	19.5	23.2
	Female	13.2	11.8	14.5
Race/	Non-Hispanic White	17.4	16.0	18.7
Ethnicity	Non-Hispanic Black	12.3	9.3	15.4
	Hispanic	20.7	17.0	24.4
	Non-Hispanic Other*	14.1	9.2	19.0
Income	Less than \$35,000	15.2	12.8	17.6
	\$35,000-\$74,999	16.4	13.9	18.8
	\$75,000 and more	21.5	19.6	23.4
Insurance	Insured	16.5	15.3	17.7
	Not Insured	23.6	18.4	28.8
Disability	Disabled	13.9	11.8	16.0
	Non-disabled	18.1	16.7	19.4
Education	High School Graduate or Less	16.6	14.5	18.6
	More than High School	17.5	16.1	18.8

Heavy Drinking

<u>Indicator Description: Among all adults, the proportion who reported consuming an average of more than two alcoholic drinks per day for men, or more than one per day for women in the previous month.</u>

Demographics		%	95%	6 CI
Total	Total		5.8	7.1
Age	18-34 years old	6.4	4.8	8.0
	35-54 years old	7.2	5.9	8.5
	55 and over	6.1	5.2	7.0
Sex	Male	6.1	5.1	7.1
	Female	6.8	5.9	7.8
Race/	Non-Hispanic White	7.2	6.4	8.1
Ethnicity	Non-Hispanic Black**	5.0	-	-
	Hispanic**	5.0	-	-
	Non-Hispanic Other**	10.0	1	-
Income	Less than \$35,000	5.3	4.0	6.6
	\$35,000-\$74,999	5.8	4.4	7.3
	\$75,000 and more	8.5	7.2	9.7
Insura	Insured	6.4	5.6	7.1
	Not Insured**	5.0	-	-
Disability	Disabled	6.1	4.8	7.5
	Non-disabled	6.6	5.8	7.4
Education	High School Graduate or Less	6.0	4.7	7.3
	More than High School	6.8	6.0	7.6

Excessive Drinking

<u>Indicator Description:</u> Among all adults, the proportion of adults who drank alcohol excessively (either heavy or binge drinkers).

Demographics		%	95%	6 CI
Total	Total		17.6	19.9
Age	18-34 years old	28.2	25.2	31.3
	35-54 years old	20.4	18.4	22.5
	55 and over	11.7	10.5	12.8
Sex	Male	22.3	20.4	24.1
	Female	15.6	14.2	17.0
Race/	Non-Hispanic White	19.6	18.2	21.0
Ethnicity	Non-Hispanic Black	13.5	10.4	16.6
	Hispanic	20.7	17.1	24.4
	Non-Hispanic Other*	14.6	9.7	19.5
Income	Less than \$35,000	16.4	14.0	18.9
	\$35,000-\$74,999	17.8	15.3	20.2
	\$75,000 and more	23.9	22.0	25.9
Insurance	Insured	18.3	17.1	19.4
	Not Insured	23.9	18.7	29.0
Disability	Disabled	15.3	13.1	17.4
	Non-disabled	19.9	18.5	21.2
Education	High School Graduate or Less	17.9	15.8	19.9
	More than High School	19.4	18.0	20.8

Seatbelt Use

<u>Indicator Description: Among all adults, the proportion who reported always or nearly always used a seatbelt when driving or riding in a car.</u>

Demograph	Demographics		95% CI	
Total		95.0	94.4	95.7
Age	18-34 years old	93.9	92.2	95.6
	35-54 years old	94.8	93.6	96.0
	55 and over	96.0	95.2	96.7
Sex	Male	94.0	93.0	95.0
	Female	96.0	95.1	96.9
Race/	Non-Hispanic White	96.3	95.6	97.0
Ethnicity	Non-Hispanic Black	89.3	86.4	92.3
	Hispanic*	94.0	91.9	96.1
	Non-Hispanic Other**	90.0	-	-
Income	Less than \$35,000	91.9	90.2	93.6
	\$35,000-\$74,999*	95.6	94.2	97.1
	\$75,000 and more	96.9	96.2	97.7
Insurance	Insured	95.3	94.6	96.0
	Not Insured*	92.0	88.8	95.1
Disability	Disabled	91.6	89.5	93.6
	Non-disabled	96.1	95.5	96.7
Education	High School Graduate or Less	92.2	90.8	93.7
	More than High School	96.7	96.2	97.3

Drinking and Driving

<u>Indicator Description: Among all adults, the proportion who reported having driven at least once when perhaps had too much to drink.</u>

Demographics		%	95% CI	
Total		3.2	2.6	3.8
Age	18-34 years old*	4.5	2.7	6.2
	35-54 years old	3.5	2.5	4.5
	55 and over	2.2	1.7	2.8
Sex	Male	3.8	3.0	4.6
	Female*	2.6	1.6	3.5
Race/	Non-Hispanic White	3.7	3.0	4.5
Ethnicity	Non-Hispanic Black	-	-	-
	Hispanic	-	-	-
	Non-Hispanic Other	-	-	-
Income	Less than \$35,000**	<0.1	-	-
	\$35,000-\$74,999**	5.0	-	-
	\$75,000 and more	4.2	3.2	5.3
Insurance	Insured	3.2	2.6	3.9
	Not Insured	-	-	-
Disability	Disabled**	5.0	-	-
	Non-disabled	3.2	2.6	3.8
Education	High School Graduate or Less**	5.0	-	-
	More than High School	3.4	2.8	4.1

Fall Indicator Description: Among adults aged 45+ years old, the proportion who reported had a fall sometime in the past 12 months.

Demographics		%	95%	6 CI
Total		21.6	20.4	22.7
Age	45-54 years old	19.8	17.5	22.1
	55 and over	22.6	21.2	24.1
Sex	Male	20.2	18.5	22.0
	Female	22.6	21.1	24.2
Race/	Non-Hispanic White	22.0	20.8	23.2
Ethnicity	Non-Hispanic Black	16.6	12.1	21.1
	Hispanic	25.3	19.6	31.1
	Non-Hispanic Other*	17.9	12.0	23.8
Income	Less than \$35,000	28.2	25.2	31.2
	\$35,000-\$74,999	22.5	20.0	25.1
	\$75,000 and more	17.9	16.3	19.6
Insurance	Insured	21.7	20.5	22.9
	Not Insured*	19.8	12.0	27.5
Disability	Disabled	37.0	34.3	39.8
	Non-disabled	15.7	14.6	16.9
Education	High School Graduate or Less	22.6	20.4	24.8
	More than High School	20.9	19.6	22.2

Fall and Injured

<u>Indicator Description: Among adults aged 45 years old who had a fall in the past 12 months, the proportion who were injured during the fall.</u>

Demograph	ics	%	95%	6 CI
Total		39.9	36.9	43.0
Age	45-54 years old	39.3	32.9	45.7
	55 and over	40.1	36.6	43.6
Sex	Male	35.7	30.9	40.5
	Female	43.1	39.2	47.0
Race/	Non-Hispanic White	37.8	34.8	40.8
Ethnicity	Non-Hispanic Black*	49.5	34.4	64.7
	Hispanic	47.1	33.6	60.5
	Non-Hispanic Other**	45.0	-	-
Income	Less than \$35,000	44.0	37.8	50.2
	\$35,000-\$74,999	36.3	30.2	42.4
	\$75,000 and more	35.5	30.8	40.2
Insurance	Insured	39.5	36.5	42.5
	Not Insured**	50.0	-	-
Disability	Disabled	49.3	44.6	53.9
	Non-disabled	31.8	27.9	35.6
Education	High School Graduate or Less	44.3	38.8	49.9
	More than High School	36.8	33.4	40.2

Insufficient Sleep

<u>Indicator Description:</u> Among all adults, the proportion who reported getting on average less than eight hours of sleep per night.

Demographics		%	95% CI	
Total		66.0	64.7	67.3
Age	18-34 years old	68.0	64.8	71.1
	35-54 years old	69.4	67.1	71.6
	55 and over	61.7	60.1	63.4
Sex	Male	67.0	65.1	68.9
	Female	65.1	63.3	66.8
Race/	Non-Hispanic White	66.0	64.6	67.5
Ethnicity	Non-Hispanic Black	65.8	61.3	70.2
	Hispanic	64.0	59.8	68.1
	Non-Hispanic Other	69.9	64.1	75.7
Income	Less than \$35,000	62.1	59.3	65.0
	\$35,000-\$74,999	64.9	62.0	67.7
	\$75,000 and more	70.5	68.6	72.4
Insurance	Insured	66.4	65.1	67.7
	Not Insured	60.7	55.1	66.3
Disability	Disabled	64.9	62.3	67.5
	Non-disabled	66.6	65.1	68.1
Education	High School Graduate or Less	61.9	59.5	64.3
	More than High School	68.5	67.0	69.9

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Routine Checkup in the past year

<u>Indicator Description: Among all adults, the proportion who reported that they did not have a routine checkup in the past year.</u>

Demographics		%	95%	6 CI
Total		81.1	80.0	82.2
Age	18-34 years old	73.8	70.9	76.6
	35-54 years old	77.0	75.0	79.1
	55 and over	89.2	88.2	90.2
Sex	Male	78.8	77.1	80.5
	Female	83.3	81.8	84.7
Race/	Non-Hispanic White	82.0	80.8	83.3
Ethnicity	Non-Hispanic Black	83.9	80.3	87.4
	Hispanic	77.4	74.0	80.9
	Non-Hispanic Other	74.0	68.4	79.6
Income	Less than \$35,000	79.9	77.3	82.4
	\$35,000-\$74,999	77.1	74.4	79.8
	\$75,000 and more	82.2	80.7	83.8
Insurance	Insured	83.0	81.9	84.1
	Not Insured	59.2	53.6	64.9
Disability	Disabled	84.6	82.3	86.9
	Non-disabled	79.9	78.7	81.2
Education	High School Graduate or Less	81.5	79.6	83.5
	More than High School	80.9	79.5	82.2

Flu Shot in the past year

<u>Indicator Description: Among all adults, the proportion who reported that they had a flu shot in the past year.</u>

Demograph	Demographics		95%	% CI
Total		35.0	33.7	36.2
Age	18-34 years old	23.4	20.5	26.3
	35-54 years old	28.7	26.5	30.8
	55 and over	47.9	46.2	49.6
Sex	Male	32.1	30.3	33.9
	Female	37.6	35.9	39.4
Race/	Non-Hispanic White	38.1	36.6	39.6
Ethnicity	Non-Hispanic Black	25.9	21.9	29.9
	Hispanic	25.3	21.6	29.0
	Non-Hispanic Other	33.1	27.3	38.9
Income	Less than \$35,000	33.8	30.9	36.7
	\$35,000-\$74,999	33.4	30.7	36.1
	\$75,000 and more	37.2	35.2	39.2
Insurance	Insured	36.1	34.8	37.4
	Not Insured	21.5	16.7	26.3
Disability	Disabled	38.3	35.6	41.0
	Non-disabled	34.1	32.6	35.6
Education	High School Graduate or Less	31.8	29.6	34.1
	More than High School	36.9	35.4	38.4

Flu Shot in the past year (65+)

<u>Indicator Description:</u> Among adults 65 and older, the proportion who reported that they had a flu shot in the past year.

Demographics		%	95% CI	
Total	Total		53.2	57.5
Sex	Male	56.2	52.8	59.5
	Female	54.7	51.8	57.6
Race/	Non-Hispanic White	55.5	53.3	57.8
Ethnicity	Non-Hispanic Black	44.7	33.6	55.8
	Hispanic*	64.4	50.7	78.0
	Non-Hispanic Other*	60.3	44.7	75.8
Income	Less than \$35,000	53.0	48.3	57.7
	\$35,000-\$74,999	57.1	53.2	61.0
	\$75,000 and more	58.0	54.3	61.7
Insurance	Insured	56.0	53.8	58.2
	Not Insured	-	-	-
Disability	Disabled	59.0	55.3	62.6
	Non-disabled	53.1	50.4	55.8
Education	High School Graduate or Less	54.5	50.7	58.4
	More than High School	55.9	53.5	58.3

Ever Had Pneumonia Shot

<u>Indicator Description: Among all adults, the proportion who reported that they ever had a pneumococcal vaccine.</u>

Demographics		%	95%	6 CI
Total	Total		30.8	33.3
Age	18-34 years old	20.1	16.8	23.3
	35-54 years old	14.3	12.6	16.0
	55 and over	51.8	50.1	53.6
Sex	Male	29.9	28.0	31.8
	Female	34.0	32.2	35.7
Race/	Non-Hispanic White	36.1	34.6	37.6
Ethnicity	Non-Hispanic Black	25.1	21.0	29.3
	Hispanic	18.5	15.0	22.0
	Non-Hispanic Other	22.8	17.6	28.0
Income	Less than \$35,000	36.3	33.4	39.2
	\$35,000-\$74,999	32.8	30.2	35.4
	\$75,000 and more	27.0	25.2	28.9
Insurance	Insured	33.1	31.8	34.4
	Not Insured	18.3	13.2	23.4
Disability	Disabled	47.3	44.4	50.1
	Non-disabled	27.2	25.8	28.6
Education	High School Graduate or Less	32.9	30.6	35.2
	More than High School	31.4	29.9	32.9

Ever Had Pneumonia Shot (65+)

<u>Indicator Description: Among adults 65 years old and over, the proportion who reported that they ever had a pneumococcal vaccine.</u>

Demographics		%	95% CI	
Total	Total		70.4	74.4
Sex	Male	68.9	65.7	72.1
	Female	75.0	72.5	77.5
Race/	Non-Hispanic White	74.3	72.3	76.3
Ethnicity	Non-Hispanic Black	57.3	46.4	68.2
	Hispanic*	65.1	51.2	79.0
	Non-Hispanic Other*	55.5	38.4	72.5
Income	Less than \$35,000	69.0	64.5	73.5
	\$35,000-\$74,999	75.0	71.6	78.4
	\$75,000 and more	74.7	71.4	77.9
Insurance	Insured	73.4	71.4	75.3
	Not Insured**	25.0	ı	-
Disability	Disabled	75.6	72.4	78.9
	Non-disabled	70.3	67.8	72.8
Education	High School Graduate or Less	70.1	66.6	73.6
	More than High School	74.3	72.0	76.5

Ever Tested for HIV

<u>Indicator Description: Among all adults, the proportion who reported that they ever had been tested for HIV, apart from tests that were part of a blood donation.</u>

Demographics		%	95%	6 CI
Total	Total		37.7	40.4
Age	18-34 years old	45.6	42.1	49.0
	35-54 years old	56.3	53.9	58.8
	55 and over	21.4	20.0	22.8
Sex	Male	36.7	34.7	38.7
	Female	41.3	39.4	43.2
Race/	Non-Hispanic White	32.4	30.9	33.9
Ethnicity	Non-Hispanic Black	62.5	57.7	67.2
	Hispanic	55.7	51.2	60.2
	Non-Hispanic Other	37.4	31.3	43.6
Income	Less than \$35,000	45.6	42.5	48.6
	\$35,000-\$74,999	37.7	34.7	40.8
	\$75,000 and more	39.2	37.0	41.3
Insurance	Insured	38.6	37.2	40.0
	Not Insured	45.0	38.9	51.1
Disability	Disabled	39.9	37.0	42.8
	Non-disabled	38.6	37.0	40.2
Education	High School Graduate or Less	36.5	34.0	39.0
	More than High School	40.6	39.0	42.3

Dentist Visit in the past year

<u>Indicator Description:</u> Among all adults, the proportion who reported that they had visited a dentist or dental clinic for any reason in the previous year.

Demographics		%	95% CI	
Total		76.5	75.4	77.7
Age	18-34 years old	72.6	69.7	75.5
	35-54 years old	77.7	75.7	79.7
	55 and over	77.9	76.4	79.4
Sex	Male	75.6	73.9	77.4
	Female	77.4	75.9	79.0
Race/	Non-Hispanic White	79.5	78.2	80.8
Ethnicity	Non-Hispanic Black	69.9	65.8	74.0
	Hispanic	69.5	65.7	73.3
	Non-Hispanic Other	70.5	64.9	76.2
Income	Less than \$35,000	63.5	60.7	66.3
	\$35,000-\$74,999	74.8	72.1	77.4
	\$75,000 and more	87.1	85.6	88.5
Insurance	Insured	77.9	76.7	79.1
	Not Insured	59.6	54.0	65.1
Disability	Disabled	66.6	63.9	69.3
	Non-disabled	79.4	78.1	80.7
Education	High School Graduate or Less	69.0	66.8	71.2
	More than High School	81.1	79.8	82.5

Breast Cancer Screening (Women 50-74)

<u>Indicator Description: Among women aged 50-74 years old, the proportion women reported had a mammogram in the past two years.</u>

Demographics		%	95%	6 CI
Total	Total		81.0	85.0
Race/	Non-Hispanic White	83.5	81.4	85.5
Ethnicity	Non-Hispanic Black**	80.0	1	-
	Hispanic**	85.0	ı	-
	Non-Hispanic Other**	70.0	ı	-
Income	Less than \$35,000	76.2	71.4	80.9
	\$35,000-\$74,999	83.3	79.7	86.9
	\$75,000 and more	86.3	83.7	88.9
Insurance	Insured	83.8	81.8	85.7
	Not Insured**	65.0	-	-
Disability	Disabled	78.0	73.2	82.8
	Non-disabled	84.6	82.5	86.7
Education	High School Graduate or Less	80.5	76.5	84.6
	More than High School	84.4	82.4	86.5

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Cervical Cancer Screening (women 21+)

<u>Indicator Description: Among women aged 21 years old and over, the proportion women who ever had a pap test.</u>

Demographics		%	95%	6 CI
Total	Total		92.6	94.9
Age	18-34 years old	83.3	79.2	87.5
	35-54 years old**	95.0	-	-
	55 and over	96.5	95.8	97.3
Race/	Non-Hispanic White	95.4	94.3	96.5
Ethnicity	Non-Hispanic Black**	95.0	1	-
	Hispanic**	90.0	ı	-
	Non-Hispanic Other**	80.0	ı	-
Income	Less than \$35,000	89.7	87.0	92.3
	\$35,000-\$74,999**	95.0	ı	-
	\$75,000 and more**	100.0	ı	-
Insurance	Insured	94.6	93.6	95.6
	Not Insured**	85.0	-	-
Disability	Disabled	91.6	89.2	93.9
	Non-disabled	94.4	93.2	95.7
Education	High School Graduate or Less	92.4	90.3	94.5
	More than High School	94.6	93.3	95.9

Cervical Cancer Screening (women 21-65)

<u>Indicator Description: Among women aged 21-65 years old, the proportion women who had a pap test in last three years.</u>

Demographics		%	95%	6 CI
Total		84.0	82.0	86.0
Age	21-34 years old	78.4	73.9	83.0
	35-54 years old	88.0	85.4	90.5
	55-65 years old	84.2	81.4	87.0
Race/	Non-Hispanic White	85.6	83.4	87.7
Ethnicity	Non-Hispanic Black**	85.0	-	-
	Hispanic*	82.2	76.4	87.9
	Non-Hispanic Other*	69.4	58.9	79.9
Income	Less than \$35,000	77.6	72.9	82.4
	\$35,000-\$74,999	84.4	80.4	88.4
	\$75,000 and more	91.0	89.1	92.9
Insurance	Insured	85.6	83.6	87.5
	Not Insured*	69.0	59.0	79.0
Disability	Disabled	78.8	73.4	84.2
	Non-disabled	85.1	83.0	87.2
Education	High School Graduate or Less	79.2	74.7	83.7
	More than High School	86.1	84.0	88.1

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Ever Had HPV Test

Indicator Description: the proportion women who reported that they ever had an HPV test.

Demographics		%	95%	6 CI
Total		50.3	48.1	52.5
Age	18-34 years old	55.1	49.8	60.5
	35-54 years old	67.9	64.3	71.5
	55 and over	32.4	29.8	35.0
Race/	Non-Hispanic White	47.5	45.0	50.0
Ethnicity	Non-Hispanic Black	57.7	50.8	64.5
	Hispanic	60.2	53.3	67.1
	Non-Hispanic Other	44.1	33.6	54.7
Income	Less than \$35,000	48.0	43.6	52.5
	\$35,000-\$74,999	50.7	46.0	55.4
	\$75,000 and more	60.5	56.9	64.0
Insurance	Insured	50.5	48.2	52.7
	Not Insured	50.1	40.0	60.1
Disability	Disabled	44.5	39.8	49.2
	Non-disabled	52.2	49.7	54.7
Education	High School Graduate or Less	42.2	38.1	46.3
	More than High School	55.0	52.4	57.5

Had PSA test in past 2 years (Men 40+)

<u>Indicator Description: Among men aged 40 years and older, the proportion who reported had PSA test</u> within past 2 years.

Demographi	Demographics		95%	6 CI
Total	Total		31.8	35.8
Age	40-54 years old	15.9	13.1	18.7
	55 and over	46.3	43.6	48.9
Race/	Non-Hispanic White	37.1	34.8	39.4
Ethnicity	Non-Hispanic Black*	25.2	17.4	33.1
	Hispanic*	18.1	11.5	24.7
	Non-Hispanic Other*	28.5	18.5	38.4
Income	Less than \$35,000	26.1	21.6	30.6
	\$35,000-\$74,999	31.9	27.8	36.1
	\$75,000 and more	37.0	34.0	39.9
Insurance	Insured	35.1	33.0	37.2
	Not Insured**	10.0	-	-
Disability	Disabled	34.8	30.5	39.0
	Non-disabled	33.4	31.0	35.7
Education	High School Graduate or Less	25.8	22.3	29.4
	More than High School	38.6	36.2	41.1

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Ever Talked about Advantages of PSA Test (Men 40+)

Indicator Description: Among men aged 40 years and older, the proportion who reported ever receiving a recommendation from a doctor, nurse, or other health professional regarding advantages to get a PSA test.

Demographi	Demographics		95%	6 CI
Total	Total		43.2	47.5
Age	40-54 years old	27.2	23.8	30.6
	55 and over	57.9	55.3	60.6
Race/	Non-Hispanic White	48.4	46.0	50.7
Ethnicity	Non-Hispanic Black	40.2	31.1	49.4
	Hispanic	30.0	22.1	37.8
	Non-Hispanic Other	33.8	24.2	43.5
Income	Less than \$35,000	35.5	30.6	40.4
	\$35,000-\$74,999	45.4	40.6	50.1
	\$75,000 and more	50.2	47.1	53.3
Insurance	Insured	46.4	44.2	48.6
	Not Insured*	28.1	17.1	39.1
Disability	Disabled	45.8	41.4	50.2
	Non-disabled	45.3	42.8	47.8
Education	High School Graduate or Less	39.0	35.0	43.0
	More than High School	49.3	46.8	51.8

Ever Talked about Disadvantages of PSA Test (Men 40+)

Indicator Description: Among men aged 40 years and older, the proportion who reported ever receiving a recommendation from a doctor, nurse, or other health professional regarding disadvantages to get a PSA test.

Demograph	Demographics		95%	6 CI
Total	Total		19.4	22.9
Age	40-54 years old	13.5	10.9	16.1
	55 and over	26.4	24.1	28.7
Race/	Non-Hispanic White	21.1	19.2	22.9
Ethnicity	Non-Hispanic Black*	24.4	15.8	33.1
	Hispanic*	19.3	12.6	26.1
	Non-Hispanic Other**	15.0	ı	-
Income	Less than \$35,000	18.1	14.1	22.1
	\$35,000-\$74,999	18.8	15.4	22.2
	\$75,000 and more	24.1	21.4	26.7
Insurance	Insured	21.4	19.6	23.1
	Not Insured**	20.0	-	-
Disability	Disabled	21.7	17.8	25.6
	Non-disabled	21.0	19.0	22.9
Education	High School Graduate or Less	17.9	14.7	21.2
	More than High School	23.1	21.1	25.1

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Ever Received Recommendation to Get PSA Test (Men 40+)

<u>Indicator Description: Among adults aged 40 years and older, the proportion who reported ever receiving a recommendation from a doctor, nurse, or other health professional to get a PSA test.</u>

Demographi	Demographics		95%	6 CI
Total	Total		38.1	42.3
Age	40-54 years old	21.5	18.4	24.7
	55 and over	52.5	49.8	55.2
Race/	Non-Hispanic White	43.0	40.7	45.3
Ethnicity	Non-Hispanic Black	32.0	23.9	40.2
	Hispanic*	24.1	16.9	31.2
	Non-Hispanic Other	34.6	24.6	44.6
Income	Less than \$35,000	30.7	26.1	35.4
	\$35,000-\$74,999	39.9	35.4	44.4
	\$75,000 and more	44.3	41.2	47.4
Insurance	Insured	41.3	39.2	43.5
	Not Insured**	20.0	-	-
Disability	Disabled	39.9	35.7	44.1
	Non-disabled	40.1	37.7	42.5
Education	High School Graduate or Less	31.6	27.9	35.3
	More than High School	45.4	42.9	47.9

Had Blood Stool Test within the past 3 years (50-75)

<u>Indicator Description: Among adults 50-75 years old, the proportion who reported had a blood stool test</u> <u>within the past 3 years.</u>

Demographics		%	95% CI	
Total	Total		13.6	16.0
Sex	Male	14.3	12.5	16.1
	Female	15.2	13.6	16.8
Race/	Non-Hispanic White	15.0	13.8	16.3
Ethnicity	Non-Hispanic Black	19.8	14.1	25.5
	Hispanic**	10.0	-	-
	Non-Hispanic Other**	15.0	-	-
Income	Less than \$35,000	14.8	12.2	17.4
	\$35,000-\$74,999	16.3	13.7	19.0
	\$75,000 and more	14.7	12.9	16.5
Insurance	Insured	15.1	13.9	16.3
	Not Insured**	10.0	ı	ı
Disability	Disabled	16.8	14.4	19.3
	Non-disabled	14.1	12.7	15.4
Education	High School Graduate or Less	14.7	12.5	17.0
	More than High School	14.9	13.5	16.2

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Had Any Permanent Teeth Extracted

Indicator Description: Among all adults, the proportion who reported that they were missing any teeth due to tooth decay or gum disease. This excludes teeth lost for other reasons, such as injury or orthodontics.

Demographics		%	95%	6 CI
Total		37.3	36.1	38.6
Age	18-34 years old	14.4	12.1	16.7
	35-54 years old	32.8	30.5	35.1
	55 and over	57.9	56.2	59.5
Sex	Male	37.3	35.4	39.2
	Female	37.4	35.6	39.1
Race/	Non-Hispanic White	36.6	35.2	38.0
Ethnicity	Non-Hispanic Black	46.8	42.2	51.3
	Hispanic	37.1	33.1	41.1
	Non-Hispanic Other	30.4	25.1	35.7
Income	Less than \$35,000	49.9	46.9	52.8
	\$35,000-\$74,999	43.2	40.3	46.1
	\$75,000 and more	27.0	25.3	28.8
Insurance	Insured	37.4	36.1	38.7
	Not Insured	38.1	32.7	43.5
Disability	Disabled	56.1	53.2	59.0
	Non-disabled	32.0	30.6	33.4
Education	High School Graduate or Less	47.9	45.4	50.3
	More than High School	30.9	29.6	32.3

Ever Had Sigmoidoscopy/ Colonoscopy (50-75)

<u>Indicator Description: Among adults 50-75 years old, the proportion who reported that they ever had sigmoidoscopy or colonoscopy.</u>

Demographics		%	95%	6 CI
Total	Total		76.6	79.6
Sex	Male	76.7	74.5	79.0
	Female	79.2	77.2	81.3
Race/	Non-Hispanic White	79.6	78.0	81.2
Ethnicity	Non-Hispanic Black	75.5	69.0	81.9
	Hispanic	70.2	63.3	77.2
	Non-Hispanic Other	71.9	63.2	80.7
Income	Less than \$35,000	70.5	66.6	74.3
	\$35,000-\$74,999	79.1	75.9	82.3
	\$75,000 and more	81.5	79.5	83.6
Insurance	Insured	78.9	77.4	80.4
	Not Insured	57.7	47.0	68.3
Disability	Disabled	79.0	75.8	82.1
	Non-disabled	77.9	76.1	79.6
Education	High School Graduate or Less	73.4	70.5	76.4
	More than High School	80.8	79.2	82.5

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Ever Tested for Hepatitis C

<u>Indicator Description:</u> Among all adults, the proportion who reported that they ever tested for hepatitis <u>C.</u>

Demographics		%	95%	6 CI
Total		28.6	26.6	30.6
Age	18-34 years old	29.9	24.9	34.9
	35-54 years old	32.8	29.2	36.4
	55 and over	25.2	22.9	27.5
Sex	Male	29.1	26.2	32.0
	Female	28.1	25.4	30.8
Race/	Non-Hispanic White	26.9	24.7	29.1
Ethnicity	Non-Hispanic Black	37.2	30.3	44.1
	Hispanic	30.6	24.3	36.9
	Non-Hispanic Other*	29.0	20.2	37.8
Income	Less than \$35,000	29.8	25.5	34.0
	\$35,000-\$74,999	26.0	22.1	29.9
	\$75,000 and more	29.9	26.9	33.0
Insurance	Insured	28.9	26.9	30.9
	Not Insured*	25.1	17.1	33.1
Disability	Disabled	34.7	30.4	39.0
	Non-disabled	26.8	24.6	29.0
Education	High School Graduate or Less	25.3	21.9	28.6
	More than High School	30.7	28.3	33.2

Had Blood Sugar Test in past three years

<u>Indicator Description:</u> Among all adults, the proportion who had a test for high blood sugar in past three <u>years.</u>

Demographics		%	95%	6 CI
Total		56.6	54.5	58.7
Age	18-34 years old	43.9	38.9	48.9
	35-54 years old	59.3	55.7	62.9
	55 and over	64.7	62.2	67.2
Sex	Male	55.9	52.7	59.1
	Female	57.2	54.4	60.1
Race/	Non-Hispanic White	56.7	54.2	59.2
Ethnicity	Non-Hispanic Black	61.5	54.6	68.4
	Hispanic	56.4	49.8	63.0
	Non-Hispanic Other	48.5	39.3	57.6
Income	Less than \$35,000	56.8	52.3	61.4
	\$35,000-\$74,999	55.2	50.5	59.9
	\$75,000 and more	58.5	55.1	61.8
Insurance	Insured	57.3	55.1	59.5
	Not Insured	47.0	38.3	55.8
Disability	Disabled	55.2	50.3	60.0
	Non-disabled	57.3	54.8	59.7
Education	High School Graduate or Less	54.8	50.9	58.7
Euucation	More than High School	57.6	55.1	60.1

Current Asthma

<u>Indicator Description:</u> Among all adults, the proportion who reported that they were ever told by a doctor, nurse, or other health care professional that they had asthma, and still had it.

Demographics		%	95% CI	
Total		10.3	9.5	11.1
Age	18-34 years old	9.1	7.3	10.8
	35-54 years old	11.8	10.4	13.3
	55 and over	10.1	9.0	11.1
Sex	Male	7.6	6.6	8.6
	Female	12.8	11.6	14.0
Race/	Non-Hispanic White	9.1	8.3	9.9
Ethnicity	Non-Hispanic Black	15.3	11.8	18.8
	Hispanic	13.0	10.3	15.7
	Non-Hispanic Other*	8.8	5.9	11.8
Income	Less than \$35,000	15.5	13.5	17.6
	\$35,000-\$74,999	8.9	7.3	10.5
	\$75,000 and more	7.5	6.5	8.5
Insurance	Insured	10.4	9.6	11.2
	Not Insured*	8.8	5.8	11.8
Disability	Disabled	18.3	16.2	20.4
	Non-disabled	7.8	7.0	8.6
Education	High School Graduate or Less	12.8	11.2	14.4
	More than High School	8.7	7.9	9.6

Chronic Obstructive Pulmonary Diseases

<u>Indicator Description: Among all adults, the proportion who reported ever told by a doctor that they had chronic obstructive pulmonary disease (COPD), emphysema or chronic bronchitis.</u>

Demographics		%	95%	% CI
Total		5.3	4.8	5.9
Age	18-34 years old**	<0.1	ı	-
	35-54 years old	3.3	2.5	4.1
	55 and over	9.7	8.7	10.7
Sex	Male	4.8	4.1	5.6
	Female	5.8	5.0	6.5
Race/	Non-Hispanic White	5.9	5.3	6.6
Ethnicity	Non-Hispanic Black*	3.6	2.3	5.0
	Hispanic*	3.8	2.4	5.3
	Non-Hispanic Other**	5.0	-	-
Income	Less than \$35,000	9.8	8.4	11.3
	\$35,000-\$74,999	5.1	3.9	6.2
	\$75,000 and more	2.9	2.3	3.6
Insurance	Insured	5.5	5.0	6.1
	Not Insured**	5.0	-	-
Disability	Disabled	14.5	12.7	16.3
	Non-disabled	2.7	2.3	3.1
Education	High School Graduate or Less	8.1	7.0	9.2
	More than High School	3.6	3.1	4.1

Arthritis

<u>Indicator Description: Among all adults, the proportion who reported ever told by a doctor that they had some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia.</u>

Demograph	Demographics		95%	6 CI
Total		26.0	24.9	27.1
Age	18-34 years old	5.6	4.1	7.1
	35-54 years old	20.5	18.5	22.4
	55 and over	44.9	43.3	46.6
Sex	Male	21.3	19.8	22.7
	Female	30.3	28.7	31.9
Race/	Non-Hispanic White	29.5	28.2	30.8
Ethnicity	Non-Hispanic Black	22.5	18.7	26.4
	Hispanic	14.8	12.1	17.5
	Non-Hispanic Other	17.6	13.0	22.2
Income	Less than \$35,000	30.9	28.4	33.4
	\$35,000-\$74,999	26.8	24.4	29.2
	\$75,000 and more	21.8	20.2	23.4
Insurance	Insured	27.0	25.9	28.2
	Not Insured	13.7	10.0	17.4
Disability	Disabled	48.2	45.4	51.0
	Non-disabled	19.5	18.4	20.6
Education	High School Graduate or Less	29.4	27.3	31.5
	More than High School	23.8	22.6	25.0

Cardiovascular Diseases

<u>Indicator Description: Among all adults, the proportion who reported ever diagnosed with a heart attack, coronary heart disease, or stroke.</u>

Demographics		%	95%	6 CI
Total		7.5	6.9	8.1
Age	18-34 years old**	<0.1	-	-
	35-54 years old	3.6	2.7	4.4
	55 and over	15.5	14.2	16.7
Sex	Male	8.8	7.8	9.8
	Female	6.4	5.6	7.1
Race/	Non-Hispanic White	8.4	7.7	9.1
Ethnicity	Non-Hispanic Black*	6.2	3.9	8.4
	Hispanic*	5.3	3.6	7.0
	Non-Hispanic Other**	5.0	-	-
Income	Less than \$35,000	11.6	9.9	13.2
	\$35,000-\$74,999	7.5	6.3	8.6
	\$75,000 and more	4.6	3.9	5.3
Insurance	Insured	7.8	7.2	8.4
	Not Insured**	5.0	-	-
Disability	Disabled	18.4	16.5	20.4
	Non-disabled	4.3	3.8	4.8
Education	High School Graduate or Less	9.8	8.5	11.1
	More than High School	6.2	5.6	6.8

Diabetes

<u>Indicator Description:</u> Among all adults, the proportion who reported that they were ever told by a <u>doctor that they had diabetes.</u>

Demographics		%	95% CI	
Total		9.7	9.0	10.4
Age	18-34 years old**	<0.1	-	-
	35-54 years old	6.6	5.5	7.7
	55 and over	18.5	17.0	19.9
Sex	Male	10.7	9.6	11.8
	Female	8.7	7.8	9.6
Race/	Non-Hispanic White	9.1	8.4	9.9
Ethnicity	Non-Hispanic Black	15.3	12.0	18.7
	Hispanic	9.1	6.9	11.2
	Non-Hispanic Other*	8.2	5.3	11.2
Income	Less than \$35,000	15.3	13.3	17.2
	\$35,000-\$74,999	9.8	8.3	11.3
	\$75,000 and more	6.0	5.2	6.8
Insurance	Insured	9.8	9.1	10.5
	Not Insured*	8.4	5.2	11.5
Disability	Disabled	20.4	18.3	22.6
	Non-disabled	6.6	5.9	7.2
Education	High School Graduate or Less	13.9	12.4	15.4
	More than High School	7.1	6.4	7.7

Pre-diabetes

<u>Indicator Description: Among all adults, the proportion who reported that they had ever diagnosed with pre-diabetes.</u>

Demographics		%	95%	6 CI
Total	Total		9.4	11.8
Age	18-34 years old*	5.6	3.4	7.7
	35-54 years old	10.4	8.2	12.7
	55 and over	14.9	13.0	16.8
Sex	Male	9.3	7.7	10.8
	Female	11.8	10.0	13.6
Race/	Non-Hispanic White	9.2	8.0	10.4
Ethnicity	Non-Hispanic Black*	13.5	8.7	18.2
	Hispanic*	15.1	10.3	19.9
	Non-Hispanic Other**	10.0	-	-
Income	Less than \$35,000	14.4	11.1	17.7
	\$35,000-\$74,999	12.6	9.6	15.6
	\$75,000 and more	8.3	6.8	9.8
Insurance	Insured	10.5	9.3	11.8
	Not Insured**	15.0	-	-
Disability	Disabled	17.0	13.5	20.5
	Non-disabled	9.0	7.7	10.2
Education	High School Graduate or Less	12.7	10.3	15.2
	More than High School	9.4	8.1	10.7

Kidney Diseases

<u>Indicator Description: Among all adults, the proportion who reported ever told by a doctor that they had kidney disease.</u>

Demograph	Demographics		95% CI	
Total		2.6	2.2	3.0
Age	18-34 years old	-	-	-
	35-54 years old**	<0.1	-	-
	55 and over	4.9	4.0	5.8
Sex	Male	2.8	2.2	3.4
	Female	2.4	1.8	2.9
Race/	Non-Hispanic White	2.5	2.0	2.9
Ethnicity	Non-Hispanic Black**	5.0	-	-
	Hispanic**	5.0	-	-
	Non-Hispanic Other	-	-	-
Income	Less than \$35,000	4.3	3.1	5.5
	\$35,000-\$74,999*	1.9	1.3	2.5
	\$75,000 and more	1.7	1.2	2.1
Insurance	Insured	2.4	2.0	2.8
	Not Insured	-	-	-
Disability	Disabled	6.3	4.9	7.8
	Non-disabled	1.5	1.1	1.8
Education	High School Graduate or Less	3.8	2.9	4.8
	More than High School	1.8	1.5	2.1

Depression

<u>Indicator Description:</u> Among all adults, the proportion who reported ever told by a doctor that they had a depressive disorder including depression, major/minor depression, or dysthymia.

Demographics		%	95%	6 CI
Total		15.5	14.5	16.5
Age	18-34 years old	19.1	16.6	21.7
	35-54 years old	15.0	13.4	16.7
	55 and over	13.8	12.6	14.9
Sex	Male	11.5	10.2	12.7
	Female	19.1	17.6	20.6
Race/	Non-Hispanic White	15.9	14.8	17.1
Ethnicity	Non-Hispanic Black	14.3	11.0	17.5
	Hispanic	15.7	12.6	18.9
	Non-Hispanic Other*	12.4	8.1	16.7
Income	Less than \$35,000	22.5	20.1	24.8
	\$35,000-\$74,999	16.7	14.5	19.0
	\$75,000 and more	11.2	9.9	12.4
Insurance	Insured	15.8	14.8	16.8
	Not Insured*	12.3	8.5	16.1
Disability	Disabled	34.3	31.4	37.1
	Non-disabled	10.1	9.2	11.0
Education	High School Graduate or Less	16.4	14.6	18.3
	More than High School	14.9	13.8	16.1

Periodontal Disease

<u>Indicator Description: Among all adults, the proportion who reported they have gum diseases.</u>

Demographics		%	95%	6 CI
Total	Total		10.6	12.8
Age	18-34 years old**	5.0	-	-
	35-54 years old	9.9	7.9	11.8
	55 and over	18.3	16.5	20.2
Sex	Male	11.4	9.8	13.0
	Female	11.9	10.3	13.4
Race/	Non-Hispanic White	13.3	11.9	14.6
Ethnicity	Non-Hispanic Black*	12.1	7.7	16.6
	Hispanic**	5.0	-	-
	Non-Hispanic Other**	10.0	-	-
Income	Less than \$35,000	13.1	10.4	15.8
	\$35,000-\$74,999	15.6	12.7	18.5
	\$75,000 and more	10.1	8.6	11.6
Insurance	Insured	12.4	11.2	13.6
	Not Insured**	5.0	-	-
Disability	Disabled	16.2	13.7	18.8
	Non-disabled	10.2	9.0	11.4
Education	High School Graduate or Less	12.3	10.2	14.4
	More than High School	11.2	10.0	12.5

Ever Had Treatment for Periodontal Disease

<u>Indicator Description:</u> Among all adults, the proportion who reported that they ever had treatment <u>for gum disease such as scaling and root planing, or deep cleaning.</u>

Demographics		%	95%	6 CI
Total	Total		16.1	18.8
Age	18-34 years old	8.6	6.3	11.0
	35-54 years old	13.8	11.6	16.0
	55 and over	26.6	24.4	28.8
Sex	Male	17.7	15.7	19.7
	Female	17.2	15.4	19.0
Race/	Non-Hispanic White	17.6	16.1	19.1
Ethnicity	Non-Hispanic Black	22.4	17.3	27.6
	Hispanic	15.7	11.7	19.7
	Non-Hispanic Other**	10.0	-	-
Income	Less than \$35,000	18.5	15.3	21.6
	\$35,000-\$74,999	22.1	18.9	25.3
	\$75,000 and more	14.5	12.8	16.3
Insurance	Insured	17.9	16.5	19.3
	Not Insured*	12.7	7.9	17.5
Disability	Disabled	22.9	19.8	26.0
	Non-disabled	15.9	14.4	17.3
Education	High School Graduate or Less	19.1	16.6	21.7
	More than High School	16.3	14.9	17.8

Suicide Ideation

<u>Indicator Description:</u> Among all adults, the proportion who reported they had ever thought of taking their own life.

Demographics		%	95%	6 CI
Total		12.4	10.9	14.0
Age	18-34 years old	22.0	17.4	26.7
	35-54 years old	11.8	9.4	14.3
	55 and over	7.2	5.9	8.6
Sex	Male	13.5	11.1	15.9
	Female	11.4	9.4	13.5
Race/	Non-Hispanic White	12.2	10.4	14.1
Ethnicity	Non-Hispanic Black**	10.0	-	-
	Hispanic*	13.5	9.3	17.7
	Non-Hispanic Other**	15.0	-	-
Income	Less than \$35,000	12.9	9.8	16.0
	\$35,000-\$74,999	15.5	12.1	19.0
	\$75,000 and more	9.3	7.4	11.2
Insurance	Insured	12.5	10.9	14.2
	Not Insured**	10.0	-	-
Disability	Disabled	23.1	18.8	27.5
	Non-disabled	9.1	7.6	10.6
Education	High School Graduate or Less	13.5	10.3	16.6
	More than High School	11.9	10.2	13.6

Suicide Attempt

<u>Indicator Description: Among adults who thought of suicide, the proportion who attempted suicide.</u>

Demographics		%	95%	6 CI
Total	Total		23.8	37.3
Age	18-34 years old*	31.9	19.7	44.2
	35-54 years old*	33.7	23.5	43.9
	55 and over*	23.4	15.0	31.9
Sex	Male*	29.9	19.6	40.3
	Female	31.3	22.6	40.0
Race/	Non-Hispanic White	29.8	21.4	38.3
Ethnicity	Non-Hispanic Black	-	-	-
	Hispanic**	40.0	-	-
	Non-Hispanic Other	-	-	-
Income	Less than \$35,000	44.3	31.6	56.9
	\$35,000-\$74,999**	20.0	-	-
	\$75,000 and more**	25.0	-	-
Insurance	Insured	30.9	23.8	38.0
	Not Insured	-	-	-
Disability	Disabled	42.5	30.7	54.3
	Non-disabled*	21.1	14.5	27.7
Education	High School Graduate or Less*	38.4	25.1	51.7
	More than High School	25.4	18.8	32.0

Suicide Campaign

Indicator Description: Among all adults, the proportion who reported they were aware of the "1 WORD, 1 VOICE, 1 LIFE" suicide prevention campaign in CT.

Demographics		%	95%	6 CI
Total		35.4	33.4	37.4
Age	18-34 years old	40.8	35.6	46.1
	35-54 years old	36.5	33.1	39.9
	55 and over	31.2	28.9	33.5
Sex	Male	30.3	27.4	33.3
	Female	40.2	37.5	42.9
Race/	Non-Hispanic White	36.1	33.8	38.4
Ethnicity	Non-Hispanic Black	41.1	34.0	48.3
	Hispanic	30.3	24.4	36.2
	Non-Hispanic Other*	32.3	22.0	42.6
Income	Less than \$35,000	36.8	32.5	41.2
	\$35,000-\$74,999	39.8	35.4	44.2
	\$75,000 and more	32.6	29.7	35.5
Insurance	Insured	36.0	33.9	38.0
	Not Insured	28.5	20.2	36.8
Disability	Disabled	38.3	34.0	42.6
	Non-disabled	34.5	32.3	36.8
Education	High School Graduate or Less	36.2	32.4	40.0
	More than High School	34.9	32.7	37.1

Public Water

<u>Indicator Description: Among all adults, the proportion who reported having public water in their home.</u>

Demographics		%	95%	6 CI
Total	Total		68.0	71.6
Age	18-34 years old	72.3	67.4	77.1
	35-54 years old	69.3	66.2	72.4
	55 and over	68.5	66.3	70.8
Sex	Male	69.7	67.0	72.5
	Female	69.8	67.4	72.2
Race/	Non-Hispanic White	64.8	62.7	66.9
Ethnicity	Non-Hispanic Black	92.5	88.7	96.3
	Hispanic	77.7	71.7	83.7
	Non-Hispanic Other	82.6	76.1	89.2
Income	Less than \$35,000	80.6	77.1	84.1
	\$35,000-\$74,999	72.3	68.5	76.0
	\$75,000 and more	63.8	60.8	66.7
Insurance	Insured	69.9	68.1	71.7
	Not Insured	65.8	56.5	75.2
Disability	Disabled	72.7	69.1	76.3
	Non-disabled	68.9	66.8	71.0
Education	High School Graduate or Less	72.0	68.5	75.4
	More than High School	68.6	66.5	70.6

Well Water

<u>Indicator Description: Among all adults, the proportion who reported having private well water in their home.</u>

Demographics		%	95%	6 CI
Total		26.7	25.0	28.4
Age	18-34 years old	21.2	16.8	25.6
	35-54 years old	27.9	24.8	30.9
	55 and over	29.1	26.9	31.2
Sex	Male	26.6	24.0	29.1
	Female	26.8	24.5	29.1
Race/	Non-Hispanic White	32.5	30.4	34.5
Ethnicity	Non-Hispanic Black	-	-	-
	Hispanic*	12.4	7.6	17.2
	Non-Hispanic Other**	15.0	-	-
Income	Less than \$35,000	13.5	10.6	16.4
	\$35,000-\$74,999	24.4	20.9	27.9
	\$75,000 and more	33.8	30.9	36.6
Insurance	Insured	27.2	25.5	29.0
	Not Insured**	20.0	-	-
Disability	Disabled	22.1	18.8	25.3
	Non-disabled	28.1	26.1	30.1
Education	High School Graduate or Less	22.9	19.7	26.1
	More than High School	28.8	26.8	30.8

Prescription Drug Safe Storage

Indicator Description: Among all adults, the proportion who reported their opioid-based pain relievers in their home are safely stored away from others who may use them for recreational or other non-medical purposes.

Demographics		%	95%	6 CI
Total	Total		42.5	46.8
Age	18-34 years old	50.0	44.5	55.4
	35-54 years old	44.0	40.2	47.9
	55 and over	41.5	39.0	44.1
Sex	Male	44.5	41.3	47.8
	Female	44.8	41.8	47.7
Race/	Non-Hispanic White	41.1	38.7	43.6
Ethnicity	Non-Hispanic Black	55.1	47.7	62.4
	Hispanic	57.8	50.9	64.6
	Non-Hispanic Other	45.7	35.7	55.7
Income	Less than \$35,000	54.6	50.1	59.2
	\$35,000-\$74,999	45.9	41.3	50.5
	\$75,000 and more	38.4	35.0	41.8
Insurance	Insured	43.9	41.7	46.1
	Not Insured	56.6	46.8	66.3
Disability	Disabled	50.4	46.0	54.7
	Non-disabled	42.8	40.3	45.3
Education	High School Graduate or Less	52.3	48.5	56.1
	More than High School	40.0	37.5	42.6

Prescription Drug Safe Disposal

<u>Indicator Description:</u> Among all adults, the proportion who reported they are aware of how to safely dispose of their prescription opioid-based pain relievers when they are finished.

Demographics		%	95%	6 CI
Total	Total		76.4	80.0
Age	18-34 years old	75.0	70.4	79.6
	35-54 years old	79.1	75.9	82.3
	55 and over	79.5	77.5	81.5
Sex	Male	75.2	72.4	78.0
	Female	81.0	78.7	83.3
Race/	Non-Hispanic White	80.0	78.1	81.9
Ethnicity	Non-Hispanic Black	77.0	71.0	83.1
	Hispanic	73.4	67.2	79.7
	Non-Hispanic Other*	69.9	59.5	80.3
Income	Less than \$35,000	79.1	75.1	83.1
	\$35,000-\$74,999	77.2	73.1	81.3
	\$75,000 and more	77.7	75.0	80.4
Insurance	Insured	78.4	76.5	80.2
	Not Insured*	77.7	70.0	85.3
Disability	Disabled	80.6	77.2	83.9
	Non-disabled	77.4	75.3	79.5
Education	High School Graduate or Less	80.3	77.2	83.4
	More than High School	77.0	74.9	79.2

Firearm

<u>Indicator Description:</u> Among all adults, the proportion who reported having at least one firearm kept in <u>or around home.</u>

Demographics		%	95%	6 CI
Total	Total		16.3	19.5
Age	18-34 years old	18.8	14.7	22.9
	35-54 years old	17.2	14.6	19.8
	55 and over	18.3	16.4	20.3
Sex	Male	23.8	21.1	26.4
	Female	12.7	11.0	14.4
Race/	Non-Hispanic White	22.0	20.0	24.0
Ethnicity	Non-Hispanic Black**	10.0	-	-
	Hispanic**	10.0	-	-
	Non-Hispanic Other**	10.0	-	-
Income	Less than \$35,000	10.5	7.8	13.2
	\$35,000-\$74,999	16.8	13.2	20.5
	\$75,000 and more	25.9	23.1	28.6
Insurance	Insured	18.6	17.0	20.3
	Not Insured	-	-	-
Disability	Disabled	16.4	13.4	19.4
	Non-disabled	18.4	16.6	20.3
Education	High School Graduate or Less	14.6	11.9	17.3
	More than High School	19.8	17.9	21.7

Air Quality Index Awareness

<u>Indicator Description:</u> Among all adults, the proportion who reported ever heard or read about the air <u>quality index or air quality alerts where they live.</u>

Demographics		%	95%	6 CI
Total		44.8	42.8	46.8
Age	18-34 years old	33.0	28.3	37.8
	35-54 years old	43.3	39.8	46.7
	55 and over	52.4	49.9	55.0
Sex	Male	48.2	45.1	51.2
	Female	41.6	39.0	44.2
Race/	Non-Hispanic White	50.4	48.1	52.7
Ethnicity	Non-Hispanic Black	31.3	24.9	37.6
	Hispanic	26.5	21.0	31.9
	Non-Hispanic Other	37.1	27.1	47.1
Income	Less than \$35,000	34.1	30.0	38.1
	\$35,000-\$74,999	44.5	40.2	48.8
	\$75,000 and more	53.3	50.2	56.3
Insurance	Insured	46.5	44.4	48.5
	Not Insured*	20.7	14.3	27.1
Disability	Disabled	41.2	37.1	45.2
	Non-disabled	46.1	43.8	48.3
Education	High School Graduate or Less	30.9	27.7	34.2
	More than High School	52.8	50.5	55.2

Sexual Violence

<u>Indicator Description: Among all adults, the proportion who reported ever been forced to have sexual intercourse since 18 years old.</u>

Demograph	Demographics		95% CI	
Total		8.7	7.4	10.0
Age	18-34 years old*	12.1	8.4	15.7
	35-54 years old	10.9	8.7	13.2
	55 and over	4.9	4.1	5.8
Sex	Male*	2.8	1.7	3.9
	Female	14.2	12.0	16.4
Race/	Non-Hispanic White	8.7	7.3	10.2
Ethnicity	Non-Hispanic Black**	10.0	-	-
	Hispanic**	10.0	-	-
	Non-Hispanic Other	-	-	-
Income	Less than \$35,000*	9.1	6.4	11.8
	\$35,000-\$74,999	9.1	6.6	11.6
	\$75,000 and more	8.0	6.1	9.9
Insurance	Insured	8.7	7.4	10.0
	Not Insured	-	-	-
Disability	Disabled	15.6	11.7	19.4
	Non-disabled	6.7	5.5	7.8
Education	High School Graduate or Less*	6.5	4.3	8.7
	More than High School	9.9	8.3	11.5

Overweight (5-17 years old)

<u>Indicator Description: Among children 5-17 years old, the proportion of children whose BMI was</u> between the 85th and 95th percentile for children of the same age and sex.

Demographics		%	95% CI	
Total		16.2	13.2	19.3
Age	5-11 years old	16.1	11.5	20.8
	12-17 years old	16.3	12.2	20.4
Sex	Male	17.2	13.0	21.4
	Female*	15.1	10.6	19.7
Race/	Non-Hispanic White	15.9	12.2	19.8
Ethnicity	Non-Hispanic Black**	25.0	ı	ı
	Hispanic**	15.0	ı	ı
	Non-Hispanic Other	ı	ı	ı
Household	Less than \$35,000**	20.0	ı	ı
Income	\$35,000-\$74,999*	21.7	13.3	30.1
	\$75,000 and more	13.3	9.6	16.9
Caregiver's	Insured	15.3	12.2	18.4
Insurance	Not Insured**	35.0	-	-
Caregiver's	High School Graduate or Less*	22.3	14.2	30.3
Education	More than High School	14.9	11.6	18.1

Obese (5-17 years old)

<u>Indicator Description: Among children 5-17 years old, the proportion of children whose BMI was above</u> the 95th percentile for children of the same age and sex.

Demographics		%	95% CI	
Total	tal 1		15.0	21.7
Age	5-11 years old	27.7	21.6	33.7
	12-17 years old*	10.1	6.9	13.3
Sex	Male	18.2	13.5	22.8
	Female	18.6	13.5	23.6
Race/	Non-Hispanic White	12.8	9.5	16.1
Ethnicity	Non-Hispanic Black**	15.0	ı	ı
	Hispanic*	31.4	21.8	41.0
	Non-Hispanic Other**	15.0	-	-
Household	Less than \$35,000*	24.8	17.0	32.6
Income	\$35,000-\$74,999**	20.0	-	-
	\$75,000 and more	15.2	10.9	19.6
Caregiver's	Insured	18.2	14.7	21.6
Insurance	Not Insured	-	-	-
Caregiver's	High School Graduate or Less*	20.3	14.1	26.5
Education	More than High School	17.9	14.0	21.8

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Ever Breastfed

<u>Indicator Description: Among all children, the proportion of children ever breastfed or given pumped</u> breast milk, even for a short period of time.

Demographics		%	95% CI	
Total		81.6	79.3	83.9
Age	0-4 years old*	84.0	78.7	89.4
	5-11 years old	83.8	79.9	87.7
	12-17 years old	78.6	74.9	82.3
Sex	Male	80.3	77.0	83.6
	Female	82.7	79.4	86.0
Race/	Non-Hispanic White	84.5	81.8	87.1
Ethnicity	Non-Hispanic Black*	75.3	67.5	83.2
	Hispanic	79.9	74.8	85.0
	Non-Hispanic Other*	79.6	72.0	87.2
Household	Less than \$35,000	69.7	63.3	76.2
Income	\$35,000-\$74,999	78.0	71.8	84.1
	\$75,000 and more	87.3	84.7	89.9
Caregiver's	Insured	81.8	79.4	84.2
Insurance	Not Insured**	80.0	ı	-
Caregiver's	High School Graduate or Less	70.4	64.8	75.9
Education	More than High School	84.9	82.5	87.4

Excessive Screen Time

<u>Indicator Description: Among children aged 2-17 years old, the proportion of children who had screen time more than 2 hours daily.</u>

Demographi	emographics % 95% CI		6 CI	
Total		56.5	53.1	59.9
Age	2-4 years old	46.5	37.5	55.5
	5-11 years old	51.3	45.7	56.8
	12-17 years old	65.9	61.2	70.6
Sex	Male	59.6	54.9	64.2
	Female	53.3	48.3	58.2
Race/	Non-Hispanic White	52.8	48.6	57.0
Ethnicity	Non-Hispanic Black*	66.7	56.2	77.3
	Hispanic	58.4	51.1	65.7
	Non-Hispanic Other	59.3	47.7	71.0
Household	Less than \$35,000	59.0	51.2	66.8
Income	\$35,000-\$74,999	62.0	54.0	70.0
	\$75,000 and more	54.3	49.6	59.0
Caregiver's	Insured	57.0	53.4	60.5
Insurance	Not Insured	53.5	41.6	65.4
Caregiver's	High School Graduate or Less	60.0	53.0	67.0
Education	More than High School	55.5	51.6	59.4

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Soda Consumption

<u>Indicator Description: Among children 2-17 years old, the proportion of children who drank soda or sugary drinks at least once daily.</u>

Demographics		%	95% CI	
Total		28.3	25.1	31.5
Age	2-4 years old*	26.9	18.2	35.6
	5-11 years old	25.5	20.2	30.8
	12-17 years old	31.7	27.2	36.2
Sex	Male	29.8	25.4	34.2
	Female	26.6	21.9	31.3
Race/	Non-Hispanic White	20.5	17.2	23.8
Ethnicity	Non-Hispanic Black	40.7	30.2	51.2
	Hispanic	38.1	30.6	45.6
	Non-Hispanic Other**	25.0	-	-
Household	Less than \$35,000	44.4	36.4	52.4
Income	\$35,000-\$74,999	30.6	22.9	38.2
	\$75,000 and more	21.8	17.9	25.8
Caregiver's	Insured	28.3	24.9	31.7
Insurance	Not Insured*	30.4	19.3	41.6
Caregiver's	High School Graduate or Less	38.5	31.1	45.8
Education	More than High School	25.4	21.9	29.0

Fast Food Consumption

<u>Indicator Description: Among children 2-17 years old, the proportion of children who ate fast food at</u> least twice weekly.

- Weekiy.					
Demographics		%	95% CI		
Total		41.3	37.9	44.7	
Age	2-4 years old	41.5	32.4	50.5	
	5-11 years old	42.6	36.8	48.3	
	12-17 years old	39.9	35.4	44.4	
Sex	Male	44.2	39.4	49.0	
	Female	38.2	33.4	43.0	
Race/	Non-Hispanic White	37.2	33.1	41.2	
Ethnicity	Non-Hispanic Black	54.3	44.0	64.5	
	Hispanic	42.9	35.2	50.6	
	Non-Hispanic Other	43.6	31.6	55.6	
Household	Less than \$35,000	42.2	34.3	50.0	
Income	\$35,000-\$74,999	43.6	35.2	52.0	
	\$75,000 and more	40.0	35.5	44.6	
Caregiver's	Insured	41.9	38.3	45.4	
Insurance	Not Insured*	34.4	22.5	46.4	
Caregiver's	High School Graduate or Less	44.9	37.6	52.2	
Education	More than High School	40.2	36.3	44.1	

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Dentist Visit in the past year

<u>Indicator Description: Among all children, the proportion of children who visited a dentist in the past year.</u>

Demographi	Demographics		95% CI	
Total		86.9	84.8	89.0
Age	0-4 years old	61.6	54.7	68.5
	5-11 years old**	95.0	-	-
	12-17 years old**	95.0	ı	-
Sex	Male	86.4	83.3	89.5
	Female	87.3	84.2	90.3
Race/	Non-Hispanic White	89.2	86.7	91.7
Ethnicity	Non-Hispanic Black**	90.0	-	-
	Hispanic	80.2	74.4	86.0
	Non-Hispanic Other**	90.0	-	-
Household	Less than \$35,000*	82.0	76.4	87.6
Income	\$35,000-\$74,999**	85.0	-	-
	\$75,000 and more	89.3	86.7	91.9
Caregiver's	Insured	87.4	85.2	89.6
Insurance	Not Insured**	80.0	-	-
Caregiver's	High School Graduate or Less	81.8	76.9	86.8
Education	More than High School	88.3	86.0	90.7

Dental Sealants

<u>Indicator Description: Among children 5-17 years old, the proportion of children who ever received</u> dental seala<u>nt on their permanent teeth.</u>

Demographics		%	95% CI	
Total		47.2	43.4	51.0
Age	5-11 years old	37.3	31.7	42.9
	12-17 years old	58.0	53.2	62.9
Sex	Male	46.9	41.6	52.1
	Female	47.6	42.1	53.2
Race/	Non-Hispanic White	52.2	47.7	56.8
Ethnicity	Non-Hispanic Black*	40.4	28.5	52.3
	Hispanic	44.3	35.5	53.1
	Non-Hispanic Other*	40.6	28.6	52.6
Household	Less than \$35,000	36.2	27.7	44.8
Income	\$35,000-\$74,999	48.6	39.2	58.0
	\$75,000 and more	51.9	46.9	56.9
Caregiver's	Insured	48.8	44.8	52.7
Insurance	Not Insured**	30.0	ı	-
Caregiver's	High School Graduate or Less	40.0	32.3	47.7
Education	More than High School	49.2	44.9	53.5

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).

Dental Decay in the past year

<u>Indicator Description: Among all children, the proportion of children told they had dental decay in the past 12 months.</u>

Demographics		%	95% CI	
Total		14.3	12.0	16.7
Age	0-4 years old	-	1	-
	5-11 years old	17.5	12.9	22.2
	12-17 years old	14.9	11.4	18.4
Sex	Male	15.3	12.1	18.6
	Female	13.6	10.1	17.2
Race/	Non-Hispanic White	10.8	8.5	13.1
Ethnicity	Non-Hispanic Black**	10.0	ı	-
	Hispanic*	18.4	12.6	24.3
	Non-Hispanic Other**	25.0	-	-
Household	Less than \$35,000*	20.7	14.4	26.9
Income	\$35,000-\$74,999*	18.9	13.0	24.9
	\$75,000 and more	10.9	7.8	14.1
Caregiver's	Insured	14.2	11.7	16.7
Insurance	Not Insured**	20.0	-	-
Caregiver's	High School Graduate or Less	20.3	14.9	25.7
Education	More than High School	12.7	10.0	15.3

Current Asthma

<u>Indicator Description: Among all children, the proportion of children ever diagnosed with asthma and still have it.</u>

Demographics		%	95% CI	
Total		9.7	8.0	11.4
Age	0-4 years old	-	-	-
	5-11 years old*	9.4	6.6	12.3
	12-17 years old	12.5	9.3	15.7
Sex	Male	10.3	8.0	12.7
	Female	9.0	6.6	11.5
Race/	Non-Hispanic White	7.8	6.1	9.6
Ethnicity	Non-Hispanic Black**	10.0	-	-
	Hispanic*	12.0	7.9	16.0
	Non-Hispanic Other**	10.0	-	-
Household	Less than \$35,000*	14.1	9.8	18.3
Income	\$35,000-\$74,999**	10.0	-	-
	\$75,000 and more	8.2	6.0	10.4
Caregiver's	Insured	8.8	7.2	10.5
Insurance	Not Insured**	20.0	-	-
Caregiver's	High School Graduate or Less*	10.3	7.1	13.4
Education	More than High School	9.6	7.6	11.6

^{&#}x27;*' estimates have a CV between 15% and 20%;

^{&#}x27;**' estimates have a CV between 20.1% and 30%;

^{&#}x27;-' estimates were suppressed because of limited validity (CV>30%).