

Prevalence of Tobacco Use Among Connecticut High School Students, 2017

Demographic Characteristics	Current Tobacco Use*		Current Cigarette Smoking		Current E-Cigarette Use		Current Cigar Smoking		Current Other Tobacco Use [§]	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Overall	17.9	(14.1-21.6)	3.5	(2.4-4.7)	14.7	(11.0-18.4)	5.3	(3.4-7.2)	4.3	(3.0-5.6)
Sex										
Male	17.8	(12.5-23.1)	4.2	(2.5-6.0)	14.8	(9.9-19.8)	7.2	(3.6-10.7)	5.0	(2.8-7.2)
Female	17.3	(14.0-20.7)	2.7	(1.4-4.0)	14.0	(10.2-17.9)	2.8	(1.6-4.1)	3.2	(1.6-4.7)
Grade										
9	11.7	(8.4-15.0)	**	—	10.2	(6.9-13.5)	**	—	**	—
10	15.2	(11.9-18.4)	3.2	(1.7-4.6)	12.0	(8.8-15.3)	3.2	(1.7-4.6)	4.2	(2.5-5.9)
11	20.9	(15.2-26.7)	4.9	(2.6-7.2)	16.8	(10.5-23.0)	8.9	(6.0-11.8)	6.4	(3.7-9.1)
12	24.4	(16.3-32.5)	5.5	(2.2-8.8)	20.4	(13.4-27.4)	8.2	(2.7-13.8)	5.0	(2.0-8.0)
Race/Ethnicity										
Non-Hispanic White	21.5	(17.2-25.7)	4.5	(2.9-6.1)	18.7	(14.3-23.0)	6.2	(3.6-8.9)	4.2	(2.5-5.8)
Non-Hispanic Black	8.9	(3.9-13.9)	**	—	**	—	**	—	**	—
Hispanic	14.1	(10.5-17.6)	**	—	9.3	(6.6-12.0)	3.4	(1.6-5.3)	4.7	(2.8-6.6)

Data Source: 2017 Connecticut Youth Tobacco Survey

Note: Current use is defined as having used a product on 1 or more of the past 30 days.

*'Tobacco' includes cigarettes, cigars, chewing tobacco, snuff, dip, pipes (other than water pipes), hookahs (water pipes) and e-cigarettes.

**Estimates based on unweighted cases fewer than 50 or with a relative standard error greater than 30% are not presented.

[§]'Other Tobacco' includes chewing tobacco, snuff, dip, pipes and hookahs.