

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Causes of Death for the Age-Specific Mortality Tables

| | Page | | Page |
|---|------|--|------|
| All causes | 2 | Hypertension with or without renal disease | 33 |
| Infectious diseases and parasitic disease | 3 | Hypertension-related | 34 |
| Septicemia | 4 | Cerebrovascular disease | 35 |
| Septicemia-related | 5 | Atherosclerosis | 36 |
| HIV infection | 6 | Atherosclerosis-related | 37 |
| All cancer | 7 | Aortic aneurysm | 38 |
| Oral cavity or pharynx | 8 | Pneumonia & influenza | 39 |
| Colorectal cancer | 9 | Chronic obstructive pulmonary disease | 40 |
| Colon cancer | 10 | COPD-related | 41 |
| Rectal cancer | 11 | Asthma | 42 |
| Pancreatic cancer | 12 | Chronic liver disease and cirrhosis | 43 |
| Lung and other respiratory cancer | 13 | Nephritis, nephrotic syndrome/nephrosis | 44 |
| Lung cancer | 14 | Nephritis-related | 45 |
| Malignant melanoma of skin | 15 | End stage renal disease | 46 |
| Female breast cancer | 16 | Unintentional injuries | 47 |
| Cervical cancer | 17 | Motor vehicle accidents | 48 |
| Endometrial cancer | 18 | Falls and fall related injuries | 49 |
| Ovarian cancer | 19 | Drowning | 50 |
| Prostate cancer | 20 | Residential fires | 51 |
| Bladder cancer | 21 | Suicide | 52 |
| Brain & other CNS cancer | 22 | Homicide and legal intervention | 53 |
| Leukemia | 23 | Homicide | 54 |
| Diabetes mellitus | 24 | All injury | 55 |
| Diabetes-related | 25 | All motor vehicle accidents | 56 |
| Diabetic ketoacidosis | 26 | All falls | 57 |
| Alzheimer's diseases | 27 | All firearms | 58 |
| Cardiovascular disease | 28 | Suicide | 59 |
| Diseases of the heart | 29 | Homicide | 60 |
| Heart disease-related | 30 | All poisoning | 61 |
| Coronary heart disease | 31 | Alcohol-induced | 62 |
| Congestive heart failure | 32 | Drug-induced | 63 |

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All causes-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 55 | 112.4 | 15.16 | 36 | 143.6 | 23.94 | 19 | 79.6 | 18.26 |
| 5-9 | 4 | 9.5 | 4.75 | 2 | 9.4 | 6.61 | 2 | 9.7 | 6.84 |
| 10-14 | 6 | 15.1 | 6.18 | 3 | 14.9 | 8.62 | 3 | 15.4 | 8.87 |
| 15-19 | 27 | 69.6 | 13.40 | 20 | 100.7 | 22.53 | 7 | 37.0 | 13.99 |
| 20-24 | 40 | 88.9 | 14.06 | 32 | 142.8 | 25.25 | 8 | 35.4 | 12.52 |
| 25-29 | 65 | 144.7 | 17.94 | 51 | 232.6 | 32.58 | 14 | 60.9 | 16.26 |
| 30-34 | 73 | 192.3 | 22.50 | 54 | 293.0 | 39.87 | 19 | 97.3 | 22.31 |
| 35-39 | 70 | 235.4 | 28.14 | 52 | 363.2 | 50.37 | 18 | 116.8 | 27.52 |
| 40-44 | 55 | 224.6 | 30.29 | 41 | 352.7 | 55.09 | 14 | 108.8 | 29.09 |
| 45-49 | 71 | 391.3 | 46.44 | 53 | 620.5 | 85.24 | 18 | 187.4 | 44.18 |
| 50-54 | 66 | 485.7 | 59.79 | 45 | 709.3 | 105.74 | 21 | 289.9 | 63.26 |
| 55-59 | 65 | 615.7 | 76.37 | 34 | 690.5 | 118.42 | 31 | 550.3 | 98.84 |
| 60-64 | 63 | 779.4 | 98.20 | 38 | 1015.8 | 164.78 | 25 | 575.8 | 115.15 |
| 65-69 | 89 | 1489.8 | 157.92 | 50 | 1947.8 | 275.46 | 39 | 1144.7 | 183.30 |
| 70-74 | 64 | 1425.7 | 178.21 | 30 | 1638.4 | 299.14 | 34 | 1279.2 | 219.37 |
| 75-79 | 68 | 2043.9 | 247.86 | 35 | 2734.4 | 462.19 | 33 | 1612.1 | 280.63 |
| 80-84 | 76 | 4189.6 | 480.58 | 36 | 5598.8 | 933.13 | 40 | 3415.9 | 540.10 |
| 85+ | 93 | 6758.7 | 700.85 | 29 | 6223.2 | 1155.61 | 64 | 7033.0 | 879.12 |
| Total deaths | 1,050 | | | 641 | | | 409 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 250.7 | - | | 312.0 | - | | 191.6 | - |
| . AAMR (1940 Std. mil.) | | 328.6 | 10.69 | | 442.2 | 18.65 | | 232.1 | 12.13 |
| . AAMR (1970 Std. mil.) | | 410.7 | 13.93 | | 532.5 | 24.44 | | 311.1 | 16.20 |
| . AAMR (2000 Std. mil.) | | 530.7 | 19.23 | | 673.6 | 33.87 | | 415.6 | 22.61 |
| . HP 2000 (1940 Std. mil.) | | 328.4 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 527.3 | - | | 688.4 | - | | 396.7 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Infectious diseases and parasitic disease (1-139)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 2 | 4.1 | 2.89 | 2 | 8.0 | 5.64 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 2 | 5.2 | 3.65 | 1 | 5.0 | 5.04 | 1 | 5.3 | 5.29 |
| 20-24 | 5 | 11.1 | 4.97 | 4 | 17.9 | 8.93 | 1 | 4.4 | 4.43 |
| 25-29 | 17 | 37.8 | 9.18 | 11 | 50.2 | 15.13 | 6 | 26.1 | 10.65 |
| 30-34 | 23 | 60.6 | 12.63 | 19 | 103.1 | 23.65 | 4 | 20.5 | 10.24 |
| 35-39 | 20 | 67.3 | 15.04 | 17 | 118.7 | 28.80 | 3 | 19.5 | 11.24 |
| 40-44 | 11 | 44.9 | 13.54 | 8 | 68.8 | 24.33 | 3 | 23.3 | 13.47 |
| 45-49 | 7 | 38.6 | 14.58 | 6 | 70.2 | 28.68 | 1 | 10.4 | 10.41 |
| 50-54 | 7 | 51.5 | 19.47 | 3 | 47.3 | 27.30 | 4 | 55.2 | 27.61 |
| 55-59 | 3 | 28.4 | 16.41 | 2 | 40.6 | 28.72 | 1 | 17.8 | 17.75 |
| 60-64 | 1 | 12.4 | 12.37 | 1 | 26.7 | 26.73 | 0 | 0.0 | 0.00 |
| 65-69 | 3 | 50.2 | 28.99 | 2 | 77.9 | 55.09 | 1 | 29.4 | 29.35 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 2 | 60.1 | 42.51 | 1 | 78.1 | 78.13 | 1 | 48.9 | 48.85 |
| 80-84 | 2 | 110.3 | 77.96 | 1 | 155.5 | 155.52 | 1 | 85.4 | 85.40 |
| 85+ | 3 | 218.0 | 125.88 | 1 | 214.6 | 214.59 | 2 | 219.8 | 155.41 |
| Total deaths | 108 | | | 79 | | | 29 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 25.8 | - | | 38.5 | - | | 13.6 | - |
| . AAMR (1940 Std. mil.) | | 27.9 | 2.81 | | 42.5 | 5.06 | | 14.6 | 2.85 |
| . AAMR (1970 Std. mil.) | | 26.7 | 2.94 | | 39.7 | 5.30 | | 15.2 | 3.12 |
| . AAMR (2000 Std. mil.) | | 33.8 | 3.95 | | 49.8 | 7.13 | | 19.6 | 4.30 |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 16.4 | - | | 25.2 | - | | 8.0 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): CT rate is higher
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Septicemia (038)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 1 | 2.2 | 2.22 | 1 | 4.5 | 4.46 | 0 | 0.0 | 0.00 |
| 25-29 | 2 | 4.5 | 3.15 | 2 | 9.1 | 6.45 | 0 | 0.0 | 0.00 |
| 30-34 | 2 | 5.3 | 3.72 | 1 | 5.4 | 5.43 | 1 | 5.1 | 5.12 |
| 35-39 | 1 | 3.4 | 3.36 | 1 | 7.0 | 6.98 | 0 | 0.0 | 0.00 |
| 40-44 | 3 | 12.3 | 7.07 | 2 | 17.2 | 12.17 | 1 | 7.8 | 7.77 |
| 45-49 | 1 | 5.5 | 5.51 | 0 | 0.0 | 0.00 | 1 | 10.4 | 10.41 |
| 50-54 | 2 | 14.7 | 10.41 | 1 | 15.8 | 15.76 | 1 | 13.8 | 13.80 |
| 55-59 | 1 | 9.5 | 9.47 | 1 | 20.3 | 20.31 | 0 | 0.0 | 0.00 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 65-69 | 1 | 16.7 | 16.74 | 1 | 39.0 | 38.96 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 2 | 60.1 | 42.51 | 1 | 78.1 | 78.13 | 1 | 48.9 | 48.85 |
| 80-84 | 2 | 110.3 | 77.96 | 1 | 155.5 | 155.52 | 1 | 85.4 | 85.40 |
| 85+ | 2 | 145.3 | 102.78 | 1 | 214.6 | 214.59 | 1 | 109.9 | 109.89 |
| Total deaths | 20 | | | 13 | | | 7 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 4.8 | - | | 6.3 | - | | 3.3 | - |
| . AAMR (1940 Std. mil.) | | 5.8 | 1.36 | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | 7.3 | 1.82 | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | 10.3 | 2.73 | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 4.0 | - | | 4.6 | - | | 3.5 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Septicemia-related (038)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 1 | 2.2 | 2.22 | 1 | 4.5 | 4.46 | 0 | 0.0 | 0.00 |
| 25-29 | 2 | 4.5 | 3.15 | 2 | 9.1 | 6.45 | 0 | 0.0 | 0.00 |
| 30-34 | 4 | 10.5 | 5.27 | 2 | 10.9 | 7.67 | 2 | 10.2 | 7.24 |
| 35-39 | 6 | 20.2 | 8.24 | 4 | 27.9 | 13.97 | 2 | 13.0 | 9.17 |
| 40-44 | 4 | 16.3 | 8.17 | 2 | 17.2 | 12.17 | 2 | 15.5 | 10.99 |
| 45-49 | 3 | 16.5 | 9.55 | 2 | 23.4 | 16.56 | 1 | 10.4 | 10.41 |
| 50-54 | 4 | 29.4 | 14.72 | 3 | 47.3 | 27.30 | 1 | 13.8 | 13.80 |
| 55-59 | 2 | 18.9 | 13.40 | 2 | 40.6 | 28.72 | 0 | 0.0 | 0.00 |
| 60-64 | 3 | 37.1 | 21.43 | 2 | 53.5 | 37.80 | 1 | 23.0 | 23.03 |
| 65-69 | 8 | 133.9 | 47.35 | 6 | 233.7 | 95.42 | 2 | 58.7 | 41.51 |
| 70-74 | 3 | 66.8 | 38.58 | 1 | 54.6 | 54.61 | 2 | 75.2 | 53.21 |
| 75-79 | 3 | 90.2 | 52.06 | 2 | 156.3 | 110.49 | 1 | 48.9 | 48.85 |
| 80-84 | 4 | 220.5 | 110.25 | 1 | 155.5 | 155.52 | 3 | 256.2 | 147.91 |
| 85+ | 10 | 726.7 | 229.82 | 2 | 429.2 | 303.48 | 8 | 879.1 | 310.82 |
| Total deaths | 58 | | | 33 | | | 25 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 13.8 | - | | 16.1 | - | | 11.7 | - |
| . AAMR (1940 Std. mil.) | | 18.4 | 2.55 | | 24.9 | 4.59 | | 12.7 | 2.70 |
| . AAMR (1970 Std. mil.) | | 24.4 | 3.46 | | 30.1 | 5.91 | | 18.9 | 3.97 |
| . AAMR (2000 Std. mil.) | | 34.0 | 5.12 | | 37.7 | 7.99 | | 29.5 | 6.37 |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----HIV infection (042-044)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 1 | 2.6 | 2.58 | 0 | 0.0 | 0.00 | 1 | 5.3 | 5.29 |
| 20-24 | 3 | 6.7 | 3.85 | 2 | 8.9 | 6.31 | 1 | 4.4 | 4.43 |
| 25-29 | 15 | 33.4 | 8.62 | 9 | 41.1 | 13.68 | 6 | 26.1 | 10.65 |
| 30-34 | 21 | 55.3 | 12.07 | 18 | 97.7 | 23.02 | 3 | 15.4 | 8.87 |
| 35-39 | 17 | 57.2 | 13.87 | 14 | 97.8 | 26.13 | 3 | 19.5 | 11.24 |
| 40-44 | 7 | 28.6 | 10.81 | 5 | 43.0 | 19.24 | 2 | 15.5 | 10.99 |
| 45-49 | 6 | 33.1 | 13.50 | 6 | 70.2 | 28.68 | 0 | 0.0 | 0.00 |
| 50-54 | 5 | 36.8 | 16.46 | 2 | 31.5 | 22.29 | 3 | 41.4 | 23.91 |
| 55-59 | 1 | 9.5 | 9.47 | 1 | 20.3 | 20.31 | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | 1 | 26.7 | 26.73 | 0 | 0.0 | 0.00 |
| 65-69 | 1 | 16.7 | 16.74 | 1 | 39.0 | 38.96 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 79 | | | 60 | | | 19 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 18.9 | - | | 29.2 | - | | 8.9 | - |
| . AAMR (1940 Std. mil.) | | 19.6 | 2.30 | | 31.2 | 4.23 | | 9.0 | 2.15 |
| . AAMR (1970 Std. mil.) | | 16.6 | 2.06 | | 26.5 | 3.83 | | 7.7 | 1.90 |
| . AAMR (2000 Std. mil.) | | 20.0 | 2.45 | | 32.3 | 4.55 | | 8.9 | 2.23 |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 8.7 | - | | 15.7 | - | | 1.8 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): CT rate is higher
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All cancer (140-208)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 1 | 2.4 | 2.38 | 0 | 0.0 | 0.00 | 1 | 4.8 | 4.84 |
| 10-14 | 1 | 2.5 | 2.52 | 0 | 0.0 | 0.00 | 1 | 5.1 | 5.12 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 2 | 4.4 | 3.14 | 1 | 4.5 | 4.46 | 1 | 4.4 | 4.43 |
| 25-29 | 4 | 8.9 | 4.45 | 2 | 9.1 | 6.45 | 2 | 8.7 | 6.15 |
| 30-34 | 5 | 13.2 | 5.89 | 3 | 16.3 | 9.40 | 2 | 10.2 | 7.24 |
| 35-39 | 5 | 16.8 | 7.52 | 1 | 7.0 | 6.98 | 4 | 25.9 | 12.97 |
| 40-44 | 6 | 24.5 | 10.00 | 2 | 17.2 | 12.17 | 4 | 31.1 | 15.55 |
| 45-49 | 12 | 66.1 | 19.09 | 6 | 70.2 | 28.68 | 6 | 62.5 | 25.50 |
| 50-54 | 11 | 81.0 | 24.41 | 9 | 141.9 | 47.29 | 2 | 27.6 | 19.52 |
| 55-59 | 17 | 161.0 | 39.06 | 11 | 223.4 | 67.36 | 6 | 106.5 | 43.48 |
| 60-64 | 17 | 210.3 | 51.01 | 9 | 240.6 | 80.19 | 8 | 184.2 | 65.14 |
| 65-69 | 24 | 401.7 | 82.01 | 12 | 467.5 | 134.95 | 12 | 352.2 | 101.68 |
| 70-74 | 10 | 222.8 | 70.45 | 4 | 218.5 | 109.23 | 6 | 225.7 | 92.16 |
| 75-79 | 19 | 571.1 | 131.02 | 7 | 546.9 | 206.70 | 12 | 586.2 | 169.23 |
| 80-84 | 14 | 771.8 | 206.27 | 8 | 1244.2 | 439.88 | 6 | 512.4 | 209.18 |
| 85+ | 6 | 436.0 | 178.02 | 1 | 214.6 | 214.59 | 5 | 549.5 | 245.72 |
| Total deaths | 154 | | | 76 | | | 78 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 36.8 | - | | 37.0 | - | | 36.5 | - |
| . AAMR (1940 Std. mil.) | | 56.9 | 4.72 | | 66.7 | 7.83 | | 49.2 | 5.78 |
| . AAMR (1970 Std. mil.) | | 72.2 | 6.09 | | 84.2 | 10.28 | | 63.1 | 7.44 |
| . AAMR (2000 Std. mil.) | | 88.1 | 7.73 | | 101.4 | 13.14 | | 78.2 | 9.45 |
| . HP 2000 (1940 Std. mil.) | | 120.0 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 134.3 | - | | 165.4 | - | | 112.2 | - |
| . HP 2000 (1940 Std. mil.) | | 130.0 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Oral cavity or pharynx (140.0-149.9)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 45-49 | 2 | 11.0 | 7.79 | 2 | 23.4 | 16.56 | 0 | 0.0 | 0.00 |
| 50-54 | 2 | 14.7 | 10.41 | 2 | 31.5 | 22.29 | 0 | 0.0 | 0.00 |
| 55-59 | 1 | 9.5 | 9.47 | 1 | 20.3 | 20.31 | 0 | 0.0 | 0.00 |
| 60-64 | 2 | 24.7 | 17.50 | 1 | 26.7 | 26.73 | 1 | 23.0 | 23.03 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 7 | | | 6 | | | 1 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 1.7 | - | | 2.9 | - | | 0.5 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 2.4 | - | | 3.7 | - | | 1.3 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | 1 | 4.6 | 4.56 | 0 | 0.0 | 0.00 |
| 30-34 | 1 | 2.6 | 2.63 | 1 | 5.4 | 5.43 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 45-49 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 2 | 18.9 | 13.40 | 1 | 20.3 | 20.31 | 1 | 17.8 | 17.75 |
| 60-64 | 3 | 37.1 | 21.43 | 0 | 0.0 | 0.00 | 3 | 69.1 | 39.89 |
| 65-69 | 1 | 16.7 | 16.74 | 1 | 39.0 | 38.96 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 4 | 120.2 | 60.11 | 1 | 78.1 | 78.13 | 3 | 146.6 | 84.61 |
| 80-84 | 1 | 55.1 | 55.13 | 0 | 0.0 | 0.00 | 1 | 85.4 | 85.40 |
| 85+ | 2 | 145.3 | 102.78 | 0 | 0.0 | 0.00 | 2 | 219.8 | 155.41 |
| Total deaths | 15 | | | 5 | | | 10 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 3.6 | - | | 2.4 | - | | 4.7 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 14.0 | - | | 17.0 | - | | 11.7 | - |
| . HP 2000 (1940 Std. mil.) | | 13.2 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Rectal cancer (154)-----

| Age | Both Sexes | | | Males | | | Females | | |
|--|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|----------------------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 45-49 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 80-84 | 1 | 55.1 | 55.13 | 0 | - | - | 1 | 85.4 | 85.40 |
| 85+ | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| Total deaths | 1 | | | - | | | 1 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 0.2 | - | | - | - | | 0.5 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 1.9 | - | | 2.5 | - | | 1.5 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs: | | | | | | | | | |
| . The CT HP 2000 target: | | | | | | | | | No comparison is available |
| . The U.S. AAMR (1940 Std. mil.): | | | | | | | | | No comparison is available |
| . THE U.S. HP 2000 target: | | | | | | | | | No comparison is available |

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 45-49 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | 0 | 0.0 | 0.00 | 1 | 23.0 | 23.03 |
| 65-69 | 1 | 16.7 | 16.74 | 1 | 39.0 | 38.96 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 2 | | | 1 | | | 1 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 0.5 | - | | 0.5 | - | | 0.5 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 6.4 | - | | 7.5 | - | | 5.4 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung and other respiratory cancer (160-165)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 1 | 4.1 | 4.08 | 1 | 8.6 | 8.60 | 0 | 0.0 | 0.00 |
| 45-49 | 3 | 16.5 | 9.55 | 2 | 23.4 | 16.56 | 1 | 10.4 | 10.41 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 3 | 28.4 | 16.41 | 3 | 60.9 | 35.18 | 0 | 0.0 | 0.00 |
| 60-64 | 2 | 24.7 | 17.50 | 2 | 53.5 | 37.80 | 0 | 0.0 | 0.00 |
| 65-69 | 8 | 133.9 | 47.35 | 3 | 116.9 | 67.47 | 5 | 146.8 | 65.63 |
| 70-74 | 3 | 66.8 | 38.58 | 2 | 109.2 | 77.24 | 1 | 37.6 | 37.62 |
| 75-79 | 6 | 180.3 | 73.62 | 2 | 156.3 | 110.49 | 4 | 195.4 | 97.70 |
| 80-84 | 4 | 220.5 | 110.25 | 4 | 622.1 | 311.04 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 30 | | | 19 | | | 11 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 7.2 | - | | 9.2 | - | | 5.2 | - |
| . AAMR (1940 Std. mil.) | | 11.9 | 2.21 | | 17.6 | 4.08 | | - | - |
| . AAMR (1970 Std. mil.) | | 16.0 | 2.96 | | 24.0 | 5.75 | | - | - |
| . AAMR (2000 Std. mil.) | | 19.4 | 3.69 | | 30.2 | 7.62 | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 40.7 | - | | 60.4 | - | | 25.7 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung cancer (162.2-.9)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 1 | 4.1 | 4.08 | 1 | 8.6 | 8.60 | 0 | 0.0 | 0.00 |
| 45-49 | 3 | 16.5 | 9.55 | 2 | 23.4 | 16.56 | 1 | 10.4 | 10.41 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 3 | 28.4 | 16.41 | 3 | 60.9 | 35.18 | 0 | 0.0 | 0.00 |
| 60-64 | 2 | 24.7 | 17.50 | 2 | 53.5 | 37.80 | 0 | 0.0 | 0.00 |
| 65-69 | 8 | 133.9 | 47.35 | 3 | 116.9 | 67.47 | 5 | 146.8 | 65.63 |
| 70-74 | 3 | 66.8 | 38.58 | 2 | 109.2 | 77.24 | 1 | 37.6 | 37.62 |
| 75-79 | 6 | 180.3 | 73.62 | 2 | 156.3 | 110.49 | 4 | 195.4 | 97.70 |
| 80-84 | 4 | 220.5 | 110.25 | 4 | 622.1 | 311.04 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 30 | | | 19 | | | 11 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 7.2 | - | | 9.2 | - | | 5.2 | - |
| . AAMR (1940 Std. mil.) | | 11.9 | 2.21 | | 17.6 | 4.08 | | - | - |
| . AAMR (1970 Std. mil.) | | 16.0 | 2.96 | | 24.0 | 5.75 | | - | - |
| . AAMR (2000 Std. mil.) | | 19.4 | 3.69 | | 30.2 | 7.62 | | - | - |
| . HP 2000 (1940 Std. mil.) | | 42.0 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 39.3 | - | | 57.9 | - | | 25.0 | - |
| . HP 2000 (1940 Std. mil.) | | 42.0 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Malignant melanoma of skin (172)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 45-49 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 65-69 | 1 | 16.7 | 16.74 | 0 | - | - | 1 | 29.4 | 29.35 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| Total deaths | 1 | | | - | | | 1 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 0.2 | - | | - | - | | 0.5 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 1.8 | - | | 2.6 | - | | 1.2 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Female breast cancer (174)-----

| Age | Both Sexes | | | Males | | | Females | | |
|--|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|----------------------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | - | - | - | 1 | 4.3 | 4.35 |
| 30-34 | 1 | 2.6 | 2.63 | - | - | - | 1 | 5.1 | 5.12 |
| 35-39 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 45-49 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | - | - | - | 1 | 23.0 | 23.03 |
| 65-69 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 80-84 | 2 | 110.3 | 77.96 | - | - | - | 2 | 170.8 | 120.77 |
| 85+ | 1 | 72.7 | 72.67 | - | - | - | 1 | 109.9 | 109.89 |
| Total deaths | 6 | | | - | | | 6 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | - | - | | - | - | | 2.8 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | 23.1 | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | 23.1 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | 20.6 | - |
| Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs: | | | | | | | | | |
| . The CT HP 2000 target: | | | | | | | | | No comparison is available |
| . The U.S. AAMR (1940 Std. mil.): | | | | | | | | | No comparison is available |
| . THE U.S. HP 2000 target: | | | | | | | | | No comparison is available |

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cervical cancer (180)-----

| Age | Both Sexes | | | Males | | | Females | | |
|--|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|----------------------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 45-49 | 1 | 5.5 | 5.51 | - | - | - | 1 | 10.4 | 10.41 |
| 50-54 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 60-64 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 65-69 | 1 | 16.7 | 16.74 | - | - | - | 1 | 29.4 | 29.35 |
| 70-74 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 75-79 | 1 | 30.1 | 30.06 | - | - | - | 1 | 48.9 | 48.85 |
| 80-84 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| Total deaths | 3 | | | - | | | 3 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | - | - | | - | - | | 1.4 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | 1.1 | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | 2.6 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | 1.3 | - |
| Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs: | | | | | | | | | |
| . The CT HP 2000 target: | | | | | | | | | No comparison is available |
| . The U.S. AAMR (1940 Std. mil.): | | | | | | | | | No comparison is available |
| . THE U.S. HP 2000 target: | | | | | | | | | No comparison is available |

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | - | - | - | - | - | 0 | - | - |
| 5-9 | 0 | - | - | - | - | - | 0 | - | - |
| 10-14 | 0 | - | - | - | - | - | 0 | - | - |
| 15-19 | 0 | - | - | - | - | - | 0 | - | - |
| 20-24 | 0 | - | - | - | - | - | 0 | - | - |
| 25-29 | 0 | - | - | - | - | - | 0 | - | - |
| 30-34 | 0 | - | - | - | - | - | 0 | - | - |
| 35-39 | 0 | - | - | - | - | - | 0 | - | - |
| 40-44 | 0 | - | - | - | - | - | 0 | - | - |
| 45-49 | 0 | - | - | - | - | - | 0 | - | - |
| 50-54 | 0 | - | - | - | - | - | 0 | - | - |
| 55-59 | 0 | - | - | - | - | - | 0 | - | - |
| 60-64 | 0 | - | - | - | - | - | 0 | - | - |
| 65-69 | 0 | - | - | - | - | - | 0 | - | - |
| 70-74 | 0 | - | - | - | - | - | 0 | - | - |
| 75-79 | 0 | - | - | - | - | - | 0 | - | - |
| 80-84 | 0 | - | - | - | - | - | 0 | - | - |
| 85+ | 0 | - | - | - | - | - | 0 | - | - |
| Total deaths | - | | | - | | | - | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | - | - | | - | - | | - | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | 1.2 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 45-49 | 2 | 11.0 | 7.79 | - | - | - | 2 | 20.8 | 14.73 |
| 50-54 | 1 | 7.4 | 7.36 | - | - | - | 1 | 13.8 | 13.80 |
| 55-59 | 2 | 18.9 | 13.40 | - | - | - | 2 | 35.5 | 25.11 |
| 60-64 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 70-74 | 1 | 22.3 | 22.28 | - | - | - | 1 | 37.6 | 37.62 |
| 75-79 | 1 | 30.1 | 30.06 | - | - | - | 1 | 48.9 | 48.85 |
| 80-84 | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | - | - | - | 0 | 0.0 | 0.00 |
| Total deaths | 7 | | | - | | | 7 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | - | - | | - | - | | 3.3 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | 6.3 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Bladder cancer (188)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 45-49 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 70-74 | 1 | 22.3 | 22.28 | 1 | 54.6 | 54.61 | 0 | - | - |
| 75-79 | 1 | 30.1 | 30.06 | 1 | 78.1 | 78.13 | 0 | - | - |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| Total deaths | 2 | | | 2 | | | - | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 0.5 | - | | 1.0 | - | | - | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 2.2 | - | | 3.8 | - | | 1.1 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Brain & other CNS cancer (191-192)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | 1 | 4.6 | 4.56 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 1 | 4.1 | 4.08 | 0 | 0.0 | 0.00 | 1 | 7.8 | 7.77 |
| 45-49 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 50-54 | 1 | 7.4 | 7.36 | 0 | 0.0 | 0.00 | 1 | 13.8 | 13.80 |
| 55-59 | 1 | 9.5 | 9.47 | 0 | 0.0 | 0.00 | 1 | 17.8 | 17.75 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 4 | | | 1 | | | 3 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 1.0 | - | | 0.5 | - | | 1.4 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 3.5 | - | | 4.4 | - | | 2.9 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Leukemia (204-208)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 1 | 2.5 | 2.52 | 0 | 0.0 | 0.00 | 1 | 5.1 | 5.12 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | 0 | 0.0 | 0.00 | 1 | 4.3 | 4.35 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 2 | 6.7 | 4.76 | 0 | 0.0 | 0.00 | 2 | 13.0 | 9.17 |
| 40-44 | 1 | 4.1 | 4.08 | 1 | 8.6 | 8.60 | 0 | 0.0 | 0.00 |
| 45-49 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 1 | 9.5 | 9.47 | 0 | 0.0 | 0.00 | 1 | 17.8 | 17.75 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 70-74 | 1 | 22.3 | 22.28 | 1 | 54.6 | 54.61 | 0 | 0.0 | 0.00 |
| 75-79 | 1 | 30.1 | 30.06 | 1 | 78.1 | 78.13 | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 8 | | | 3 | | | 5 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 1.9 | - | | 1.5 | - | | 2.3 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 5.1 | - | | 6.5 | - | | 3.9 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes mellitus (250)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | 1 | 4.6 | 4.56 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 1 | 4.1 | 4.08 | 0 | 0.0 | 0.00 | 1 | 7.8 | 7.77 |
| 45-49 | 1 | 5.5 | 5.51 | 1 | 11.7 | 11.71 | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 3 | 28.4 | 16.41 | 1 | 20.3 | 20.31 | 2 | 35.5 | 25.11 |
| 60-64 | 4 | 49.5 | 24.74 | 1 | 26.7 | 26.73 | 3 | 69.1 | 39.89 |
| 65-69 | 3 | 50.2 | 28.99 | 1 | 39.0 | 38.96 | 2 | 58.7 | 41.51 |
| 70-74 | 4 | 89.1 | 44.55 | 1 | 54.6 | 54.61 | 3 | 112.9 | 65.16 |
| 75-79 | 3 | 90.2 | 52.06 | 1 | 78.1 | 78.13 | 2 | 97.7 | 69.09 |
| 80-84 | 4 | 220.5 | 110.25 | 1 | 155.5 | 155.52 | 3 | 256.2 | 147.91 |
| 85+ | 3 | 218.0 | 125.88 | 0 | 0.0 | 0.00 | 3 | 329.7 | 190.34 |
| Total deaths | 27 | | | 8 | | | 19 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 6.4 | - | | 3.9 | - | | 8.9 | - |
| . AAMR (1940 Std. mil.) | | 10.0 | 1.97 | | - | - | | 12.0 | 2.88 |
| . AAMR (1970 Std. mil.) | | 14.1 | 2.78 | | - | - | | 17.3 | 4.02 |
| . AAMR (2000 Std. mil.) | | 18.5 | 3.81 | | - | - | | 23.0 | 5.45 |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 11.6 | - | | 12.3 | - | | 10.9 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | 1 | 4.6 | 4.56 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 3 | 10.1 | 5.83 | 1 | 7.0 | 6.98 | 2 | 13.0 | 9.17 |
| 40-44 | 5 | 20.4 | 9.13 | 3 | 25.8 | 14.90 | 2 | 15.5 | 10.99 |
| 45-49 | 3 | 16.5 | 9.55 | 2 | 23.4 | 16.56 | 1 | 10.4 | 10.41 |
| 50-54 | 4 | 29.4 | 14.72 | 4 | 63.1 | 31.53 | 0 | 0.0 | 0.00 |
| 55-59 | 13 | 123.1 | 34.15 | 5 | 101.5 | 45.41 | 8 | 142.0 | 50.21 |
| 60-64 | 17 | 210.3 | 51.01 | 7 | 187.1 | 70.72 | 10 | 230.3 | 72.83 |
| 65-69 | 20 | 334.8 | 74.86 | 11 | 428.5 | 129.20 | 9 | 264.2 | 88.05 |
| 70-74 | 13 | 289.6 | 80.32 | 5 | 273.1 | 122.12 | 8 | 301.0 | 106.41 |
| 75-79 | 9 | 270.5 | 90.17 | 3 | 234.4 | 135.32 | 6 | 293.1 | 119.66 |
| 80-84 | 11 | 606.4 | 182.83 | 6 | 933.1 | 380.95 | 5 | 427.0 | 190.95 |
| 85+ | 6 | 436.0 | 178.02 | 1 | 214.6 | 214.59 | 5 | 549.5 | 245.72 |
| Total deaths | 105 | | | 49 | | | 56 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 25.1 | - | | 23.9 | - | | 26.2 | - |
| . AAMR (1940 Std. mil.) | | 41.1 | 4.10 | | 45.1 | 6.57 | | 38.1 | 5.25 |
| . AAMR (1970 Std. mil.) | | 53.9 | 5.39 | | 58.5 | 8.76 | | 50.6 | 6.86 |
| . AAMR (2000 Std. mil.) | | 65.4 | 6.79 | | 70.5 | 11.17 | | 61.6 | 8.54 |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | 34.0 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | - | - | 0 | - | - | 0 | - | - |
| 5-9 | 0 | - | - | 0 | - | - | 0 | - | - |
| 10-14 | 0 | - | - | 0 | - | - | 0 | - | - |
| 15-19 | 0 | - | - | 0 | - | - | 0 | - | - |
| 20-24 | 0 | - | - | 0 | - | - | 0 | - | - |
| 25-29 | 0 | - | - | 0 | - | - | 0 | - | - |
| 30-34 | 0 | - | - | 0 | - | - | 0 | - | - |
| 35-39 | 0 | - | - | 0 | - | - | 0 | - | - |
| 40-44 | 0 | - | - | 0 | - | - | 0 | - | - |
| 45-49 | 0 | - | - | 0 | - | - | 0 | - | - |
| 50-54 | 0 | - | - | 0 | - | - | 0 | - | - |
| 55-59 | 0 | - | - | 0 | - | - | 0 | - | - |
| 60-64 | 0 | - | - | 0 | - | - | 0 | - | - |
| 65-69 | 0 | - | - | 0 | - | - | 0 | - | - |
| 70-74 | 0 | - | - | 0 | - | - | 0 | - | - |
| 75-79 | 0 | - | - | 0 | - | - | 0 | - | - |
| 80-84 | 0 | - | - | 0 | - | - | 0 | - | - |
| 85+ | 0 | - | - | 0 | - | - | 0 | - | - |
| Total deaths | - | | | - | | | - | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | - | - | | - | - | | - | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 0.6 | - | | 0.6 | - | | 0.4 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Alzheimer's diseases (331.0)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 45-49 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 85+ | 1 | 72.7 | 72.67 | 1 | 214.6 | 214.59 | 0 | - | - |
| Total deaths | 1 | | | 1 | | | - | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 0.2 | - | | 0.5 | - | | - | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 2.1 | - | | 2.4 | - | | 2.0 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cardiovascular disease (390-459)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 2 | 4.1 | 2.89 | 1 | 4.0 | 3.99 | 1 | 4.2 | 4.19 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 1 | 2.6 | 2.58 | 0 | 0.0 | 0.00 | 1 | 5.3 | 5.29 |
| 20-24 | 1 | 2.2 | 2.22 | 1 | 4.5 | 4.46 | 0 | 0.0 | 0.00 |
| 25-29 | 6 | 13.4 | 5.45 | 5 | 22.8 | 10.20 | 1 | 4.3 | 4.35 |
| 30-34 | 8 | 21.1 | 7.45 | 5 | 27.1 | 12.13 | 3 | 15.4 | 8.87 |
| 35-39 | 6 | 20.2 | 8.24 | 4 | 27.9 | 13.97 | 2 | 13.0 | 9.17 |
| 40-44 | 12 | 49.0 | 14.15 | 10 | 86.0 | 27.21 | 2 | 15.5 | 10.99 |
| 45-49 | 19 | 104.7 | 24.02 | 13 | 152.2 | 42.21 | 6 | 62.5 | 25.50 |
| 50-54 | 20 | 147.2 | 32.91 | 9 | 141.9 | 47.29 | 11 | 151.8 | 45.78 |
| 55-59 | 22 | 208.4 | 44.43 | 9 | 182.8 | 60.93 | 13 | 230.8 | 64.01 |
| 60-64 | 25 | 309.3 | 61.86 | 15 | 401.0 | 103.53 | 10 | 230.3 | 72.83 |
| 65-69 | 35 | 585.9 | 99.03 | 20 | 779.1 | 174.22 | 15 | 440.3 | 113.68 |
| 70-74 | 33 | 735.1 | 127.97 | 17 | 928.5 | 225.18 | 16 | 602.0 | 150.49 |
| 75-79 | 30 | 901.7 | 164.63 | 18 | 1406.3 | 331.46 | 12 | 586.2 | 169.23 |
| 80-84 | 43 | 2370.5 | 361.49 | 21 | 3265.9 | 712.69 | 22 | 1878.7 | 400.55 |
| 85+ | 53 | 3851.7 | 529.08 | 16 | 3433.5 | 858.37 | 37 | 4065.9 | 668.44 |
| Total deaths | 316 | | | 164 | | | 152 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 75.4 | - | | 79.8 | - | | 71.2 | - |
| . AAMR (1940 Std. mil.) | | 109.3 | 6.39 | | 137.6 | 11.09 | | 88.5 | 7.60 |
| . AAMR (1970 Std. mil.) | | 156.6 | 9.13 | | 193.6 | 16.18 | | 130.4 | 10.77 |
| . AAMR (2000 Std. mil.) | | 217.3 | 13.29 | | 263.2 | 23.59 | | 185.1 | 15.77 |
| . HP 2000 (1940 Std. mil.) | | 84.4 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 197.2 | - | | 260.2 | - | | 147.5 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diseases of the heart (390-398,402,404-429)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 5 | 11.1 | 4.98 | 4 | 18.2 | 9.12 | 1 | 4.3 | 4.35 |
| 30-34 | 6 | 15.8 | 6.45 | 5 | 27.1 | 12.13 | 1 | 5.1 | 5.12 |
| 35-39 | 4 | 13.5 | 6.73 | 2 | 14.0 | 9.88 | 2 | 13.0 | 9.17 |
| 40-44 | 8 | 32.7 | 11.55 | 7 | 60.2 | 22.76 | 1 | 7.8 | 7.77 |
| 45-49 | 10 | 55.1 | 17.43 | 9 | 105.4 | 35.12 | 1 | 10.4 | 10.41 |
| 50-54 | 17 | 125.1 | 30.34 | 8 | 126.1 | 44.58 | 9 | 124.2 | 41.41 |
| 55-59 | 16 | 151.6 | 37.89 | 5 | 101.5 | 45.41 | 11 | 195.3 | 58.88 |
| 60-64 | 21 | 259.8 | 56.69 | 12 | 320.8 | 92.60 | 9 | 207.3 | 69.09 |
| 65-69 | 27 | 452.0 | 86.98 | 16 | 623.3 | 155.82 | 11 | 322.9 | 97.35 |
| 70-74 | 25 | 556.9 | 111.38 | 12 | 655.4 | 189.19 | 13 | 489.1 | 135.65 |
| 75-79 | 23 | 691.3 | 144.15 | 15 | 1171.9 | 302.58 | 8 | 390.8 | 138.17 |
| 80-84 | 32 | 1764.1 | 311.84 | 15 | 2332.8 | 602.33 | 17 | 1451.8 | 352.10 |
| 85+ | 38 | 2761.6 | 448.00 | 12 | 2575.1 | 743.37 | 26 | 2857.1 | 560.33 |
| Total deaths | 233 | | | 123 | | | 110 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 55.6 | - | | 59.9 | - | | 51.5 | - |
| . AAMR (1940 Std. mil.) | | 81.6 | 5.55 | | 103.5 | 9.63 | | 65.6 | 6.61 |
| . AAMR (1970 Std. mil.) | | 116.9 | 7.91 | | 145.6 | 14.04 | | 96.7 | 9.34 |
| . AAMR (2000 Std. mil.) | | 161.1 | 11.43 | | 197.7 | 20.42 | | 135.7 | 13.51 |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 157.5 | - | | 213.9 | - | | 112.6 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Heart disease-related (390-398,402,404-429)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 4 | 8.2 | 4.09 | 2 | 8.0 | 5.64 | 2 | 8.4 | 5.93 |
| 5-9 | 1 | 2.4 | 2.38 | 0 | 0.0 | 0.00 | 1 | 4.8 | 4.84 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 2 | 5.2 | 3.65 | 2 | 10.1 | 7.12 | 0 | 0.0 | 0.00 |
| 20-24 | 1 | 2.2 | 2.22 | 1 | 4.5 | 4.46 | 0 | 0.0 | 0.00 |
| 25-29 | 16 | 35.6 | 8.90 | 13 | 59.3 | 16.45 | 3 | 13.0 | 7.53 |
| 30-34 | 15 | 39.5 | 10.20 | 9 | 48.8 | 16.28 | 6 | 30.7 | 12.54 |
| 35-39 | 25 | 84.1 | 16.82 | 15 | 104.8 | 27.05 | 10 | 64.9 | 20.51 |
| 40-44 | 18 | 73.5 | 17.33 | 15 | 129.1 | 33.32 | 3 | 23.3 | 13.47 |
| 45-49 | 21 | 115.7 | 25.26 | 15 | 175.6 | 45.35 | 6 | 62.5 | 25.50 |
| 50-54 | 28 | 206.1 | 38.94 | 15 | 236.4 | 61.05 | 13 | 179.5 | 49.77 |
| 55-59 | 30 | 284.2 | 51.88 | 14 | 284.3 | 75.99 | 16 | 284.0 | 71.01 |
| 60-64 | 35 | 433.0 | 73.19 | 20 | 534.6 | 119.54 | 15 | 345.5 | 89.20 |
| 65-69 | 47 | 786.7 | 114.76 | 28 | 1090.8 | 206.14 | 19 | 557.7 | 127.94 |
| 70-74 | 39 | 868.8 | 139.12 | 17 | 928.5 | 225.18 | 22 | 827.7 | 176.46 |
| 75-79 | 39 | 1172.2 | 187.71 | 20 | 1562.5 | 349.39 | 19 | 928.2 | 212.94 |
| 80-84 | 46 | 2535.8 | 373.89 | 22 | 3421.5 | 729.46 | 24 | 2049.5 | 418.36 |
| 85+ | 65 | 4723.8 | 585.92 | 19 | 4077.3 | 935.39 | 46 | 5054.9 | 745.31 |
| Total deaths | 432 | | | 227 | | | 205 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 103.1 | - | | 110.5 | - | | 96.1 | - |
| . AAMR (1940 Std. mil.) | | 146.3 | 7.34 | | 181.5 | 12.55 | | 118.4 | 8.75 |
| . AAMR (1970 Std. mil.) | | 201.7 | 10.22 | | 242.7 | 17.68 | | 170.3 | 12.25 |
| . AAMR (2000 Std. mil.) | | 275.7 | 14.71 | | 323.2 | 25.41 | | 239.5 | 17.75 |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | 1 | 4.6 | 4.56 | 0 | 0.0 | 0.00 |
| 30-34 | 1 | 2.6 | 2.63 | 1 | 5.4 | 5.43 | 0 | 0.0 | 0.00 |
| 35-39 | 2 | 6.7 | 4.76 | 2 | 14.0 | 9.88 | 0 | 0.0 | 0.00 |
| 40-44 | 7 | 28.6 | 10.81 | 6 | 51.6 | 21.07 | 1 | 7.8 | 7.77 |
| 45-49 | 6 | 33.1 | 13.50 | 6 | 70.2 | 28.68 | 0 | 0.0 | 0.00 |
| 50-54 | 10 | 73.6 | 23.27 | 5 | 78.8 | 35.25 | 5 | 69.0 | 30.87 |
| 55-59 | 11 | 104.2 | 31.42 | 4 | 81.2 | 40.62 | 7 | 124.3 | 46.97 |
| 60-64 | 15 | 185.6 | 47.92 | 9 | 240.6 | 80.19 | 6 | 138.2 | 56.41 |
| 65-69 | 19 | 318.0 | 72.96 | 10 | 389.6 | 123.19 | 9 | 264.2 | 88.05 |
| 70-74 | 18 | 401.0 | 94.51 | 10 | 546.1 | 172.71 | 8 | 301.0 | 106.41 |
| 75-79 | 15 | 450.9 | 116.41 | 10 | 781.3 | 247.05 | 5 | 244.3 | 109.24 |
| 80-84 | 27 | 1488.4 | 286.45 | 15 | 2332.8 | 602.33 | 12 | 1024.8 | 295.82 |
| 85+ | 26 | 1889.5 | 370.57 | 8 | 1716.7 | 606.96 | 18 | 1978.0 | 466.22 |
| Total deaths | 158 | | | 87 | | | 71 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 37.7 | - | | 42.3 | - | | 33.3 | - |
| . AAMR (1940 Std. mil.) | | 56.3 | 4.63 | | 75.5 | 8.28 | | 42.5 | 5.34 |
| . AAMR (1970 Std. mil.) | | 82.3 | 6.69 | | 108.9 | 12.31 | | 64.2 | 7.66 |
| . AAMR (2000 Std. mil.) | | 114.1 | 9.70 | | 149.4 | 17.97 | | 90.5 | 11.11 |
| . HP 2000 (1940 Std. mil.) | | 84.0 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 125.6 | - | | 174.7 | - | | 87.0 | - |
| . HP 2000 (1940 Std. mil.) | | 100.0 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Congestive heart failure (CHF) (428.0)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 45-49 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | 1 | 26.7 | 26.73 | 0 | 0.0 | 0.00 |
| 65-69 | 1 | 16.7 | 16.74 | 1 | 39.0 | 38.96 | 0 | 0.0 | 0.00 |
| 70-74 | 1 | 22.3 | 22.28 | 1 | 54.6 | 54.61 | 0 | 0.0 | 0.00 |
| 75-79 | 1 | 30.1 | 30.06 | 0 | 0.0 | 0.00 | 1 | 48.9 | 48.85 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 3 | 218.0 | 125.88 | 0 | 0.0 | 0.00 | 3 | 329.7 | 190.34 |
| Total deaths | 7 | | | 3 | | | 4 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 1.7 | - | | 1.5 | - | | 1.9 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 5.7 | - | | 6.6 | - | | 5.2 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension with or without renal disease (401,403)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 40-44 | 1 | 4.1 | 4.08 | 1 | 8.6 | 8.60 | 0 | - | - |
| 45-49 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 55-59 | 1 | 9.5 | 9.47 | 1 | 20.3 | 20.31 | 0 | - | - |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 70-74 | 1 | 22.3 | 22.28 | 1 | 54.6 | 54.61 | 0 | - | - |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| Total deaths | 3 | | | 3 | | | - | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 0.7 | - | | 1.5 | - | | - | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 1.8 | - | | 2.1 | - | | 1.7 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension-related (401,403)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 4 | 16.3 | 8.17 | 3 | 25.8 | 14.90 | 1 | 7.8 | 7.77 |
| 45-49 | 5 | 27.6 | 12.32 | 3 | 35.1 | 20.28 | 2 | 20.8 | 14.73 |
| 50-54 | 5 | 36.8 | 16.46 | 2 | 31.5 | 22.29 | 3 | 41.4 | 23.91 |
| 55-59 | 6 | 56.8 | 23.20 | 2 | 40.6 | 28.72 | 4 | 71.0 | 35.51 |
| 60-64 | 4 | 49.5 | 24.74 | 0 | 0.0 | 0.00 | 4 | 92.1 | 46.06 |
| 65-69 | 6 | 100.4 | 41.00 | 2 | 77.9 | 55.09 | 4 | 117.4 | 58.70 |
| 70-74 | 8 | 178.2 | 63.01 | 5 | 273.1 | 122.12 | 3 | 112.9 | 65.16 |
| 75-79 | 4 | 120.2 | 60.11 | 0 | 0.0 | 0.00 | 4 | 195.4 | 97.70 |
| 80-84 | 6 | 330.8 | 135.03 | 5 | 777.6 | 347.76 | 1 | 85.4 | 85.40 |
| 85+ | 4 | 290.7 | 145.35 | 0 | 0.0 | 0.00 | 4 | 439.6 | 219.78 |
| Total deaths | 52 | | | 22 | | | 30 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 12.4 | - | | 10.7 | - | | 14.1 | - |
| . AAMR (1940 Std. mil.) | | 19.6 | 2.78 | | 19.6 | 4.25 | | 20.1 | 3.80 |
| . AAMR (1970 Std. mil.) | | 25.9 | 3.68 | | 26.1 | 5.94 | | 26.3 | 4.86 |
| . AAMR (2000 Std. mil.) | | 33.1 | 4.90 | | 33.8 | 8.04 | | 33.0 | 6.26 |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cerebrovascular disease (430-438)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 0 | 0.0 | 0.00 | 1 | 4.2 | 4.19 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 1 | 2.2 | 2.22 | 1 | 4.5 | 4.46 | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | 1 | 4.6 | 4.56 | 0 | 0.0 | 0.00 |
| 30-34 | 2 | 5.3 | 3.72 | 0 | 0.0 | 0.00 | 2 | 10.2 | 7.24 |
| 35-39 | 1 | 3.4 | 3.36 | 1 | 7.0 | 6.98 | 0 | 0.0 | 0.00 |
| 40-44 | 2 | 8.2 | 5.78 | 1 | 8.6 | 8.60 | 1 | 7.8 | 7.77 |
| 45-49 | 8 | 44.1 | 15.59 | 4 | 46.8 | 23.42 | 4 | 41.6 | 20.82 |
| 50-54 | 3 | 22.1 | 12.75 | 1 | 15.8 | 15.76 | 2 | 27.6 | 19.52 |
| 55-59 | 5 | 47.4 | 21.18 | 3 | 60.9 | 35.18 | 2 | 35.5 | 25.11 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 65-69 | 5 | 83.7 | 37.43 | 3 | 116.9 | 67.47 | 2 | 58.7 | 41.51 |
| 70-74 | 4 | 89.1 | 44.55 | 2 | 109.2 | 77.24 | 2 | 75.2 | 53.21 |
| 75-79 | 6 | 180.3 | 73.62 | 2 | 156.3 | 110.49 | 4 | 195.4 | 97.70 |
| 80-84 | 9 | 496.1 | 165.38 | 5 | 777.6 | 347.76 | 4 | 341.6 | 170.79 |
| 85+ | 11 | 799.4 | 241.03 | 1 | 214.6 | 214.59 | 10 | 1098.9 | 347.50 |
| Total deaths | 59 | | | 25 | | | 34 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 14.1 | - | | 12.2 | - | | 15.9 | - |
| . AAMR (1940 Std. mil.) | | 19.2 | 2.59 | | 20.8 | 4.28 | | 17.8 | 3.24 |
| . AAMR (1970 Std. mil.) | | 27.8 | 3.79 | | 28.4 | 6.18 | | 26.9 | 4.75 |
| . AAMR (2000 Std. mil.) | | 40.4 | 5.80 | | 38.1 | 8.81 | | 40.9 | 7.49 |
| . HP 2000 (1940 Std. mil.) | | 16.8 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 28.1 | - | | 31.0 | - | | 26.1 | - |
| . HP 2000 (1940 Std. mil.) | | 20.0 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Atherosclerosis (440)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 45-49 | 1 | 5.5 | 5.51 | 0 | 0.0 | 0.00 | 1 | 10.4 | 10.41 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | 1 | 26.7 | 26.73 | 0 | 0.0 | 0.00 |
| 65-69 | 1 | 16.7 | 16.74 | 0 | 0.0 | 0.00 | 1 | 29.4 | 29.35 |
| 70-74 | 1 | 22.3 | 22.28 | 1 | 54.6 | 54.61 | 0 | 0.0 | 0.00 |
| 75-79 | 1 | 30.1 | 30.06 | 1 | 78.1 | 78.13 | 0 | 0.0 | 0.00 |
| 80-84 | 1 | 55.1 | 55.13 | 1 | 155.5 | 155.52 | 0 | 0.0 | 0.00 |
| 85+ | 2 | 145.3 | 102.78 | 2 | 429.2 | 303.48 | 0 | 0.0 | 0.00 |
| Total deaths | 8 | | | 6 | | | 2 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 1.9 | - | | 2.9 | - | | 0.9 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 3.0 | - | | 3.4 | - | | 2.7 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Atherosclerosis-related (440)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 45-49 | 1 | 5.5 | 5.51 | 0 | 0.0 | 0.00 | 1 | 10.4 | 10.41 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 1 | 9.5 | 9.47 | 0 | 0.0 | 0.00 | 1 | 17.8 | 17.75 |
| 60-64 | 4 | 49.5 | 24.74 | 2 | 53.5 | 37.80 | 2 | 46.1 | 32.57 |
| 65-69 | 2 | 33.5 | 23.67 | 0 | 0.0 | 0.00 | 2 | 58.7 | 41.51 |
| 70-74 | 3 | 66.8 | 38.58 | 3 | 163.8 | 94.60 | 0 | 0.0 | 0.00 |
| 75-79 | 1 | 30.1 | 30.06 | 1 | 78.1 | 78.13 | 0 | 0.0 | 0.00 |
| 80-84 | 4 | 220.5 | 110.25 | 3 | 466.6 | 269.37 | 1 | 85.4 | 85.40 |
| 85+ | 9 | 654.1 | 218.02 | 4 | 858.4 | 429.18 | 5 | 549.5 | 245.72 |
| Total deaths | 25 | | | 13 | | | 12 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 6.0 | - | | 6.3 | - | | 5.6 | - |
| . AAMR (1940 Std. mil.) | | 8.3 | 1.74 | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | 13.7 | 2.76 | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | 20.9 | 4.41 | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pneumonia & influenza (480-487)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 45-49 | 1 | 5.5 | 5.51 | 1 | 11.7 | 11.71 | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 1 | 9.5 | 9.47 | 1 | 20.3 | 20.31 | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | 1 | 26.7 | 26.73 | 0 | 0.0 | 0.00 |
| 65-69 | 2 | 33.5 | 23.67 | 2 | 77.9 | 55.09 | 0 | 0.0 | 0.00 |
| 70-74 | 2 | 44.6 | 31.50 | 1 | 54.6 | 54.61 | 1 | 37.6 | 37.62 |
| 75-79 | 4 | 120.2 | 60.11 | 1 | 78.1 | 78.13 | 3 | 146.6 | 84.61 |
| 80-84 | 4 | 220.5 | 110.25 | 2 | 311.0 | 219.94 | 2 | 170.8 | 120.77 |
| 85+ | 11 | 799.4 | 241.03 | 4 | 858.4 | 429.18 | 7 | 769.2 | 290.74 |
| Total deaths | 26 | | | 13 | | | 13 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 6.2 | - | | 6.3 | - | | 6.1 | - |
| . AAMR (1940 Std. mil.) | | 7.9 | 1.63 | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | 14.4 | 2.83 | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | 23.5 | 4.77 | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 13.7 | - | | 18.2 | - | | 10.9 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | 1 | 4.6 | 4.56 | 0 | 0.0 | 0.00 |
| 30-34 | 1 | 2.6 | 2.63 | 1 | 5.4 | 5.43 | 0 | 0.0 | 0.00 |
| 35-39 | 1 | 3.4 | 3.36 | 0 | 0.0 | 0.00 | 1 | 6.5 | 6.49 |
| 40-44 | 1 | 4.1 | 4.08 | 0 | 0.0 | 0.00 | 1 | 7.8 | 7.77 |
| 45-49 | 1 | 5.5 | 5.51 | 0 | 0.0 | 0.00 | 1 | 10.4 | 10.41 |
| 50-54 | 1 | 7.4 | 7.36 | 1 | 15.8 | 15.76 | 0 | 0.0 | 0.00 |
| 55-59 | 3 | 28.4 | 16.41 | 1 | 20.3 | 20.31 | 2 | 35.5 | 25.11 |
| 60-64 | 1 | 12.4 | 12.37 | 1 | 26.7 | 26.73 | 0 | 0.0 | 0.00 |
| 65-69 | 1 | 16.7 | 16.74 | 1 | 39.0 | 38.96 | 0 | 0.0 | 0.00 |
| 70-74 | 3 | 66.8 | 38.58 | 1 | 54.6 | 54.61 | 2 | 75.2 | 53.21 |
| 75-79 | 3 | 90.2 | 52.06 | 2 | 156.3 | 110.49 | 1 | 48.9 | 48.85 |
| 80-84 | 3 | 165.4 | 95.48 | 0 | 0.0 | 0.00 | 3 | 256.2 | 147.91 |
| 85+ | 4 | 290.7 | 145.35 | 1 | 214.6 | 214.59 | 3 | 329.7 | 190.34 |
| Total deaths | 25 | | | 11 | | | 14 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 6.0 | - | | 5.4 | - | | 6.6 | - |
| . AAMR (1940 Std. mil.) | | 8.1 | 1.70 | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | 11.6 | 2.46 | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | 16.4 | 3.64 | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | 20.0 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 19.5 | - | | 26.9 | - | | 14.8 | - |
| . HP 2000 (1940 Std. mil.) | | 25.0 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----COPD-related (490-496)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | 1 | 4.6 | 4.56 | 0 | 0.0 | 0.00 |
| 30-34 | 1 | 2.6 | 2.63 | 1 | 5.4 | 5.43 | 0 | 0.0 | 0.00 |
| 35-39 | 2 | 6.7 | 4.76 | 0 | 0.0 | 0.00 | 2 | 13.0 | 9.17 |
| 40-44 | 1 | 4.1 | 4.08 | 0 | 0.0 | 0.00 | 1 | 7.8 | 7.77 |
| 45-49 | 1 | 5.5 | 5.51 | 0 | 0.0 | 0.00 | 1 | 10.4 | 10.41 |
| 50-54 | 5 | 36.8 | 16.46 | 3 | 47.3 | 27.30 | 2 | 27.6 | 19.52 |
| 55-59 | 6 | 56.8 | 23.20 | 2 | 40.6 | 28.72 | 4 | 71.0 | 35.51 |
| 60-64 | 1 | 12.4 | 12.37 | 1 | 26.7 | 26.73 | 0 | 0.0 | 0.00 |
| 65-69 | 6 | 100.4 | 41.00 | 5 | 194.8 | 87.11 | 1 | 29.4 | 29.35 |
| 70-74 | 7 | 155.9 | 58.94 | 2 | 109.2 | 77.24 | 5 | 188.1 | 84.13 |
| 75-79 | 11 | 330.6 | 99.69 | 6 | 468.8 | 191.37 | 5 | 244.3 | 109.24 |
| 80-84 | 7 | 385.9 | 145.85 | 3 | 466.6 | 269.37 | 4 | 341.6 | 170.79 |
| 85+ | 7 | 508.7 | 192.28 | 3 | 643.8 | 371.68 | 4 | 439.6 | 219.78 |
| Total deaths | 57 | | | 28 | | | 29 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 13.6 | - | | 13.6 | - | | 13.6 | - |
| . AAMR (1940 Std. mil.) | | 20.0 | 2.74 | | 24.1 | 4.70 | | 17.3 | 3.34 |
| . AAMR (1970 Std. mil.) | | 28.7 | 3.93 | | 35.2 | 6.99 | | 24.5 | 4.66 |
| . AAMR (2000 Std. mil.) | | 39.4 | 5.61 | | 48.0 | 10.21 | | 34.1 | 6.61 |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Asthma (493)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | 1 | 4.6 | 4.56 | 0 | 0.0 | 0.00 |
| 30-34 | 1 | 2.6 | 2.63 | 1 | 5.4 | 5.43 | 0 | 0.0 | 0.00 |
| 35-39 | 1 | 3.4 | 3.36 | 0 | 0.0 | 0.00 | 1 | 6.5 | 6.49 |
| 40-44 | 1 | 4.1 | 4.08 | 0 | 0.0 | 0.00 | 1 | 7.8 | 7.77 |
| 45-49 | 1 | 5.5 | 5.51 | 0 | 0.0 | 0.00 | 1 | 10.4 | 10.41 |
| 50-54 | 1 | 7.4 | 7.36 | 1 | 15.8 | 15.76 | 0 | 0.0 | 0.00 |
| 55-59 | 1 | 9.5 | 9.47 | 0 | 0.0 | 0.00 | 1 | 17.8 | 17.75 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 1 | 55.1 | 55.13 | 0 | 0.0 | 0.00 | 1 | 85.4 | 85.40 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 9 | | | 4 | | | 5 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 2.1 | - | | 1.9 | - | | 2.3 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 1.4 | - | | 1.3 | - | | 1.6 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic liver disease and cirrhosis (571)-----

| Age | Both Sexes | | | Males | | | Females | | |
|--|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|----------------------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 1 | 2.6 | 2.63 | 0 | 0.0 | 0.00 | 1 | 5.1 | 5.12 |
| 35-39 | 4 | 13.5 | 6.73 | 3 | 21.0 | 12.10 | 1 | 6.5 | 6.49 |
| 40-44 | 3 | 12.3 | 7.07 | 3 | 25.8 | 14.90 | 0 | 0.0 | 0.00 |
| 45-49 | 6 | 33.1 | 13.50 | 6 | 70.2 | 28.68 | 0 | 0.0 | 0.00 |
| 50-54 | 11 | 81.0 | 24.41 | 11 | 173.4 | 52.28 | 0 | 0.0 | 0.00 |
| 55-59 | 2 | 18.9 | 13.40 | 2 | 40.6 | 28.72 | 0 | 0.0 | 0.00 |
| 60-64 | 5 | 61.9 | 27.66 | 4 | 106.9 | 53.46 | 1 | 23.0 | 23.03 |
| 65-69 | 4 | 67.0 | 33.48 | 4 | 155.8 | 77.91 | 0 | 0.0 | 0.00 |
| 70-74 | 2 | 44.6 | 31.50 | 1 | 54.6 | 54.61 | 1 | 37.6 | 37.62 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 1 | 55.1 | 55.13 | 0 | 0.0 | 0.00 | 1 | 85.4 | 85.40 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 39 | | | 34 | | | 5 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 9.3 | - | | 16.5 | - | | 2.3 | - |
| . AAMR (1940 Std. mil.) | | 14.7 | 2.40 | | 28.4 | 4.97 | | - | - |
| . AAMR (1970 Std. mil.) | | 15.7 | 2.64 | | 29.7 | 5.36 | | - | - |
| . AAMR (2000 Std. mil.) | | 17.7 | 2.97 | | 32.9 | 5.82 | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 9.2 | - | | 12.8 | - | | 5.5 | - |
| . HP 2000 (1940 Std. mil.) | | 6.0 | - | | - | - | | - | - |
| Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs: | | | | | | | | | |
| . The CT HP 2000 target: | | | | | | | | | No comparison is available |
| . The U.S. AAMR (1940 Std. mil.): | | | | | | | | | CT rate is higher |
| . THE U.S. HP 2000 target: | | | | | | | | | CT rate is higher |

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis, nephrotic syndrome/nephrosis (580-589)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 1 | 2.2 | 2.22 | 1 | 4.5 | 4.46 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 1 | 2.6 | 2.63 | 1 | 5.4 | 5.43 | 0 | 0.0 | 0.00 |
| 35-39 | 2 | 6.7 | 4.76 | 0 | 0.0 | 0.00 | 2 | 13.0 | 9.17 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 45-49 | 2 | 11.0 | 7.79 | 1 | 11.7 | 11.71 | 1 | 10.4 | 10.41 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 1 | 9.5 | 9.47 | 0 | 0.0 | 0.00 | 1 | 17.8 | 17.75 |
| 60-64 | 2 | 24.7 | 17.50 | 2 | 53.5 | 37.80 | 0 | 0.0 | 0.00 |
| 65-69 | 3 | 50.2 | 28.99 | 2 | 77.9 | 55.09 | 1 | 29.4 | 29.35 |
| 70-74 | 3 | 66.8 | 38.58 | 1 | 54.6 | 54.61 | 2 | 75.2 | 53.21 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 1 | 55.1 | 55.13 | 1 | 155.5 | 155.52 | 0 | 0.0 | 0.00 |
| 85+ | 1 | 72.7 | 72.67 | 1 | 214.6 | 214.59 | 0 | 0.0 | 0.00 |
| Total deaths | 18 | | | 11 | | | 7 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 4.3 | - | | 5.4 | - | | 3.3 | - |
| . AAMR (1940 Std. mil.) | | 6.3 | 1.56 | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | 7.7 | 1.97 | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | 9.2 | 2.44 | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 4.5 | - | | 5.7 | - | | 3.7 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis-related (580-589)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 2 | 4.1 | 2.89 | 1 | 4.0 | 3.99 | 1 | 4.2 | 4.19 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 1 | 2.2 | 2.22 | 1 | 4.5 | 4.46 | 0 | 0.0 | 0.00 |
| 25-29 | 1 | 2.2 | 2.23 | 0 | 0.0 | 0.00 | 1 | 4.3 | 4.35 |
| 30-34 | 3 | 7.9 | 4.56 | 3 | 16.3 | 9.40 | 0 | 0.0 | 0.00 |
| 35-39 | 6 | 20.2 | 8.24 | 4 | 27.9 | 13.97 | 2 | 13.0 | 9.17 |
| 40-44 | 1 | 4.1 | 4.08 | 1 | 8.6 | 8.60 | 0 | 0.0 | 0.00 |
| 45-49 | 7 | 38.6 | 14.58 | 5 | 58.5 | 26.18 | 2 | 20.8 | 14.73 |
| 50-54 | 7 | 51.5 | 19.47 | 5 | 78.8 | 35.25 | 2 | 27.6 | 19.52 |
| 55-59 | 9 | 85.3 | 28.42 | 2 | 40.6 | 28.72 | 7 | 124.3 | 46.97 |
| 60-64 | 9 | 111.3 | 37.11 | 4 | 106.9 | 53.46 | 5 | 115.2 | 51.50 |
| 65-69 | 14 | 234.3 | 62.63 | 9 | 350.6 | 116.87 | 5 | 146.8 | 65.63 |
| 70-74 | 8 | 178.2 | 63.01 | 3 | 163.8 | 94.60 | 5 | 188.1 | 84.13 |
| 75-79 | 4 | 120.2 | 60.11 | 1 | 78.1 | 78.13 | 3 | 146.6 | 84.61 |
| 80-84 | 6 | 330.8 | 135.03 | 4 | 622.1 | 311.04 | 2 | 170.8 | 120.77 |
| 85+ | 5 | 363.4 | 162.50 | 2 | 429.2 | 303.48 | 3 | 329.7 | 190.34 |
| Total deaths | 83 | | | 45 | | | 38 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 19.8 | - | | 21.9 | - | | 17.8 | - |
| . AAMR (1940 Std. mil.) | | 30.6 | 3.48 | | 37.2 | 5.77 | | 25.6 | 4.30 |
| . AAMR (1970 Std. mil.) | | 38.0 | 4.39 | | 46.1 | 7.53 | | 32.3 | 5.39 |
| . AAMR (2000 Std. mil.) | | 46.0 | 5.54 | | 57.1 | 9.98 | | 38.5 | 6.58 |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----End stage renal disease (ESRD) (581-588,591,593.8,593.9)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 1 | 2.2 | 2.22 | 1 | 4.5 | 4.46 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 1 | 2.6 | 2.63 | 1 | 5.4 | 5.43 | 0 | 0.0 | 0.00 |
| 35-39 | 2 | 6.7 | 4.76 | 0 | 0.0 | 0.00 | 2 | 13.0 | 9.17 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 45-49 | 2 | 11.0 | 7.79 | 1 | 11.7 | 11.71 | 1 | 10.4 | 10.41 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 2 | 18.9 | 13.40 | 0 | 0.0 | 0.00 | 2 | 35.5 | 25.11 |
| 60-64 | 2 | 24.7 | 17.50 | 2 | 53.5 | 37.80 | 0 | 0.0 | 0.00 |
| 65-69 | 4 | 67.0 | 33.48 | 2 | 77.9 | 55.09 | 2 | 58.7 | 41.51 |
| 70-74 | 3 | 66.8 | 38.58 | 1 | 54.6 | 54.61 | 2 | 75.2 | 53.21 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 1 | 55.1 | 55.13 | 1 | 155.5 | 155.52 | 0 | 0.0 | 0.00 |
| 85+ | 1 | 72.7 | 72.67 | 1 | 214.6 | 214.59 | 0 | 0.0 | 0.00 |
| Total deaths | 20 | | | 11 | | | 9 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 4.8 | - | | 5.4 | - | | 4.2 | - |
| . AAMR (1940 Std. mil.) | | 7.2 | 1.69 | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | 8.8 | 2.10 | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | 10.2 | 2.55 | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 5.1 | - | | 6.5 | - | | 4.3 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 3 | 6.1 | 3.54 | 2 | 8.0 | 5.64 | 1 | 4.2 | 4.19 |
| 5-9 | 2 | 4.8 | 3.36 | 1 | 4.7 | 4.68 | 1 | 4.8 | 4.84 |
| 10-14 | 2 | 5.0 | 3.57 | 2 | 9.9 | 7.03 | 0 | 0.0 | 0.00 |
| 15-19 | 7 | 18.1 | 6.82 | 6 | 30.2 | 12.34 | 1 | 5.3 | 5.29 |
| 20-24 | 11 | 24.4 | 7.37 | 9 | 40.2 | 13.39 | 2 | 8.9 | 6.26 |
| 25-29 | 8 | 17.8 | 6.30 | 8 | 36.5 | 12.90 | 0 | 0.0 | 0.00 |
| 30-34 | 7 | 18.4 | 6.97 | 7 | 38.0 | 14.35 | 0 | 0.0 | 0.00 |
| 35-39 | 3 | 10.1 | 5.83 | 3 | 21.0 | 12.10 | 0 | 0.0 | 0.00 |
| 40-44 | 4 | 16.3 | 8.17 | 4 | 34.4 | 17.21 | 0 | 0.0 | 0.00 |
| 45-49 | 4 | 22.0 | 11.02 | 3 | 35.1 | 20.28 | 1 | 10.4 | 10.41 |
| 50-54 | 4 | 29.4 | 14.72 | 4 | 63.1 | 31.53 | 0 | 0.0 | 0.00 |
| 55-59 | 1 | 9.5 | 9.47 | 1 | 20.3 | 20.31 | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | 0 | 0.0 | 0.00 | 1 | 23.0 | 23.03 |
| 65-69 | 2 | 33.5 | 23.67 | 1 | 39.0 | 38.96 | 1 | 29.4 | 29.35 |
| 70-74 | 1 | 22.3 | 22.28 | 1 | 54.6 | 54.61 | 0 | 0.0 | 0.00 |
| 75-79 | 1 | 30.1 | 30.06 | 0 | 0.0 | 0.00 | 1 | 48.9 | 48.85 |
| 80-84 | 1 | 55.1 | 55.13 | 0 | 0.0 | 0.00 | 1 | 85.4 | 85.40 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 62 | | | 52 | | | 10 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 14.8 | - | | 25.3 | - | | 4.7 | - |
| . AAMR (1940 Std. mil.) | | 15.9 | 2.13 | | 26.9 | 3.98 | | - | - |
| . AAMR (1970 Std. mil.) | | 15.7 | 2.25 | | 25.3 | 3.96 | | - | - |
| . AAMR (2000 Std. mil.) | | 16.7 | 2.54 | | 26.6 | 4.31 | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 33.8 | - | | 49.4 | - | | 18.9 | - |
| . HP 2000 (1940 Std. mil.) | | 29.3 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Motor vehicle accidents (E810-E825)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 1 | 2.4 | 2.38 | 0 | 0.0 | 0.00 | 1 | 4.8 | 4.84 |
| 10-14 | 2 | 5.0 | 3.57 | 2 | 9.9 | 7.03 | 0 | 0.0 | 0.00 |
| 15-19 | 7 | 18.1 | 6.82 | 6 | 30.2 | 12.34 | 1 | 5.3 | 5.29 |
| 20-24 | 9 | 20.0 | 6.67 | 8 | 35.7 | 12.62 | 1 | 4.4 | 4.43 |
| 25-29 | 5 | 11.1 | 4.98 | 5 | 22.8 | 10.20 | 0 | 0.0 | 0.00 |
| 30-34 | 4 | 10.5 | 5.27 | 4 | 21.7 | 10.85 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 3 | 12.3 | 7.07 | 3 | 25.8 | 14.90 | 0 | 0.0 | 0.00 |
| 45-49 | 1 | 5.5 | 5.51 | 1 | 11.7 | 11.71 | 0 | 0.0 | 0.00 |
| 50-54 | 1 | 7.4 | 7.36 | 1 | 15.8 | 15.76 | 0 | 0.0 | 0.00 |
| 55-59 | 1 | 9.5 | 9.47 | 1 | 20.3 | 20.31 | 0 | 0.0 | 0.00 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 65-69 | 1 | 16.7 | 16.74 | 0 | 0.0 | 0.00 | 1 | 29.4 | 29.35 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 36 | | | 32 | | | 4 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 8.6 | - | | 15.6 | - | | 1.9 | - |
| . AAMR (1940 Std. mil.) | | 8.5 | 1.47 | | 15.0 | 2.76 | | - | - |
| . AAMR (1970 Std. mil.) | | 8.1 | 1.45 | | 13.9 | 2.61 | | - | - |
| . AAMR (2000 Std. mil.) | | 7.7 | 1.43 | | 13.5 | 2.65 | | - | - |
| . HP 2000 (1940 Std. mil.) | | 10.8 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 18.8 | - | | 26.7 | - | | 11.3 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 45-49 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | 0 | 0.0 | 0.00 | 1 | 23.0 | 23.03 |
| 65-69 | 1 | 16.7 | 16.74 | 1 | 39.0 | 38.96 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 1 | 30.1 | 30.06 | 0 | 0.0 | 0.00 | 1 | 48.9 | 48.85 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 4 | | | 2 | | | 2 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 1.0 | - | | 1.0 | - | | 0.9 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | 2.3 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 2.7 | - | | 3.6 | - | | 1.6 | - |
| . HP 2000 (1940 Std. mil.) | | 2.3 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Drowning (E830,E832,E910)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 5-9 | 1 | 2.4 | 2.38 | 1 | 4.7 | 4.68 | 0 | - | - |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 30-34 | 1 | 2.6 | 2.63 | 1 | 5.4 | 5.43 | 0 | - | - |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 45-49 | 2 | 11.0 | 7.79 | 2 | 23.4 | 16.56 | 0 | - | - |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| Total deaths | 4 | | | 4 | | | - | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 1.0 | - | | 1.9 | - | | - | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | 1.0 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 1.9 | - | | 3.2 | - | | 0.6 | - |
| . HP 2000 (1940 Std. mil.) | | 1.3 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Residential fires (E890-E899)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 0 | - | - | 1 | 4.2 | 4.19 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 45-49 | 1 | 5.5 | 5.51 | 0 | - | - | 1 | 10.4 | 10.41 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | - | - | 0 | 0.0 | 0.00 |
| Total deaths | 2 | | | - | | | 2 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 0.5 | - | | - | - | | 0.9 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | 0.5 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 1.7 | - | | 2.0 | - | | 1.1 | - |
| . HP 2000 (1940 Std. mil.) | | 1.2 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): No comparison is available
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 1 | 2.5 | 2.52 | 1 | 5.0 | 4.97 | 0 | 0.0 | 0.00 |
| 15-19 | 2 | 5.2 | 3.65 | 1 | 5.0 | 5.04 | 1 | 5.3 | 5.29 |
| 20-24 | 4 | 8.9 | 4.45 | 4 | 17.9 | 8.93 | 0 | 0.0 | 0.00 |
| 25-29 | 7 | 15.6 | 5.89 | 5 | 22.8 | 10.20 | 2 | 8.7 | 6.15 |
| 30-34 | 4 | 10.5 | 5.27 | 3 | 16.3 | 9.40 | 1 | 5.1 | 5.12 |
| 35-39 | 7 | 23.5 | 8.90 | 7 | 48.9 | 18.48 | 0 | 0.0 | 0.00 |
| 40-44 | 5 | 20.4 | 9.13 | 5 | 43.0 | 19.24 | 0 | 0.0 | 0.00 |
| 45-49 | 4 | 22.0 | 11.02 | 4 | 46.8 | 23.42 | 0 | 0.0 | 0.00 |
| 50-54 | 1 | 7.4 | 7.36 | 1 | 15.8 | 15.76 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | 1 | 26.7 | 26.73 | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 70-74 | 1 | 22.3 | 22.28 | 1 | 54.6 | 54.61 | 0 | 0.0 | 0.00 |
| 75-79 | 1 | 30.1 | 30.06 | 1 | 78.1 | 78.13 | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 38 | | | 34 | | | 4 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 9.1 | - | | 16.5 | - | | 1.9 | - |
| . AAMR (1940 Std. mil.) | | 9.7 | 1.63 | | 18.8 | 3.40 | | - | - |
| . AAMR (1970 Std. mil.) | | 9.0 | 1.62 | | 17.8 | 3.51 | | - | - |
| . AAMR (2000 Std. mil.) | | 10.5 | 1.95 | | 21.3 | 4.31 | | - | - |
| . HP 2000 (1940 Std. mil.) | | 6.7 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 11.4 | - | | 18.8 | - | | 4.6 | - |
| . HP 2000 (1940 Std. mil.) | | 10.5 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide and legal intervention (E960-E978)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 2 | 4.1 | 2.89 | 2 | 8.0 | 5.64 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 1 | 2.5 | 2.52 | 0 | 0.0 | 0.00 | 1 | 5.1 | 5.12 |
| 15-19 | 11 | 28.4 | 8.56 | 10 | 50.4 | 15.93 | 1 | 5.3 | 5.29 |
| 20-24 | 8 | 17.8 | 6.29 | 8 | 35.7 | 12.62 | 0 | 0.0 | 0.00 |
| 25-29 | 11 | 24.5 | 7.38 | 9 | 41.1 | 13.68 | 2 | 8.7 | 6.15 |
| 30-34 | 6 | 15.8 | 6.45 | 3 | 16.3 | 9.40 | 3 | 15.4 | 8.87 |
| 35-39 | 3 | 10.1 | 5.83 | 2 | 14.0 | 9.88 | 1 | 6.5 | 6.49 |
| 40-44 | 2 | 8.2 | 5.78 | 2 | 17.2 | 12.17 | 0 | 0.0 | 0.00 |
| 45-49 | 4 | 22.0 | 11.02 | 3 | 35.1 | 20.28 | 1 | 10.4 | 10.41 |
| 50-54 | 1 | 7.4 | 7.36 | 1 | 15.8 | 15.76 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 1 | 55.1 | 55.13 | 1 | 155.5 | 155.52 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 50 | | | 41 | | | 9 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 11.9 | - | | 20.0 | - | | 4.2 | - |
| . AAMR (1940 Std. mil.) | | 11.5 | 1.66 | | 19.4 | 3.15 | | - | - |
| . AAMR (1970 Std. mil.) | | 10.6 | 1.61 | | 18.4 | 3.27 | | - | - |
| . AAMR (2000 Std. mil.) | | 10.9 | 1.82 | | 19.2 | 3.98 | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 9.4 | - | | 14.6 | - | | 4.2 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide (E960-E969)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|------------------|---------------------|---------------|------------------|---------------------|---------------|------------------|---------------------|---------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 2 | 4.1 | 2.89 | 2 | 8.0 | 5.64 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 1 | 2.5 | 2.52 | 0 | 0.0 | 0.00 | 1 | 5.1 | 5.12 |
| 15-19 | 11 | 28.4 | 8.56 | 10 | 50.4 | 15.93 | 1 | 5.3 | 5.29 |
| 20-24 | 8 | 17.8 | 6.29 | 8 | 35.7 | 12.62 | 0 | 0.0 | 0.00 |
| 25-29 | 11 | 24.5 | 7.38 | 9 | 41.1 | 13.68 | 2 | 8.7 | 6.15 |
| 30-34 | 6 | 15.8 | 6.45 | 3 | 16.3 | 9.40 | 3 | 15.4 | 8.87 |
| 35-39 | 3 | 10.1 | 5.83 | 2 | 14.0 | 9.88 | 1 | 6.5 | 6.49 |
| 40-44 | 2 | 8.2 | 5.78 | 2 | 17.2 | 12.17 | 0 | 0.0 | 0.00 |
| 45-49 | 4 | 22.0 | 11.02 | 3 | 35.1 | 20.28 | 1 | 10.4 | 10.41 |
| 50-54 | 1 | 7.4 | 7.36 | 1 | 15.8 | 15.76 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 1 | 55.1 | 55.13 | 1 | 155.5 | 155.52 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 50 | | | 41 | | | 9 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 11.9 | - | | 20.0 | - | | 4.2 | - |
| . AAMR (1940 Std. mil.) | | 11.5 | 1.66 | | 19.4 | 3.15 | | - | - |
| . AAMR (1970 Std. mil.) | | 10.6 | 1.61 | | 18.4 | 3.27 | | - | - |
| . AAMR (2000 Std. mil.) | | 10.9 | 1.82 | | 19.2 | 3.98 | | - | - |
| . HP 2000 (1940 Std. mil.) | | 5.0 | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 9.3 | - | | 14.3 | - | | 4.2 | - |
| . HP 2000 (1940 Std. mil.) | | 7.2 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
- . THE U.S. HP 2000 target: CT rate is higher

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All injury (E800-E999)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 5 | 10.2 | 4.57 | 4 | 16.0 | 7.98 | 1 | 4.2 | 4.19 |
| 5-9 | 2 | 4.8 | 3.36 | 1 | 4.7 | 4.68 | 1 | 4.8 | 4.84 |
| 10-14 | 4 | 10.1 | 5.05 | 3 | 14.9 | 8.62 | 1 | 5.1 | 5.12 |
| 15-19 | 20 | 51.6 | 11.54 | 17 | 85.6 | 20.77 | 3 | 15.9 | 9.16 |
| 20-24 | 24 | 53.3 | 10.89 | 22 | 98.2 | 20.93 | 2 | 8.9 | 6.26 |
| 25-29 | 26 | 57.9 | 11.35 | 22 | 100.4 | 21.40 | 4 | 17.4 | 8.69 |
| 30-34 | 18 | 47.4 | 11.17 | 14 | 76.0 | 20.30 | 4 | 20.5 | 10.24 |
| 35-39 | 14 | 47.1 | 12.58 | 13 | 90.8 | 25.18 | 1 | 6.5 | 6.49 |
| 40-44 | 11 | 44.9 | 13.54 | 11 | 94.6 | 28.54 | 0 | 0.0 | 0.00 |
| 45-49 | 13 | 71.6 | 19.87 | 11 | 128.8 | 38.83 | 2 | 20.8 | 14.73 |
| 50-54 | 6 | 44.2 | 18.03 | 6 | 94.6 | 38.61 | 0 | 0.0 | 0.00 |
| 55-59 | 1 | 9.5 | 9.47 | 1 | 20.3 | 20.31 | 0 | 0.0 | 0.00 |
| 60-64 | 2 | 24.7 | 17.50 | 1 | 26.7 | 26.73 | 1 | 23.0 | 23.03 |
| 65-69 | 2 | 33.5 | 23.67 | 1 | 39.0 | 38.96 | 1 | 29.4 | 29.35 |
| 70-74 | 2 | 44.6 | 31.50 | 2 | 109.2 | 77.24 | 0 | 0.0 | 0.00 |
| 75-79 | 2 | 60.1 | 42.51 | 1 | 78.1 | 78.13 | 1 | 48.9 | 48.85 |
| 80-84 | 2 | 110.3 | 77.96 | 1 | 155.5 | 155.52 | 1 | 85.4 | 85.40 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 154 | | | 131 | | | 23 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 36.8 | - | | 63.8 | - | | 10.8 | - |
| . AAMR (1940 Std. mil.) | | 38.0 | 3.20 | | 67.1 | 6.19 | | 11.0 | 2.39 |
| . AAMR (1970 Std. mil.) | | 36.1 | 3.24 | | 63.2 | 6.29 | | 11.3 | 2.63 |
| . AAMR (2000 Std. mil.) | | 39.1 | 3.72 | | 69.1 | 7.37 | | 12.1 | 3.01 |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 55.6 | - | | 84.4 | - | | 28.1 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 1 | 2.4 | 2.38 | 0 | 0.0 | 0.00 | 1 | 4.8 | 4.84 |
| 10-14 | 2 | 5.0 | 3.57 | 2 | 9.9 | 7.03 | 0 | 0.0 | 0.00 |
| 15-19 | 7 | 18.1 | 6.82 | 6 | 30.2 | 12.34 | 1 | 5.3 | 5.29 |
| 20-24 | 9 | 20.0 | 6.67 | 8 | 35.7 | 12.62 | 1 | 4.4 | 4.43 |
| 25-29 | 5 | 11.1 | 4.98 | 5 | 22.8 | 10.20 | 0 | 0.0 | 0.00 |
| 30-34 | 3 | 7.9 | 4.56 | 3 | 16.3 | 9.40 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 3 | 12.3 | 7.07 | 3 | 25.8 | 14.90 | 0 | 0.0 | 0.00 |
| 45-49 | 1 | 5.5 | 5.51 | 1 | 11.7 | 11.71 | 0 | 0.0 | 0.00 |
| 50-54 | 1 | 7.4 | 7.36 | 1 | 15.8 | 15.76 | 0 | 0.0 | 0.00 |
| 55-59 | 1 | 9.5 | 9.47 | 1 | 20.3 | 20.31 | 0 | 0.0 | 0.00 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 65-69 | 1 | 16.7 | 16.74 | 0 | 0.0 | 0.00 | 1 | 29.4 | 29.35 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 35 | | | 31 | | | 4 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 8.4 | - | | 15.1 | - | | 1.9 | - |
| . AAMR (1940 Std. mil.) | | 8.3 | 1.46 | | 14.6 | 2.73 | | - | - |
| . AAMR (1970 Std. mil.) | | 7.9 | 1.44 | | 13.6 | 2.60 | | - | - |
| . AAMR (2000 Std. mil.) | | 7.5 | 1.42 | | 13.1 | 2.62 | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 18.6 | - | | 26.1 | - | | 10.9 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 1 | 2.0 | 2.04 | 1 | 4.0 | 3.99 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 35-39 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 40-44 | 1 | 4.1 | 4.08 | 1 | 8.6 | 8.60 | 0 | 0.0 | 0.00 |
| 45-49 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | 0 | 0.0 | 0.00 | 1 | 23.0 | 23.03 |
| 65-69 | 1 | 16.7 | 16.74 | 1 | 39.0 | 38.96 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 1 | 30.1 | 30.06 | 0 | 0.0 | 0.00 | 1 | 48.9 | 48.85 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 5 | | | 3 | | | 2 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 1.2 | - | | 1.5 | - | | 0.9 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 2.6 | - | | 3.6 | - | | 1.4 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 2 | 5.0 | 3.57 | 1 | 5.0 | 4.97 | 1 | 5.1 | 5.12 |
| 15-19 | 12 | 31.0 | 8.94 | 10 | 50.4 | 15.93 | 2 | 10.6 | 7.48 |
| 20-24 | 9 | 20.0 | 6.67 | 9 | 40.2 | 13.39 | 0 | 0.0 | 0.00 |
| 25-29 | 14 | 31.2 | 8.33 | 10 | 45.6 | 14.43 | 4 | 17.4 | 8.69 |
| 30-34 | 5 | 13.2 | 5.89 | 3 | 16.3 | 9.40 | 2 | 10.2 | 7.24 |
| 35-39 | 3 | 10.1 | 5.83 | 2 | 14.0 | 9.88 | 1 | 6.5 | 6.49 |
| 40-44 | 1 | 4.1 | 4.08 | 1 | 8.6 | 8.60 | 0 | 0.0 | 0.00 |
| 45-49 | 3 | 16.5 | 9.55 | 3 | 35.1 | 20.28 | 0 | 0.0 | 0.00 |
| 50-54 | 1 | 7.4 | 7.36 | 1 | 15.8 | 15.76 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | 1 | 26.7 | 26.73 | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 70-74 | 1 | 22.3 | 22.28 | 1 | 54.6 | 54.61 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 1 | 55.1 | 55.13 | 1 | 155.5 | 155.52 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 53 | | | 43 | | | 10 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 12.7 | - | | 20.9 | - | | 4.7 | - |
| . AAMR (1940 Std. mil.) | | 12.4 | 1.77 | | 21.4 | 3.46 | | - | - |
| . AAMR (1970 Std. mil.) | | 11.8 | 1.79 | | 21.0 | 3.75 | | - | - |
| . AAMR (2000 Std. mil.) | | 11.9 | 1.97 | | 21.7 | 4.43 | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 13.6 | - | | 23.7 | - | | 4.1 | - |
| . HP 2000 (1940 Std. mil.) | | 11.6 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: No comparison is available
- . The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
- . THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm suicide (E955.0-.4)-----

| Age | Both Sexes | | | Males | | | Females | | |
|--|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|----------------------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 1 | 2.5 | 2.52 | 1 | 5.0 | 4.97 | 0 | 0.0 | 0.00 |
| 15-19 | 2 | 5.2 | 3.65 | 1 | 5.0 | 5.04 | 1 | 5.3 | 5.29 |
| 20-24 | 1 | 2.2 | 2.22 | 1 | 4.5 | 4.46 | 0 | 0.0 | 0.00 |
| 25-29 | 3 | 6.7 | 3.86 | 1 | 4.6 | 4.56 | 2 | 8.7 | 6.15 |
| 30-34 | 1 | 2.6 | 2.63 | 1 | 5.4 | 5.43 | 0 | 0.0 | 0.00 |
| 35-39 | 2 | 6.7 | 4.76 | 2 | 14.0 | 9.88 | 0 | 0.0 | 0.00 |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 45-49 | 2 | 11.0 | 7.79 | 2 | 23.4 | 16.56 | 0 | 0.0 | 0.00 |
| 50-54 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 1 | 12.4 | 12.37 | 1 | 26.7 | 26.73 | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 70-74 | 1 | 22.3 | 22.28 | 1 | 54.6 | 54.61 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 14 | | | 11 | | | 3 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 3.3 | - | | 5.4 | - | | 1.4 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 6.8 | - | | 12.0 | - | | 1.8 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs: | | | | | | | | | |
| . The CT HP 2000 target: | | | | | | | | | No comparison is available |
| . The U.S. AAMR (1940 Std. mil.): | | | | | | | | | No comparison is available |
| . THE U.S. HP 2000 target: | | | | | | | | | No comparison is available |

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 1 | 2.5 | 2.52 | 0 | 0.0 | 0.00 | 1 | 5.1 | 5.12 |
| 15-19 | 10 | 25.8 | 8.16 | 9 | 45.3 | 15.11 | 1 | 5.3 | 5.29 |
| 20-24 | 7 | 15.6 | 5.88 | 7 | 31.2 | 11.81 | 0 | 0.0 | 0.00 |
| 25-29 | 10 | 22.3 | 7.04 | 8 | 36.5 | 12.90 | 2 | 8.7 | 6.15 |
| 30-34 | 4 | 10.5 | 5.27 | 2 | 10.9 | 7.67 | 2 | 10.2 | 7.24 |
| 35-39 | 1 | 3.4 | 3.36 | 0 | 0.0 | 0.00 | 1 | 6.5 | 6.49 |
| 40-44 | 1 | 4.1 | 4.08 | 1 | 8.6 | 8.60 | 0 | 0.0 | 0.00 |
| 45-49 | 1 | 5.5 | 5.51 | 1 | 11.7 | 11.71 | 0 | 0.0 | 0.00 |
| 50-54 | 1 | 7.4 | 7.36 | 1 | 15.8 | 15.76 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 1 | 55.1 | 55.13 | 1 | 155.5 | 155.52 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 37 | | | 30 | | | 7 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 8.8 | - | | 14.6 | - | | 3.3 | - |
| . AAMR (1940 Std. mil.) | | 8.3 | 1.39 | | 14.0 | 2.66 | | - | - |
| . AAMR (1970 Std. mil.) | | 7.8 | 1.39 | | 13.6 | 2.89 | | - | - |
| . AAMR (2000 Std. mil.) | | 7.7 | 1.55 | | 13.8 | 3.56 | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 6.1 | - | | 9.9 | - | | 1.9 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All poisoning (E850-E869, E950-E952, E962, E972, E980-E982)-----

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 1 | 2.2 | 2.22 | 1 | 4.5 | 4.46 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 3 | 7.9 | 4.56 | 2 | 10.9 | 7.67 | 1 | 5.1 | 5.12 |
| 35-39 | 1 | 3.4 | 3.36 | 1 | 7.0 | 6.98 | 0 | 0.0 | 0.00 |
| 40-44 | 1 | 4.1 | 4.08 | 1 | 8.6 | 8.60 | 0 | 0.0 | 0.00 |
| 45-49 | 1 | 5.5 | 5.51 | 1 | 11.7 | 11.71 | 0 | 0.0 | 0.00 |
| 50-54 | 1 | 7.4 | 7.36 | 1 | 15.8 | 15.76 | 0 | 0.0 | 0.00 |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 75-79 | 1 | 30.1 | 30.06 | 1 | 78.1 | 78.13 | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 9 | | | 8 | | | 1 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 2.1 | - | | 3.9 | - | | 0.5 | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 5.0 | - | | 6.9 | - | | 3.2 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
 U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
 for
 Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 25-29 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 30-34 | 2 | 5.3 | 3.72 | 0 | 0.0 | 0.00 | 2 | 10.2 | 7.24 |
| 35-39 | 3 | 10.1 | 5.83 | 2 | 14.0 | 9.88 | 1 | 6.5 | 6.49 |
| 40-44 | 2 | 8.2 | 5.78 | 2 | 17.2 | 12.17 | 0 | 0.0 | 0.00 |
| 45-49 | 7 | 38.6 | 14.58 | 7 | 82.0 | 30.98 | 0 | 0.0 | 0.00 |
| 50-54 | 7 | 51.5 | 19.47 | 7 | 110.3 | 41.70 | 0 | 0.0 | 0.00 |
| 55-59 | 4 | 37.9 | 18.94 | 4 | 81.2 | 40.62 | 0 | 0.0 | 0.00 |
| 60-64 | 3 | 37.1 | 21.43 | 3 | 80.2 | 46.30 | 0 | 0.0 | 0.00 |
| 65-69 | 4 | 67.0 | 33.48 | 4 | 155.8 | 77.91 | 0 | 0.0 | 0.00 |
| 70-74 | 1 | 22.3 | 22.28 | 1 | 54.6 | 54.61 | 0 | 0.0 | 0.00 |
| 75-79 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 |
| Total deaths | 33 | | | 30 | | | 3 | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 7.9 | - | | 14.6 | - | | 1.4 | - |
| . AAMR (1940 Std. mil.) | | 12.3 | 2.19 | | 25.4 | 4.73 | | - | - |
| . AAMR (1970 Std. mil.) | | 12.8 | 2.35 | | 26.9 | 5.15 | | - | - |
| . AAMR (2000 Std. mil.) | | 14.1 | 2.54 | | 29.5 | 5.54 | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 7.4 | - | | 11.6 | - | | 3.2 | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
 . The CT HP 2000 target: No comparison is available
 . The U.S. AAMR (1940 Std. mil.): CT rate is higher
 . THE U.S. HP 2000 target: No comparison is available

Connecticut Hispanic Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
 U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
 for
 Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

| Age | Both Sexes | | | Males | | | Females | | |
|----------------------------|---------------|------------------|------------|---------------|------------------|------------|---------------|------------------|------------|
| | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error | No. Of Deaths | Rate per 100,000 | Std. Error |
| 0-4 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 5-9 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 10-14 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 15-19 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 20-24 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 25-29 | 2 | 4.5 | 3.15 | 2 | 9.1 | 6.45 | 0 | - | - |
| 30-34 | 2 | 5.3 | 3.72 | 2 | 10.9 | 7.67 | 0 | - | - |
| 35-39 | 1 | 3.4 | 3.36 | 1 | 7.0 | 6.98 | 0 | - | - |
| 40-44 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 45-49 | 1 | 5.5 | 5.51 | 1 | 11.7 | 11.71 | 0 | - | - |
| 50-54 | 1 | 7.4 | 7.36 | 1 | 15.8 | 15.76 | 0 | - | - |
| 55-59 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 60-64 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 65-69 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 70-74 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 75-79 | 1 | 30.1 | 30.06 | 1 | 78.1 | 78.13 | 0 | - | - |
| 80-84 | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| 85+ | 0 | 0.0 | 0.00 | 0 | 0.0 | 0.00 | 0 | - | - |
| Total deaths | 8 | | | 8 | | | - | | |
| Connecticut | | | | | | | | | |
| . Crude rate | | 1.9 | - | | 3.9 | - | | - | - |
| . AAMR (1940 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (1970 Std. mil.) | | - | - | | - | - | | - | - |
| . AAMR (2000 Std. mil.) | | - | - | | - | - | | - | - |
| . HP 2000 (1940 Std. mil.) | | - | - | | - | - | | - | - |
| United States | | | | | | | | | |
| . AAMR (1940 Std. mil.) | | 4.1 | - | | 5.5 | - | | 2.7 | - |
| . HP 2000 (1940 Std. mil.) | | 3.0 | - | | - | - | | - | - |

Comparison of the CT Hispanic Resident AAMR (1940 Std. mil.) vs:
 . The CT HP 2000 target: No comparison is available
 . The U.S. AAMR (1940 Std. mil.): No comparison is available
 . THE U.S. HP 2000 target: No comparison is available