

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Infectious diseases and parasitic disease (1-139)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	18	4.5	1.07	10	4.9	1.55	8	4.2	1.47
5-9	5	1.4	0.63	2	1.1	0.78	3	1.7	1.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	7	1.9	0.71	5	2.6	1.17	2	1.1	0.78
20-24	12	2.7	0.79	10	4.5	1.42	2	0.9	0.65
25-29	48	9.5	1.38	37	14.6	2.40	11	4.4	1.32
30-34	81	15.3	1.70	67	25.4	3.10	14	5.3	1.41
35-39	70	14.8	1.76	58	24.6	3.23	12	5.0	1.45
40-44	60	13.6	1.75	47	21.7	3.16	13	5.8	1.60
45-49	49	13.6	1.95	40	22.6	3.58	9	4.9	1.64
50-54	51	18.0	2.51	38	27.4	4.44	13	9.0	2.48
55-59	37	13.6	2.23	26	19.7	3.86	11	7.8	2.36
60-64	42	14.9	2.30	28	21.0	3.96	14	9.4	2.52
65-69	83	31.4	3.44	44	37.0	5.58	39	26.8	4.29
70-74	87	39.4	4.22	51	54.6	7.65	36	28.2	4.70
75-79	149	89.6	7.34	67	104.0	12.70	82	80.6	8.90
80-84	156	148.4	11.88	68	193.5	23.46	88	125.8	13.41
85+	263	297.9	18.37	67	294.9	36.03	196	298.9	21.35
Total deaths	1,218			665			553		
Connecticut									
. Crude rate		20.7	-		23.3	-		18.2	-
. AAMR (1940 Std. mil.)		12.2	0.40		17.4	0.71		7.5	0.41
. AAMR (1970 Std. mil.)		14.8	0.44		19.9	0.78		10.3	0.48
. AAMR (2000 Std. mil.)		20.1	0.58		26.4	1.07		14.6	0.63
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		16.4	-		25.2	-		8.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Septicemia (038)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	1.5	0.62	3	1.5	0.85	3	1.6	0.90
5-9	1	0.3	0.28	0	0.0	0.00	1	0.6	0.58
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.27	1	0.5	0.52	0	0.0	0.00
20-24	3	0.7	0.39	3	1.3	0.78	0	0.0	0.00
25-29	7	1.4	0.53	5	2.0	0.88	2	0.8	0.56
30-34	5	0.9	0.42	3	1.1	0.66	2	0.8	0.53
35-39	6	1.3	0.52	2	0.8	0.60	4	1.7	0.84
40-44	10	2.3	0.71	7	3.2	1.22	3	1.3	0.77
45-49	12	3.3	0.96	6	3.4	1.39	6	3.3	1.34
50-54	16	5.6	1.41	8	5.8	2.04	8	5.5	1.95
55-59	15	5.5	1.42	10	7.6	2.40	5	3.6	1.59
60-64	27	9.6	1.84	15	11.2	2.90	12	8.1	2.33
65-69	59	22.3	2.90	31	26.1	4.68	28	19.2	3.63
70-74	68	30.8	3.73	39	41.8	6.69	29	22.7	4.22
75-79	128	77.0	6.81	57	88.4	11.72	71	69.8	8.28
80-84	133	126.5	10.97	55	156.5	21.10	78	111.5	12.62
85+	225	254.8	16.99	57	250.9	33.23	168	256.2	19.77
Total deaths	722			302			420		
Connecticut									
. Crude rate		12.3	-		10.6	-		13.8	-
. AAMR (1940 Std. mil.)		5.3	0.23		6.3	0.38		4.6	0.29
. AAMR (1970 Std. mil.)		8.1	0.31		9.4	0.54		7.2	0.38
. AAMR (2000 Std. mil.)		11.9	0.45		13.5	0.81		10.8	0.53
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.0	-		4.6	-		3.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is higher
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----HIV infection (042-044)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	1.0	0.50	2	1.0	0.69	2	1.0	0.73
5-9	1	0.3	0.28	0	0.0	0.00	1	0.6	0.58
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.27	0	0.0	0.00	1	0.5	0.55
20-24	8	1.8	0.64	6	2.7	1.10	2	0.9	0.65
25-29	38	7.5	1.22	30	11.9	2.16	8	3.2	1.13
30-34	72	13.6	1.60	61	23.1	2.96	11	4.2	1.25
35-39	57	12.0	1.59	50	21.2	3.00	7	2.9	1.11
40-44	42	9.5	1.46	37	17.1	2.80	5	2.2	0.99
45-49	30	8.3	1.52	29	16.4	3.05	1	0.5	0.55
50-54	29	10.2	1.90	25	18.0	3.60	4	2.8	1.38
55-59	9	3.3	1.10	9	6.8	2.27	0	0.0	0.00
60-64	7	2.5	0.94	6	4.5	1.84	1	0.7	0.67
65-69	6	2.3	0.93	4	3.4	1.68	2	1.4	0.97
70-74	1	0.5	0.45	1	1.1	1.07	0	0.0	0.00
75-79	1	0.6	0.60	0	0.0	0.00	1	1.0	0.98
80-84	1	1.0	0.95	1	2.8	2.85	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	307			261			46		
Connecticut									
. Crude rate		5.2	-		9.1	-		1.5	-
. AAMR (1940 Std. mil.)		4.9	0.28		8.4	0.52		1.5	0.22
. AAMR (1970 Std. mil.)		4.2	0.25		7.2	0.46		1.3	0.20
. AAMR (2000 Std. mil.)		5.1	0.29		8.8	0.55		1.4	0.22
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.7	-		15.7	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	1.0	0.50	1	0.5	0.49	3	1.6	0.90
5-9	15	4.2	1.09	8	4.4	1.55	7	4.0	1.53
10-14	4	1.2	0.60	2	1.2	0.83	2	1.2	0.87
15-19	8	2.1	0.76	5	2.6	1.17	3	1.6	0.95
20-24	20	4.5	1.01	10	4.5	1.42	10	4.6	1.45
25-29	43	8.5	1.30	22	8.7	1.85	21	8.4	1.83
30-34	82	15.5	1.71	41	15.5	2.43	41	15.5	2.42
35-39	131	27.6	2.41	56	23.7	3.17	75	31.4	3.63
40-44	217	49.0	3.33	83	38.3	4.20	134	59.4	5.13
45-49	357	99.2	5.25	176	99.6	7.50	181	98.9	7.35
50-54	494	173.9	7.82	243	175.0	11.22	251	172.8	10.91
55-59	880	322.7	10.88	439	332.7	15.88	441	313.4	14.92
60-64	1,424	504.9	13.38	767	574.7	20.75	657	442.2	17.25
65-69	1,936	731.7	16.63	1,067	897.6	27.48	869	596.4	20.23
70-74	2,173	983.4	21.10	1,181	1264.4	36.79	992	777.7	24.69
75-79	2,030	1221.2	27.10	1,062	1647.9	50.57	968	951.0	30.57
80-84	1,636	1556.3	38.48	806	2293.3	80.78	830	1186.2	41.17
85+	1,547	1752.1	44.55	612	2693.7	108.88	935	1425.8	46.63
Total deaths	13,001			6,581			6,420		
Connecticut									
. Crude rate		220.7	-		230.4	-		211.5	-
. AAMR (1940 Std. mil.)		124.8	1.21		148.9	1.92		108.9	1.57
. AAMR (1970 Std. mil.)		164.8	1.48		204.0	2.53		140.5	1.85
. AAMR (2000 Std. mil.)		206.3	1.83		262.3	3.37		174.4	2.21
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		134.3	-		165.4	-		112.2	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.2	0.20	1	0.4	0.40	0	0.0	0.00
30-34	6	1.1	0.46	1	0.4	0.38	5	1.9	0.84
35-39	7	1.5	0.56	6	2.5	1.04	1	0.4	0.42
40-44	21	4.7	1.04	8	3.7	1.30	13	5.8	1.60
45-49	29	8.1	1.50	15	8.5	2.19	14	7.7	2.04
50-54	38	13.4	2.17	19	13.7	3.14	19	13.1	3.00
55-59	68	24.9	3.02	34	25.8	4.42	34	24.2	4.14
60-64	148	52.5	4.31	89	66.7	7.07	59	39.7	5.17
65-69	209	79.0	5.46	117	98.4	9.10	92	63.1	6.58
70-74	277	125.4	7.53	155	165.9	13.33	122	95.6	8.66
75-79	285	171.4	10.16	143	221.9	18.56	142	139.5	11.71
80-84	244	232.1	14.86	106	301.6	29.29	138	197.2	16.79
85+	328	371.5	20.51	110	484.2	46.16	218	332.4	22.52
Total deaths	1,661			804			857		
Connecticut									
. Crude rate		28.2	-		28.1	-		28.2	-
. AAMR (1940 Std. mil.)		13.8	0.38		17.0	0.63		11.6	0.47
. AAMR (1970 Std. mil.)		19.8	0.50		24.8	0.88		16.6	0.60
. AAMR (2000 Std. mil.)		26.5	0.66		33.4	1.23		22.4	0.77
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		14.0	-		17.0	-		11.7	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colon cancer (153)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.2	0.20	1	0.4	0.40	0	0.0	0.00
30-34	6	1.1	0.46	1	0.4	0.38	5	1.9	0.84
35-39	5	1.1	0.47	4	1.7	0.85	1	0.4	0.42
40-44	19	4.3	0.98	7	3.2	1.22	12	5.3	1.54
45-49	23	6.4	1.33	10	5.7	1.79	13	7.1	1.97
50-54	32	11.3	1.99	14	10.1	2.69	18	12.4	2.92
55-59	60	22.0	2.84	31	23.5	4.22	29	20.6	3.83
60-64	122	43.3	3.92	73	54.7	6.40	49	33.0	4.71
65-69	178	67.3	5.04	100	84.1	8.41	78	53.5	6.06
70-74	223	100.9	6.76	122	130.6	11.83	101	79.2	7.88
75-79	240	144.4	9.32	120	186.2	17.00	120	117.9	10.76
80-84	205	195.0	13.62	89	253.2	26.84	116	165.8	15.39
85+	270	305.8	18.61	93	409.3	42.45	177	269.9	20.29
Total deaths	1,384			665			719		
Connecticut									
. Crude rate		23.5	-		23.3	-		23.7	-
. AAMR (1940 Std. mil.)		11.6	0.35		14.0	0.57		9.9	0.44
. AAMR (1970 Std. mil.)		16.5	0.45		20.5	0.80		14.0	0.55
. AAMR (2000 Std. mil.)		22.1	0.60		27.7	1.12		18.8	0.71
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.7	-		14.2	-		10.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Rectal cancer (154)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.4	0.30	2	0.8	0.60	0	0.0	0.00
40-44	2	0.5	0.32	1	0.5	0.46	1	0.4	0.44
45-49	5	1.4	0.62	4	2.3	1.13	1	0.5	0.55
50-54	6	2.1	0.86	5	3.6	1.61	1	0.7	0.69
55-59	8	2.9	1.04	3	2.3	1.31	5	3.6	1.59
60-64	22	7.8	1.66	13	9.7	2.70	9	6.1	2.02
65-69	31	11.7	2.10	17	14.3	3.47	14	9.6	2.57
70-74	45	20.4	3.04	27	28.9	5.56	18	14.1	3.33
75-79	40	24.1	3.80	19	29.5	6.76	21	20.6	4.50
80-84	34	32.3	5.55	14	39.8	10.65	20	28.6	6.39
85+	45	51.0	7.60	11	48.4	14.60	34	51.8	8.89
Total deaths	240			116			124		
Connecticut									
. Crude rate		4.1	-		4.1	-		4.1	-
. AAMR (1940 Std. mil.)		2.0	0.14		2.6	0.25		1.6	0.17
. AAMR (1970 Std. mil.)		2.9	0.19		3.6	0.33		2.3	0.22
. AAMR (2000 Std. mil.)		3.8	0.25		4.6	0.45		3.2	0.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.5	-		1.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.2	0.19	1	0.4	0.38	0	0.0	0.00
35-39	4	0.8	0.42	1	0.4	0.42	3	1.3	0.73
40-44	4	0.9	0.45	3	1.4	0.80	1	0.4	0.44
45-49	16	4.4	1.11	10	5.7	1.79	6	3.3	1.34
50-54	23	8.1	1.69	15	10.8	2.79	8	5.5	1.95
55-59	34	12.5	2.14	23	17.4	3.63	11	7.8	2.36
60-64	69	24.5	2.95	40	30.0	4.74	29	19.5	3.62
65-69	118	44.6	4.11	77	64.8	7.38	41	28.1	4.39
70-74	118	53.4	4.92	63	67.4	8.50	55	43.1	5.81
75-79	104	62.6	6.13	41	63.6	9.94	63	61.9	7.80
80-84	85	80.9	8.77	43	122.3	18.66	42	60.0	9.26
85+	96	108.7	11.10	25	110.0	22.01	71	108.3	12.85
Total deaths	672			342			330		
Connecticut									
. Crude rate		11.4	-		12.0	-		10.9	-
. AAMR (1940 Std. mil.)		6.1	0.26		7.9	0.44		4.7	0.30
. AAMR (1970 Std. mil.)		8.3	0.33		10.6	0.58		6.6	0.38
. AAMR (2000 Std. mil.)		10.6	0.41		13.2	0.74		8.6	0.48
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.4	-		7.5	-		5.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung and other respiratory cancer (160-165)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.27	1	0.5	0.52	0	0.0	0.00
20-24	1	0.2	0.23	0	0.0	0.00	1	0.5	0.46
25-29	1	0.2	0.20	1	0.4	0.40	0	0.0	0.00
30-34	3	0.6	0.33	2	0.8	0.54	1	0.4	0.38
35-39	12	2.5	0.73	8	3.4	1.20	4	1.7	0.84
40-44	32	7.2	1.28	20	9.2	2.06	12	5.3	1.54
45-49	107	29.7	2.88	68	38.5	4.66	39	21.3	3.41
50-54	155	54.6	4.38	91	65.5	6.87	64	44.1	5.51
55-59	311	114.0	6.47	175	132.6	10.02	136	96.6	8.29
60-64	463	164.2	7.63	287	215.0	12.69	176	118.5	8.93
65-69	615	232.4	9.37	359	302.0	15.94	256	175.7	10.98
70-74	654	296.0	11.57	400	428.2	21.41	254	199.1	12.49
75-79	539	324.2	13.97	337	522.9	28.49	202	198.5	13.96
80-84	326	310.1	17.18	209	594.7	41.13	117	167.2	15.46
85+	194	219.7	15.77	109	479.8	45.95	85	129.6	14.06
Total deaths	3,414			2,067			1,347		
Connecticut									
. Crude rate		57.9	-		72.4	-		44.4	-
. AAMR (1940 Std. mil.)		35.2	0.65		48.5	1.11		25.4	0.77
. AAMR (1970 Std. mil.)		45.1	0.79		64.3	1.42		31.9	0.90
. AAMR (2000 Std. mil.)		53.4	0.92		78.9	1.79		36.9	1.02
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		40.7	-		60.4	-		25.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.23	0	0.0	0.00	1	0.5	0.46
25-29	1	0.2	0.20	1	0.4	0.40	0	0.0	0.00
30-34	3	0.6	0.33	2	0.8	0.54	1	0.4	0.38
35-39	12	2.5	0.73	8	3.4	1.20	4	1.7	0.84
40-44	30	6.8	1.24	19	8.8	2.01	11	4.9	1.47
45-49	106	29.5	2.86	68	38.5	4.66	38	20.8	3.37
50-54	152	53.5	4.34	90	64.8	6.83	62	42.7	5.42
55-59	304	111.5	6.39	170	128.8	9.88	134	95.2	8.23
60-64	452	160.3	7.54	281	210.5	12.56	171	115.1	8.80
65-69	600	226.8	9.26	347	291.9	15.67	253	173.6	10.92
70-74	639	289.2	11.44	391	418.6	21.17	248	194.4	12.35
75-79	523	314.6	13.76	325	504.3	27.97	198	194.5	13.82
80-84	306	291.1	16.64	193	549.1	39.53	113	161.5	15.19
85+	185	209.5	15.40	104	457.7	44.89	81	123.5	13.72
Total deaths	3,314			1,999			1,315		
Connecticut									
. Crude rate		56.2	-		70.0	-		43.3	-
. AAMR (1940 Std. mil.)		34.3	0.64		47.0	1.09		24.8	0.76
. AAMR (1970 Std. mil.)		43.9	0.78		62.2	1.40		31.1	0.89
. AAMR (2000 Std. mil.)		51.8	0.91		76.1	1.76		36.0	1.01
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		39.3	-		57.9	-		25.0	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.23	1	0.4	0.45	0	0.0	0.00
25-29	1	0.2	0.20	1	0.4	0.40	0	0.0	0.00
30-34	7	1.3	0.50	3	1.1	0.66	4	1.5	0.76
35-39	9	1.9	0.63	4	1.7	0.85	5	2.1	0.94
40-44	11	2.5	0.75	8	3.7	1.30	3	1.3	0.77
45-49	7	1.9	0.74	4	2.3	1.13	3	1.6	0.95
50-54	12	4.2	1.22	5	3.6	1.61	7	4.8	1.82
55-59	18	6.6	1.56	8	6.1	2.14	10	7.1	2.25
60-64	15	5.3	1.37	8	6.0	2.12	7	4.7	1.78
65-69	26	9.8	1.93	15	12.6	3.26	11	7.5	2.28
70-74	23	10.4	2.17	16	17.1	4.28	7	5.5	2.07
75-79	18	10.8	2.55	8	12.4	4.39	10	9.8	3.11
80-84	9	8.6	2.85	5	14.2	6.36	4	5.7	2.86
85+	14	15.9	4.24	3	13.2	7.62	11	16.8	5.06
Total deaths	171			89			82		
Connecticut									
. Crude rate		2.9	-		3.1	-		2.7	-
. AAMR (1940 Std. mil.)		2.0	0.16		2.3	0.25		1.7	0.21
. AAMR (1970 Std. mil.)		2.3	0.18		2.7	0.29		2.0	0.23
. AAMR (2000 Std. mil.)		2.8	0.21		3.3	0.36		2.4	0.27
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.8	-		2.6	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.3	0.25	-	-	-	1	0.5	0.52
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	4	0.8	0.40	-	-	-	4	1.6	0.80
30-34	12	2.3	0.66	-	-	-	12	4.5	1.31
35-39	29	6.1	1.14	-	-	-	29	12.2	2.26
40-44	57	12.9	1.71	-	-	-	57	25.3	3.35
45-49	54	15.0	2.04	-	-	-	54	29.5	4.02
50-54	57	20.1	2.66	-	-	-	57	39.3	5.20
55-59	95	34.8	3.57	-	-	-	95	67.5	6.93
60-64	153	54.2	4.39	-	-	-	153	103.0	8.33
65-69	144	54.4	4.54	-	-	-	144	98.8	8.24
70-74	176	79.7	6.00	-	-	-	176	138.0	10.40
75-79	151	90.8	7.39	-	-	-	151	148.4	12.07
80-84	113	107.5	10.11	-	-	-	113	161.5	15.19
85+	155	175.5	14.10	-	-	-	155	236.4	18.99
Total deaths	1,201			-			1,201		
Connecticut									
. Crude rate		-	-		-	-		39.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		22.6	0.74
. AAMR (1970 Std. mil.)		-	-		-	-		27.6	0.84
. AAMR (2000 Std. mil.)		-	-		-	-		33.7	0.99
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		23.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	2	0.4	0.27	-	-	-	2	0.8	0.53
35-39	6	1.3	0.52	-	-	-	6	2.5	1.03
40-44	7	1.6	0.60	-	-	-	7	3.1	1.17
45-49	8	2.2	0.79	-	-	-	8	4.4	1.55
50-54	3	1.1	0.61	-	-	-	3	2.1	1.19
55-59	3	1.1	0.64	-	-	-	3	2.1	1.23
60-64	6	2.1	0.87	-	-	-	6	4.0	1.65
65-69	11	4.2	1.25	-	-	-	11	7.5	2.28
70-74	6	2.7	1.11	-	-	-	6	4.7	1.92
75-79	14	8.4	2.25	-	-	-	14	13.8	3.68
80-84	5	4.8	2.13	-	-	-	5	7.1	3.20
85+	8	9.1	3.20	-	-	-	8	12.2	4.31
Total deaths	79			-			79		
Connecticut									
. Crude rate		-	-		-	-		2.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.6	0.20
. AAMR (1970 Std. mil.)		-	-		-	-		1.8	0.21
. AAMR (2000 Std. mil.)		-	-		-	-		2.3	0.27
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is higher
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	1	0.2	0.23	-	-	-	1	0.4	0.44
45-49	1	0.3	0.28	-	-	-	1	0.5	0.55
50-54	0	0.0	0.00	-	-	-	0	0.0	0.00
55-59	5	1.8	0.82	-	-	-	5	3.6	1.59
60-64	12	4.3	1.23	-	-	-	12	8.1	2.33
65-69	16	6.0	1.51	-	-	-	16	11.0	2.75
70-74	14	6.3	1.69	-	-	-	14	11.0	2.93
75-79	11	6.6	2.00	-	-	-	11	10.8	3.26
80-84	11	10.5	3.16	-	-	-	11	15.7	4.74
85+	13	14.7	4.08	-	-	-	13	19.8	5.50
Total deaths	84			-			84		
Connecticut									
. Crude rate		-	-		-	-		2.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.3	0.16
. AAMR (1970 Std. mil.)		-	-		-	-		1.8	0.20
. AAMR (2000 Std. mil.)		-	-		-	-		2.2	0.24
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	2	0.4	0.28	-	-	-	2	0.8	0.56
30-34	2	0.4	0.27	-	-	-	2	0.8	0.53
35-39	1	0.2	0.21	-	-	-	1	0.4	0.42
40-44	9	2.0	0.68	-	-	-	9	4.0	1.33
45-49	14	3.9	1.04	-	-	-	14	7.7	2.04
50-54	22	7.7	1.65	-	-	-	22	15.2	3.23
55-59	25	9.2	1.83	-	-	-	25	17.8	3.55
60-64	41	14.5	2.27	-	-	-	41	27.6	4.31
65-69	58	21.9	2.88	-	-	-	58	39.8	5.23
70-74	53	24.0	3.29	-	-	-	53	41.5	5.71
75-79	47	28.3	4.12	-	-	-	47	46.2	6.74
80-84	26	24.7	4.85	-	-	-	26	37.2	7.29
85+	30	34.0	6.20	-	-	-	30	45.7	8.35
Total deaths	330			-			330		
Connecticut									
. Crude rate		-	-		-	-		10.9	-
. AAMR (1940 Std. mil.)		-	-		-	-		6.3	0.39
. AAMR (1970 Std. mil.)		-	-		-	-		7.8	0.45
. AAMR (2000 Std. mil.)		-	-		-	-		9.2	0.52
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Bladder cancer (188)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.4	0.30	2	0.8	0.60	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	2	0.6	0.39	0	0.0	0.00	2	1.1	0.77
50-54	4	1.4	0.70	3	2.2	1.25	1	0.7	0.69
55-59	11	4.0	1.22	8	6.1	2.14	3	2.1	1.23
60-64	24	8.5	1.74	19	14.2	3.27	5	3.4	1.51
65-69	27	10.2	1.96	20	16.8	3.76	7	4.8	1.82
70-74	55	24.9	3.36	41	43.9	6.86	14	11.0	2.93
75-79	54	32.5	4.42	41	63.6	9.94	13	12.8	3.54
80-84	60	57.1	7.37	40	113.8	18.00	20	28.6	6.39
85+	78	88.3	10.00	49	215.7	30.81	29	44.2	8.21
Total deaths	317			223			94		
Connecticut									
. Crude rate		5.4	-		7.8	-		3.1	-
. AAMR (1940 Std. mil.)		2.4	0.15		4.3	0.30		1.1	0.14
. AAMR (1970 Std. mil.)		3.6	0.21		6.9	0.46		1.7	0.19
. AAMR (2000 Std. mil.)		5.1	0.29		10.1	0.71		2.4	0.25
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.8	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.5	0.36	1	0.5	0.49	1	0.5	0.52
5-9	5	1.4	0.63	3	1.6	0.95	2	1.2	0.82
10-14	2	0.6	0.43	1	0.6	0.59	1	0.6	0.62
15-19	5	1.3	0.60	3	1.6	0.90	2	1.1	0.78
20-24	4	0.9	0.45	3	1.3	0.78	1	0.5	0.46
25-29	6	1.2	0.49	2	0.8	0.56	4	1.6	0.80
30-34	10	1.9	0.60	8	3.0	1.07	2	0.8	0.53
35-39	17	3.6	0.87	7	3.0	1.12	10	4.2	1.33
40-44	10	2.3	0.71	6	2.8	1.13	4	1.8	0.89
45-49	15	4.2	1.08	8	4.5	1.60	7	3.8	1.45
50-54	15	5.3	1.36	8	5.8	2.04	7	4.8	1.82
55-59	26	9.5	1.87	15	11.4	2.93	11	7.8	2.36
60-64	32	11.3	2.01	21	15.7	3.43	11	7.4	2.23
65-69	56	21.2	2.83	32	26.9	4.76	24	16.5	3.36
70-74	56	25.3	3.39	31	33.2	5.96	25	19.6	3.92
75-79	74	44.5	5.17	35	54.3	9.18	39	38.3	6.14
80-84	63	59.9	7.55	30	85.4	15.58	33	47.2	8.21
85+	63	71.4	8.99	23	101.2	21.11	40	61.0	9.64
Total deaths	461			237			224		
Connecticut									
. Crude rate		7.8	-		8.3	-		7.4	-
. AAMR (1940 Std. mil.)		4.6	0.24		5.7	0.39		3.8	0.31
. AAMR (1970 Std. mil.)		5.9	0.29		7.4	0.49		4.8	0.35
. AAMR (2000 Std. mil.)		7.5	0.35		9.5	0.64		6.2	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.1	-		6.5	-		3.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes mellitus (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.23	0	0.0	0.00	1	0.5	0.46
25-29	3	0.6	0.34	2	0.8	0.56	1	0.4	0.40
30-34	5	0.9	0.42	3	1.1	0.66	2	0.8	0.53
35-39	7	1.5	0.56	6	2.5	1.04	1	0.4	0.42
40-44	15	3.4	0.88	11	5.1	1.53	4	1.8	0.89
45-49	18	5.0	1.18	8	4.5	1.60	10	5.5	1.73
50-54	22	7.7	1.65	15	10.8	2.79	7	4.8	1.82
55-59	35	12.8	2.17	20	15.2	3.39	15	10.7	2.75
60-64	77	27.3	3.11	43	32.2	4.91	34	22.9	3.92
65-69	113	42.7	4.02	63	53.0	6.68	50	34.3	4.85
70-74	126	57.0	5.08	57	61.0	8.08	69	54.1	6.51
75-79	159	95.7	7.59	72	111.7	13.17	87	85.5	9.16
80-84	137	130.3	11.13	42	119.5	18.44	95	135.8	13.93
85+	158	178.9	14.24	35	154.0	26.04	123	187.6	16.91
Total deaths	876			377			499		
Connecticut									
. Crude rate		14.9	-		13.2	-		16.4	-
. AAMR (1940 Std. mil.)		7.5	0.28		8.5	0.46		6.5	0.35
. AAMR (1970 Std. mil.)		10.5	0.36		11.6	0.60		9.5	0.45
. AAMR (2000 Std. mil.)		14.0	0.48		15.0	0.81		12.9	0.58
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.6	-		12.3	-		10.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.3	0.25	0	0.0	0.00	1	0.5	0.52
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.27	0	0.0	0.00	1	0.5	0.55
20-24	1	0.2	0.23	0	0.0	0.00	1	0.5	0.46
25-29	5	1.0	0.44	3	1.2	0.68	2	0.8	0.56
30-34	9	1.7	0.57	7	2.7	1.00	2	0.8	0.53
35-39	20	4.2	0.94	13	5.5	1.53	7	2.9	1.11
40-44	36	8.1	1.36	25	11.5	2.30	11	4.9	1.47
45-49	61	17.0	2.17	31	17.5	3.15	30	16.4	2.99
50-54	72	25.3	2.99	45	32.4	4.83	27	18.6	3.58
55-59	161	59.0	4.65	89	67.4	7.15	72	51.2	6.03
60-64	353	125.2	6.66	207	155.1	10.78	146	98.3	8.13
65-69	524	198.0	8.65	273	229.7	13.90	251	172.3	10.87
70-74	682	308.6	11.82	343	367.2	19.83	339	265.8	14.43
75-79	775	466.2	16.75	385	597.4	30.45	390	383.2	19.40
80-84	673	640.2	24.68	256	728.4	45.52	417	596.0	29.18
85+	853	966.1	33.08	245	1078.3	68.89	608	927.2	37.60
Total deaths	4,227			1,922			2,305		
Connecticut									
. Crude rate		71.7	-		67.3	-		75.9	-
. AAMR (1940 Std. mil.)		34.2	0.59		40.8	0.97		29.2	0.72
. AAMR (1970 Std. mil.)		49.8	0.78		59.1	1.35		43.2	0.95
. AAMR (2000 Std. mil.)		67.4	1.04		79.6	1.90		59.2	1.24
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Alzheimer's diseases (331.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	4	1.5	0.73	3	2.3	1.31	1	0.7	0.71
60-64	9	3.2	1.06	4	3.0	1.50	5	3.4	1.51
65-69	19	7.2	1.65	9	7.6	2.52	10	6.9	2.17
70-74	29	13.1	2.44	16	17.1	4.28	13	10.2	2.83
75-79	74	44.5	5.17	36	55.9	9.31	38	37.3	6.06
80-84	85	80.9	8.77	29	82.5	15.32	56	80.0	10.69
85+	169	191.4	14.72	52	228.9	31.74	117	178.4	16.50
Total deaths	389			149			240		
Connecticut									
. Crude rate		6.6	-		5.2	-		7.9	-
. AAMR (1940 Std. mil.)		2.2	0.12		2.5	0.21		1.9	0.15
. AAMR (1970 Std. mil.)		4.0	0.21		4.6	0.38		3.6	0.24
. AAMR (2000 Std. mil.)		6.5	0.33		7.6	0.64		5.9	0.38
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.1	-		2.4	-		2.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cardiovascular disease (390-459)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	29	7.3	1.36	17	8.3	2.02	12	6.2	1.80
5-9	5	1.4	0.63	2	1.1	0.78	3	1.7	1.00
10-14	3	0.9	0.52	3	1.8	1.02	0	0.0	0.00
15-19	5	1.3	0.60	2	1.0	0.74	3	1.6	0.95
20-24	10	2.3	0.72	7	3.1	1.19	3	1.4	0.80
25-29	29	5.8	1.07	15	5.9	1.53	14	5.6	1.49
30-34	56	10.6	1.42	40	15.2	2.40	16	6.0	1.51
35-39	81	17.1	1.90	60	25.4	3.28	21	8.8	1.92
40-44	168	38.0	2.93	122	56.2	5.09	46	20.4	3.01
45-49	271	75.3	4.58	204	115.4	8.08	67	36.6	4.47
50-54	400	140.8	7.04	270	194.4	11.83	130	89.5	7.85
55-59	704	258.2	9.73	493	373.6	16.83	211	149.9	10.32
60-64	1,291	457.7	12.74	882	660.8	22.25	409	275.3	13.61
65-69	1,962	741.5	16.74	1,225	1030.5	29.44	737	505.8	18.63
70-74	2,819	1275.8	24.03	1,611	1724.8	42.97	1,208	947.0	27.25
75-79	3,487	2097.7	35.52	1,913	2968.4	67.87	1,574	1546.4	38.98
80-84	3,892	3702.5	59.35	1,703	4845.5	117.42	2,189	3128.4	66.86
85+	7,566	8569.0	98.51	2,089	9194.5	201.17	5,477	8352.3	112.86
Total deaths	22,778			10,658			12,120		
Connecticut									
. Crude rate		386.6	-		373.1	-		399.2	-
. AAMR (1940 Std. mil.)		162.2	1.24		217.9	2.23		119.8	1.37
. AAMR (1970 Std. mil.)		254.2	1.73		331.7	3.23		198.7	1.92
. AAMR (2000 Std. mil.)		372.6	2.48		472.7	4.79		304.6	2.79
. HP 2000 (1940 Std. mil.)		84.4	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		197.2	-		260.2	-		147.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Heart disease-related (390-398,402,404-429)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	77	19.4	2.21	35	17.2	2.90	42	21.8	3.36
5-9	19	5.3	1.23	10	5.5	1.74	9	5.2	1.73
10-14	7	2.1	0.80	4	2.4	1.18	3	1.8	1.07
15-19	18	4.8	1.13	11	5.7	1.73	7	3.8	1.45
20-24	20	4.5	1.01	11	4.9	1.49	9	4.1	1.38
25-29	53	10.5	1.45	32	12.6	2.24	21	8.4	1.83
30-34	93	17.6	1.82	64	24.2	3.03	29	10.9	2.03
35-39	159	33.5	2.66	106	44.9	4.36	53	22.2	3.05
40-44	256	57.8	3.61	185	85.3	6.27	71	31.5	3.73
45-49	385	107.0	5.45	270	152.7	9.29	115	62.8	5.86
50-54	572	201.3	8.42	375	270.0	13.94	197	135.7	9.67
55-59	974	357.2	11.44	648	491.0	19.29	326	231.7	12.83
60-64	1,818	644.6	15.12	1,183	886.3	25.77	635	427.4	16.96
65-69	2,710	1024.2	19.67	1,625	1367.0	33.91	1,085	744.6	22.60
70-74	3,854	1744.2	28.10	2,202	2357.5	50.24	1,652	1295.1	31.86
75-79	4,631	2785.9	40.94	2,526	3919.6	77.99	2,105	2068.1	45.08
80-84	4,921	4681.4	66.73	2,199	6256.8	133.42	2,722	3890.1	74.56
85+	9,004	10197.6	107.47	2,640	11619.7	226.15	6,364	9704.9	121.65
Total deaths	29,571			14,126			15,445		
Connecticut									
. Crude rate		501.9	-		494.6	-		508.8	-
. AAMR (1940 Std. mil.)		219.4	1.47		291.9	2.59		165.3	1.67
. AAMR (1970 Std. mil.)		335.3	2.01		439.3	3.71		263.3	2.27
. AAMR (2000 Std. mil.)		482.0	2.82		621.2	5.47		391.5	3.18
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									No comparison is available
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.8	0.44	0	0.0	0.00	3	1.6	0.90
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.27	0	0.0	0.00	1	0.5	0.55
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	5	1.0	0.44	3	1.2	0.68	2	0.8	0.56
30-34	19	3.6	0.82	17	6.4	1.56	2	0.8	0.53
35-39	36	7.6	1.26	33	14.0	2.44	3	1.3	0.73
40-44	102	23.0	2.28	88	40.6	4.32	14	6.2	1.66
45-49	166	46.1	3.58	132	74.7	6.50	34	18.6	3.19
50-54	247	86.9	5.53	187	134.6	9.85	60	41.3	5.33
55-59	465	170.5	7.91	346	262.2	14.10	119	84.6	7.75
60-64	853	302.4	10.36	615	460.8	18.58	238	160.2	10.38
65-69	1,247	471.3	13.35	803	675.5	23.84	444	304.7	14.46
70-74	1,788	809.2	19.14	1,078	1154.1	35.15	710	556.6	20.89
75-79	2,087	1255.5	27.48	1,193	1851.2	53.60	894	878.3	29.38
80-84	2,255	2145.2	45.17	1,016	2890.8	90.69	1,239	1770.7	50.31
85+	4,423	5009.3	75.32	1,264	5563.4	156.48	3,159	4817.4	85.71
Total deaths	13,697			6,775			6,922		
Connecticut									
. Crude rate		232.5	-		237.2	-		228.0	-
. AAMR (1940 Std. mil.)		98.8	0.97		140.1	1.79		67.1	1.00
. AAMR (1970 Std. mil.)		153.8	1.35		210.8	2.57		112.7	1.44
. AAMR (2000 Std. mil.)		223.4	1.92		297.0	3.78		173.2	2.09
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		125.6	-		174.7	-		87.0	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.8	0.44	3	1.5	0.85	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.27	0	0.0	0.00	1	0.5	0.55
20-24	1	0.2	0.23	1	0.4	0.45	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.2	0.19	0	0.0	0.00	1	0.4	0.38
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	3	0.8	0.48	3	1.7	0.98	0	0.0	0.00
50-54	3	1.1	0.61	1	0.7	0.72	2	1.4	0.97
55-59	4	1.5	0.73	4	3.0	1.52	0	0.0	0.00
60-64	13	4.6	1.28	9	6.7	2.25	4	2.7	1.35
65-69	39	14.7	2.36	22	18.5	3.95	17	11.7	2.83
70-74	57	25.8	3.42	36	38.5	6.42	21	16.5	3.59
75-79	113	68.0	6.39	62	96.2	12.22	51	50.1	7.02
80-84	136	129.4	11.09	56	159.3	21.29	80	114.3	12.78
85+	388	439.4	22.31	107	471.0	45.53	281	428.5	25.56
Total deaths	762			304			458		
Connecticut									
. Crude rate		12.9	-		10.6	-		15.1	-
. AAMR (1940 Std. mil.)		4.1	0.17		5.3	0.32		3.3	0.19
. AAMR (1970 Std. mil.)		7.7	0.28		9.5	0.55		6.5	0.32
. AAMR (2000 Std. mil.)		12.8	0.46		15.3	0.91		11.2	0.53
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.7	-		6.6	-		5.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	2	0.5	0.32	2	0.9	0.65	0	0.0	0.00
45-49	2	0.6	0.39	1	0.6	0.57	1	0.5	0.55
50-54	4	1.4	0.70	2	1.4	1.02	2	1.4	0.97
55-59	7	2.6	0.97	6	4.5	1.86	1	0.7	0.71
60-64	11	3.9	1.18	7	5.2	1.98	4	2.7	1.35
65-69	15	5.7	1.46	11	9.3	2.79	4	2.7	1.37
70-74	31	14.0	2.52	12	12.8	3.71	19	14.9	3.42
75-79	31	18.6	3.35	12	18.6	5.38	19	18.7	4.28
80-84	33	31.4	5.46	7	19.9	7.53	26	37.2	7.29
85+	73	82.7	9.68	16	70.4	17.61	57	86.9	11.51
Total deaths	209			76			133		
Connecticut									
. Crude rate		3.5	-		2.7	-		4.4	-
. AAMR (1940 Std. mil.)		1.5	0.12		1.6	0.19		1.3	0.14
. AAMR (1970 Std. mil.)		2.3	0.16		2.4	0.27		2.2	0.20
. AAMR (2000 Std. mil.)		3.4	0.24		3.3	0.40		3.3	0.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.8	-		2.1	-		1.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Atherosclerosis (440)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.3	0.25	0	0.0	0.00	1	0.5	0.52
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	4	1.1	0.56	2	1.1	0.80	2	1.1	0.77
50-54	1	0.4	0.35	0	0.0	0.00	1	0.7	0.69
55-59	4	1.5	0.73	4	3.0	1.52	0	0.0	0.00
60-64	8	2.8	1.00	6	4.5	1.84	2	1.3	0.95
65-69	15	5.7	1.46	7	5.9	2.23	8	5.5	1.94
70-74	31	14.0	2.52	19	20.3	4.67	12	9.4	2.72
75-79	40	24.1	3.80	22	34.1	7.28	18	17.7	4.17
80-84	54	51.4	6.99	21	59.8	13.04	33	47.2	8.21
85+	230	260.5	17.18	54	237.7	32.34	176	268.4	20.23
Total deaths	388			135			253		
Connecticut									
. Crude rate		6.6	-		4.7	-		8.3	-
. AAMR (1940 Std. mil.)		2.0	0.12		2.3	0.21		1.8	0.14
. AAMR (1970 Std. mil.)		3.8	0.20		4.2	0.37		3.5	0.23
. AAMR (2000 Std. mil.)		6.5	0.33		6.9	0.62		6.2	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.0	-		3.4	-		2.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Aortic aneurysm (441)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.27	1	0.5	0.52	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.4	0.28	0	0.0	0.00	2	0.8	0.56
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	3	0.7	0.39	2	0.9	0.65	1	0.4	0.44
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	3	1.1	0.61	3	2.2	1.25	0	0.0	0.00
55-59	10	3.7	1.16	6	4.5	1.86	4	2.8	1.42
60-64	27	9.6	1.84	21	15.7	3.43	6	4.0	1.65
65-69	45	17.0	2.54	37	31.1	5.12	8	5.5	1.94
70-74	55	24.9	3.36	40	42.8	6.77	15	11.8	3.04
75-79	74	44.5	5.17	46	71.4	10.52	28	27.5	5.20
80-84	83	79.0	8.67	55	156.5	21.10	28	40.0	7.56
85+	82	92.9	10.26	28	123.2	23.29	54	82.3	11.21
Total deaths	385			239			146		
Connecticut									
. Crude rate		6.5	-		8.4	-		4.8	-
. AAMR (1940 Std. mil.)		2.9	0.16		4.8	0.32		1.5	0.15
. AAMR (1970 Std. mil.)		4.4	0.23		7.4	0.48		2.5	0.22
. AAMR (2000 Std. mil.)		6.1	0.31		10.1	0.68		3.7	0.31
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.7	-		5.9	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	10	2.5	0.80	2	1.0	0.69	8	4.2	1.47
5-9	1	0.3	0.28	0	0.0	0.00	1	0.6	0.58
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.27	0	0.0	0.00	1	0.5	0.55
20-24	4	0.9	0.45	2	0.9	0.63	2	0.9	0.65
25-29	3	0.6	0.34	3	1.2	0.68	0	0.0	0.00
30-34	4	0.8	0.38	2	0.8	0.54	2	0.8	0.53
35-39	11	2.3	0.70	6	2.5	1.04	5	2.1	0.94
40-44	11	2.5	0.75	6	2.8	1.13	5	2.2	0.99
45-49	15	4.2	1.08	7	4.0	1.50	8	4.4	1.55
50-54	20	7.0	1.57	12	8.6	2.49	8	5.5	1.95
55-59	30	11.0	2.01	22	16.7	3.55	8	5.7	2.01
60-64	52	18.4	2.56	39	29.2	4.68	13	8.8	2.43
65-69	97	36.7	3.72	57	47.9	6.35	40	27.5	4.34
70-74	173	78.3	5.95	110	117.8	11.23	63	49.4	6.22
75-79	251	151.0	9.53	128	198.6	17.56	123	120.8	10.90
80-84	409	389.1	19.24	183	520.7	38.49	226	323.0	21.48
85+	1,030	1166.5	36.35	328	1443.7	79.71	702	1070.5	40.40
Total deaths	2,122			907			1,215		
Connecticut									
. Crude rate		36.0	-		31.8	-		40.0	-
. AAMR (1940 Std. mil.)		12.4	0.32		16.2	0.57		10.0	0.38
. AAMR (1970 Std. mil.)		21.9	0.49		28.5	0.95		18.2	0.56
. AAMR (2000 Std. mil.)		35.6	0.78		45.9	1.59		30.2	0.87
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.7	-		18.2	-		10.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.5	0.36	2	1.0	0.69	0	0.0	0.00
5-9	1	0.3	0.28	0	0.0	0.00	1	0.6	0.58
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.23	1	0.4	0.45	0	0.0	0.00
25-29	2	0.4	0.28	1	0.4	0.40	1	0.4	0.40
30-34	4	0.8	0.38	2	0.8	0.54	2	0.8	0.53
35-39	3	0.6	0.37	0	0.0	0.00	3	1.3	0.73
40-44	4	0.9	0.45	2	0.9	0.65	2	0.9	0.63
45-49	12	3.3	0.96	5	2.8	1.26	7	3.8	1.45
50-54	21	7.4	1.61	9	6.5	2.16	12	8.3	2.39
55-59	66	24.2	2.98	32	24.2	4.29	34	24.2	4.14
60-64	129	45.7	4.03	63	47.2	5.95	66	44.4	5.47
65-69	221	83.5	5.62	130	109.4	9.59	91	62.4	6.55
70-74	354	160.2	8.51	189	202.3	14.72	165	129.3	10.07
75-79	422	253.9	12.36	229	355.3	23.48	193	189.6	13.65
80-84	336	319.6	17.44	183	520.7	38.49	153	218.7	17.68
85+	329	372.6	20.54	159	699.8	55.50	170	259.2	19.88
Total deaths	1,907			1,007			900		
Connecticut									
. Crude rate		32.4	-		35.3	-		29.6	-
. AAMR (1940 Std. mil.)		14.9	0.38		19.7	0.65		12.1	0.47
. AAMR (1970 Std. mil.)		22.3	0.52		30.9	0.98		17.6	0.62
. AAMR (2000 Std. mil.)		30.2	0.70		43.8	1.44		23.3	0.78
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		19.5	-		26.9	-		14.8	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.3	0.25	1	0.5	0.49	0	0.0	0.00
5-9	1	0.3	0.28	0	0.0	0.00	1	0.6	0.58
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.23	1	0.4	0.45	0	0.0	0.00
25-29	1	0.2	0.20	1	0.4	0.40	0	0.0	0.00
30-34	2	0.4	0.27	1	0.4	0.38	1	0.4	0.38
35-39	2	0.4	0.30	0	0.0	0.00	2	0.8	0.59
40-44	1	0.2	0.23	0	0.0	0.00	1	0.4	0.44
45-49	3	0.8	0.48	3	1.7	0.98	0	0.0	0.00
50-54	5	1.8	0.79	3	2.2	1.25	2	1.4	0.97
55-59	9	3.3	1.10	2	1.5	1.07	7	5.0	1.88
60-64	3	1.1	0.61	3	2.2	1.30	0	0.0	0.00
65-69	7	2.6	1.00	1	0.8	0.84	6	4.1	1.68
70-74	11	5.0	1.50	2	2.1	1.51	9	7.1	2.35
75-79	13	7.8	2.17	5	7.8	3.47	8	7.9	2.78
80-84	8	7.6	2.69	2	5.7	4.02	6	8.6	3.50
85+	13	14.7	4.08	5	22.0	9.84	8	12.2	4.31
Total deaths	81			30			51		
Connecticut									
. Crude rate		1.4	-		1.1	-		1.7	-
. AAMR (1940 Std. mil.)		0.8	0.11		0.8	0.15		0.9	0.15
. AAMR (1970 Std. mil.)		1.1	0.12		1.0	0.18		1.1	0.17
. AAMR (2000 Std. mil.)		1.3	0.15		1.3	0.25		1.4	0.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.4	-		1.3	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.4	0.28	1	0.4	0.40	1	0.4	0.40
30-34	17	3.2	0.78	13	4.9	1.37	4	1.5	0.76
35-39	24	5.1	1.03	18	7.6	1.80	6	2.5	1.03
40-44	32	7.2	1.28	20	9.2	2.06	12	5.3	1.54
45-49	35	9.7	1.64	27	15.3	2.94	8	4.4	1.55
50-54	66	23.2	2.86	42	30.2	4.67	24	16.5	3.37
55-59	58	21.3	2.79	42	31.8	4.91	16	11.4	2.84
60-64	77	27.3	3.11	54	40.5	5.51	23	15.5	3.23
65-69	92	34.8	3.63	51	42.9	6.01	41	28.1	4.39
70-74	74	33.5	3.89	38	40.7	6.60	36	28.2	4.70
75-79	61	36.7	4.70	32	49.7	8.78	29	28.5	5.29
80-84	29	27.6	5.12	13	37.0	10.26	16	22.9	5.72
85+	21	23.8	5.19	11	48.4	14.60	10	15.2	4.82
Total deaths	588			362			226		
Connecticut									
. Crude rate		10.0	-		12.7	-		7.4	-
. AAMR (1940 Std. mil.)		7.2	0.32		10.0	0.55		4.8	0.35
. AAMR (1970 Std. mil.)		8.2	0.35		11.4	0.60		5.6	0.39
. AAMR (2000 Std. mil.)		9.6	0.40		13.4	0.72		6.5	0.45
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		9.2	-		12.8	-		5.5	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis, nephrotic syndrome/nephrosis (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.8	0.44	3	1.5	0.85	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.3	0.30	0	0.0	0.00	1	0.6	0.62
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.23	1	0.4	0.45	0	0.0	0.00
25-29	1	0.2	0.20	0	0.0	0.00	1	0.4	0.40
30-34	3	0.6	0.33	2	0.8	0.54	1	0.4	0.38
35-39	4	0.8	0.42	1	0.4	0.42	3	1.3	0.73
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	9	2.5	0.83	6	3.4	1.39	3	1.6	0.95
50-54	4	1.4	0.70	3	2.2	1.25	1	0.7	0.69
55-59	9	3.3	1.10	7	5.3	2.00	2	1.4	1.00
60-64	40	14.2	2.24	23	17.2	3.59	17	11.4	2.78
65-69	42	15.9	2.45	13	10.9	3.03	29	19.9	3.70
70-74	54	24.4	3.33	28	30.0	5.67	26	20.4	4.00
75-79	75	45.1	5.21	52	80.7	11.19	23	22.6	4.71
80-84	86	81.8	8.82	39	111.0	17.77	47	67.2	9.80
85+	158	178.9	14.24	64	281.7	35.21	94	143.3	14.79
Total deaths	490			242			248		
Connecticut									
. Crude rate		8.3	-		8.5	-		8.2	-
. AAMR (1940 Std. mil.)		3.5	0.19		4.7	0.32		2.8	0.22
. AAMR (1970 Std. mil.)		5.5	0.26		7.6	0.49		4.3	0.30
. AAMR (2000 Std. mil.)		8.0	0.36		11.4	0.76		6.2	0.40
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.5	-		5.7	-		3.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	38	9.6	1.55	25	12.3	2.45	13	6.7	1.87
5-9	14	3.9	1.05	10	5.5	1.74	4	2.3	1.15
10-14	26	7.8	1.53	18	10.6	2.50	8	4.9	1.74
15-19	135	36.1	3.11	107	55.8	5.39	28	15.4	2.91
20-24	184	41.7	3.08	143	64.1	5.36	41	18.8	2.94
25-29	134	26.6	2.30	108	42.7	4.11	26	10.4	2.04
30-34	113	21.4	2.01	95	36.0	3.69	18	6.8	1.60
35-39	87	18.3	1.97	73	30.9	3.62	14	5.9	1.57
40-44	78	17.6	2.00	62	28.6	3.63	16	7.1	1.77
45-49	49	13.6	1.95	36	20.4	3.39	13	7.1	1.97
50-54	48	16.9	2.44	37	26.6	4.38	11	7.6	2.28
55-59	59	21.6	2.82	44	33.3	5.03	15	10.7	2.75
60-64	66	23.4	2.88	48	36.0	5.19	18	12.1	2.86
65-69	77	29.1	3.32	40	33.6	5.32	37	25.4	4.17
70-74	108	48.9	4.70	61	65.3	8.36	47	36.8	5.37
75-79	116	69.8	6.48	60	93.1	12.02	56	55.0	7.35
80-84	121	115.1	10.46	52	148.0	20.52	69	98.6	11.87
85+	219	248.0	16.76	73	321.3	37.61	146	222.6	18.43
Total deaths	1,672			1,092			580		
Connecticut									
. Crude rate		28.4	-		38.2	-		19.1	-
. AAMR (1940 Std. mil.)		22.8	0.62		34.4	1.09		11.6	0.60
. AAMR (1970 Std. mil.)		24.3	0.63		35.8	1.12		13.6	0.64
. AAMR (2000 Std. mil.)		27.4	0.68		40.0	1.26		16.3	0.69
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		33.8	-		49.4	-		18.9	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	1.0	0.50	3	1.5	0.85	1	0.5	0.52
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	0.9	0.52	3	1.8	1.02	0	0.0	0.00
15-19	4	1.1	0.53	4	2.1	1.04	0	0.0	0.00
20-24	2	0.5	0.32	1	0.4	0.45	1	0.5	0.46
25-29	4	0.8	0.40	3	1.2	0.68	1	0.4	0.40
30-34	1	0.2	0.19	1	0.4	0.38	0	0.0	0.00
35-39	4	0.8	0.42	4	1.7	0.85	0	0.0	0.00
40-44	6	1.4	0.55	5	2.3	1.03	1	0.4	0.44
45-49	1	0.3	0.28	1	0.6	0.57	0	0.0	0.00
50-54	3	1.1	0.61	2	1.4	1.02	1	0.7	0.69
55-59	6	2.2	0.90	4	3.0	1.52	2	1.4	1.00
60-64	19	6.7	1.55	11	8.2	2.48	8	5.4	1.90
65-69	14	5.3	1.41	9	7.6	2.52	5	3.4	1.53
70-74	30	13.6	2.48	19	20.3	4.67	11	8.6	2.60
75-79	39	23.5	3.76	15	23.3	6.01	24	23.6	4.81
80-84	50	47.6	6.73	20	56.9	12.72	30	42.9	7.83
85+	137	155.2	13.26	33	145.2	25.28	104	158.6	15.55
Total deaths	327			138			189		
Connecticut									
. Crude rate		5.5	-		4.8	-		6.2	-
. AAMR (1940 Std. mil.)		2.3	0.16		3.1	0.29		1.7	0.16
. AAMR (1970 Std. mil.)		3.6	0.21		4.4	0.38		2.9	0.23
. AAMR (2000 Std. mil.)		5.4	0.30		6.2	0.56		4.7	0.34
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.7	-		3.6	-		1.6	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	0.9	0.52	3	1.8	1.02	0	0.0	0.00
15-19	29	7.8	1.44	24	12.5	2.55	5	2.7	1.23
20-24	53	12.0	1.65	49	22.0	3.14	4	1.8	0.92
25-29	58	11.5	1.51	47	18.6	2.71	11	4.4	1.32
30-34	54	10.2	1.39	44	16.7	2.51	10	3.8	1.19
35-39	50	10.5	1.49	40	17.0	2.68	10	4.2	1.33
40-44	47	10.6	1.55	41	18.9	2.95	6	2.7	1.09
45-49	44	12.2	1.84	34	19.2	3.30	10	5.5	1.73
50-54	31	10.9	1.96	22	15.8	3.38	9	6.2	2.07
55-59	29	10.6	1.97	18	13.6	3.21	11	7.8	2.36
60-64	31	11.0	1.97	25	18.7	3.75	6	4.0	1.65
65-69	30	11.3	2.07	26	21.9	4.29	4	2.7	1.37
70-74	28	12.7	2.39	22	23.6	5.02	6	4.7	1.92
75-79	19	11.4	2.62	13	20.2	5.59	6	5.9	2.41
80-84	15	14.3	3.68	10	28.5	9.00	5	7.1	3.20
85+	15	17.0	4.39	13	57.2	15.87	2	3.0	2.16
Total deaths	536			431			105		
Connecticut									
. Crude rate		9.1	-		15.1	-		3.5	-
. AAMR (1940 Std. mil.)		8.2	0.37		13.6	0.68		3.0	0.31
. AAMR (1970 Std. mil.)		7.9	0.35		13.3	0.66		2.9	0.30
. AAMR (2000 Std. mil.)		8.7	0.38		15.0	0.74		3.3	0.33
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.4	-		18.8	-		4.6	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	7	1.8	0.67	6	2.9	1.20	1	0.5	0.52
5-9	3	0.8	0.49	1	0.5	0.55	2	1.2	0.82
10-14	3	0.9	0.52	2	1.2	0.83	1	0.6	0.62
15-19	23	6.2	1.28	19	9.9	2.27	4	2.2	1.10
20-24	28	6.4	1.20	22	9.9	2.10	6	2.8	1.12
25-29	24	4.8	0.97	21	8.3	1.81	3	1.2	0.69
30-34	31	5.9	1.05	18	6.8	1.61	13	4.9	1.36
35-39	21	4.4	0.97	13	5.5	1.53	8	3.4	1.19
40-44	14	3.2	0.85	7	3.2	1.22	7	3.1	1.17
45-49	14	3.9	1.04	11	6.2	1.88	3	1.6	0.95
50-54	3	1.1	0.61	2	1.4	1.02	1	0.7	0.69
55-59	4	1.5	0.73	3	2.3	1.31	1	0.7	0.71
60-64	7	2.5	0.94	4	3.0	1.50	3	2.0	1.17
65-69	8	3.0	1.07	6	5.0	2.06	2	1.4	0.97
70-74	11	5.0	1.50	4	4.3	2.14	7	5.5	2.07
75-79	1	0.6	0.60	1	1.6	1.55	0	0.0	0.00
80-84	2	1.9	1.35	1	2.8	2.85	1	1.4	1.43
85+	3	3.4	1.96	0	0.0	0.00	3	4.6	2.64
Total deaths	207			141			66		
Connecticut									
. Crude rate		3.5	-		4.9	-		2.2	-
. AAMR (1940 Std. mil.)		3.5	0.25		5.0	0.43		2.0	0.26
. AAMR (1970 Std. mil.)		3.2	0.24		4.6	0.40		1.9	0.25
. AAMR (2000 Std. mil.)		3.3	0.24		4.6	0.39		2.0	0.26
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		9.4	-		14.6	-		4.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide (E960-E969)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	7	1.8	0.67	6	2.9	1.20	1	0.5	0.52
5-9	3	0.8	0.49	1	0.5	0.55	2	1.2	0.82
10-14	3	0.9	0.52	2	1.2	0.83	1	0.6	0.62
15-19	23	6.2	1.28	19	9.9	2.27	4	2.2	1.10
20-24	28	6.4	1.20	22	9.9	2.10	6	2.8	1.12
25-29	24	4.8	0.97	21	8.3	1.81	3	1.2	0.69
30-34	31	5.9	1.05	18	6.8	1.61	13	4.9	1.36
35-39	20	4.2	0.94	13	5.5	1.53	7	2.9	1.11
40-44	14	3.2	0.85	7	3.2	1.22	7	3.1	1.17
45-49	14	3.9	1.04	11	6.2	1.88	3	1.6	0.95
50-54	3	1.1	0.61	2	1.4	1.02	1	0.7	0.69
55-59	4	1.5	0.73	3	2.3	1.31	1	0.7	0.71
60-64	7	2.5	0.94	4	3.0	1.50	3	2.0	1.17
65-69	8	3.0	1.07	6	5.0	2.06	2	1.4	0.97
70-74	11	5.0	1.50	4	4.3	2.14	7	5.5	2.07
75-79	1	0.6	0.60	1	1.6	1.55	0	0.0	0.00
80-84	2	1.9	1.35	1	2.8	2.85	1	1.4	1.43
85+	3	3.4	1.96	0	0.0	0.00	3	4.6	2.64
Total deaths	206			141			65		
Connecticut									
. Crude rate		3.5	-		4.9	-		2.1	-
. AAMR (1940 Std. mil.)		3.5	0.25		5.0	0.43		2.0	0.26
. AAMR (1970 Std. mil.)		3.2	0.24		4.6	0.40		1.8	0.25
. AAMR (2000 Std. mil.)		3.3	0.23		4.6	0.39		2.0	0.25
. HP 2000 (1940 Std. mil.)		5.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		9.3	-		14.3	-		4.2	-
. HP 2000 (1940 Std. mil.)		7.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All injury (E800-E999)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	45	11.4	1.69	31	15.2	2.73	14	7.3	1.94
5-9	17	4.8	1.16	11	6.0	1.82	6	3.5	1.41
10-14	33	9.9	1.73	24	14.1	2.88	9	5.5	1.85
15-19	188	50.3	3.67	151	78.7	6.40	37	20.3	3.34
20-24	267	60.6	3.71	216	96.9	6.59	51	23.4	3.28
25-29	220	43.7	2.95	180	71.1	5.30	40	16.0	2.52
30-34	206	39.0	2.71	162	61.4	4.82	44	16.6	2.50
35-39	165	34.8	2.71	133	56.4	4.89	32	13.4	2.37
40-44	146	33.0	2.73	115	53.0	4.94	31	13.7	2.47
45-49	110	30.6	2.92	83	46.9	5.15	27	14.8	2.84
50-54	82	28.9	3.19	61	43.9	5.62	21	14.5	3.16
55-59	92	33.7	3.52	65	49.3	6.11	27	19.2	3.69
60-64	107	37.9	3.67	79	59.2	6.66	28	18.8	3.56
65-69	116	43.8	4.07	73	61.4	7.19	43	29.5	4.50
70-74	148	67.0	5.51	88	94.2	10.04	60	47.0	6.07
75-79	136	81.8	7.02	74	114.8	13.35	62	60.9	7.74
80-84	138	131.3	11.18	63	179.3	22.58	75	107.2	12.38
85+	237	268.4	17.44	86	378.5	40.82	151	230.3	18.74
Total deaths	2,453			1,695			758		
Connecticut									
. Crude rate		41.6	-		59.3	-		25.0	-
. AAMR (1940 Std. mil.)		35.0	0.77		54.0	1.37		16.8	0.73
. AAMR (1970 Std. mil.)		36.0	0.77		54.6	1.37		18.6	0.75
. AAMR (2000 Std. mil.)		40.1	0.82		60.6	1.53		21.9	0.81
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		55.6	-		84.4	-		28.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	1.0	0.50	3	1.5	0.85	1	0.5	0.52
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	0.9	0.52	3	1.8	1.02	0	0.0	0.00
15-19	3	0.8	0.46	3	1.6	0.90	0	0.0	0.00
20-24	3	0.7	0.39	3	1.3	0.78	0	0.0	0.00
25-29	4	0.8	0.40	3	1.2	0.68	1	0.4	0.40
30-34	2	0.4	0.27	2	0.8	0.54	0	0.0	0.00
35-39	5	1.1	0.47	5	2.1	0.95	0	0.0	0.00
40-44	4	0.9	0.45	4	1.8	0.92	0	0.0	0.00
45-49	1	0.3	0.28	1	0.6	0.57	0	0.0	0.00
50-54	3	1.1	0.61	2	1.4	1.02	1	0.7	0.69
55-59	7	2.6	0.97	5	3.8	1.69	2	1.4	1.00
60-64	19	6.7	1.55	11	8.2	2.48	8	5.4	1.90
65-69	13	4.9	1.36	8	6.7	2.38	5	3.4	1.53
70-74	26	11.8	2.31	17	18.2	4.41	9	7.1	2.35
75-79	27	16.2	3.13	9	14.0	4.66	18	17.7	4.17
80-84	38	36.1	5.86	14	39.8	10.65	24	34.3	7.00
85+	102	115.5	11.44	24	105.6	21.56	78	118.9	13.47
Total deaths	264			117			147		
Connecticut									
. Crude rate		4.5	-		4.1	-		4.8	-
. AAMR (1940 Std. mil.)		2.1	0.15		2.8	0.28		1.3	0.14
. AAMR (1970 Std. mil.)		3.0	0.20		3.8	0.36		2.3	0.21
. AAMR (2000 Std. mil.)		4.4	0.27		5.1	0.49		3.7	0.30
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.6	-		3.6	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	7	2.1	0.80	6	3.5	1.44	1	0.6	0.62
15-19	39	10.4	1.67	32	16.7	2.95	7	3.8	1.45
20-24	49	11.1	1.59	45	20.2	3.01	4	1.8	0.92
25-29	43	8.5	1.30	37	14.6	2.40	6	2.4	0.98
30-34	44	8.3	1.25	31	11.7	2.11	13	4.9	1.36
35-39	39	8.2	1.32	30	12.7	2.32	9	3.8	1.26
40-44	32	7.2	1.28	25	11.5	2.30	7	3.1	1.17
45-49	29	8.1	1.50	23	13.0	2.71	6	3.3	1.34
50-54	14	4.9	1.32	12	8.6	2.49	2	1.4	0.97
55-59	8	2.9	1.04	6	4.5	1.86	2	1.4	1.00
60-64	19	6.7	1.55	18	13.5	3.18	1	0.7	0.67
65-69	18	6.8	1.60	15	12.6	3.26	3	2.1	1.19
70-74	17	7.7	1.87	14	15.0	4.01	3	2.4	1.36
75-79	4	2.4	1.20	4	6.2	3.10	0	0.0	0.00
80-84	10	9.5	3.01	8	22.8	8.05	2	2.9	2.02
85+	6	6.8	2.77	6	26.4	10.78	0	0.0	0.00
Total deaths	378			312			66		
Connecticut									
. Crude rate		6.4	-		10.9	-		2.2	-
. AAMR (1940 Std. mil.)		6.2	0.33		10.4	0.61		2.1	0.27
. AAMR (1970 Std. mil.)		5.8	0.31		10.0	0.59		1.9	0.25
. AAMR (2000 Std. mil.)		6.2	0.32		10.7	0.62		2.1	0.27
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.6	-		23.7	-		4.1	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	0.9	0.52	2	1.2	0.83	1	0.6	0.62
15-19	19	5.1	1.17	15	7.8	2.02	4	2.2	1.10
20-24	23	5.2	1.09	20	9.0	2.01	3	1.4	0.80
25-29	17	3.4	0.82	15	5.9	1.53	2	0.8	0.56
30-34	17	3.2	0.78	11	4.2	1.26	6	2.3	0.92
35-39	12	2.5	0.73	8	3.4	1.20	4	1.7	0.84
40-44	8	1.8	0.64	4	1.8	0.92	4	1.8	0.89
45-49	8	2.2	0.79	6	3.4	1.39	2	1.1	0.77
50-54	2	0.7	0.50	2	1.4	1.02	0	0.0	0.00
55-59	1	0.4	0.37	1	0.8	0.76	0	0.0	0.00
60-64	3	1.1	0.61	2	1.5	1.06	1	0.7	0.67
65-69	2	0.8	0.53	1	0.8	0.84	1	0.7	0.69
70-74	5	2.3	1.01	2	2.1	1.51	3	2.4	1.36
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	2	1.9	1.35	1	2.8	2.85	1	1.4	1.43
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	122			90			32		
Connecticut									
. Crude rate		2.1	-		3.2	-		1.1	-
. AAMR (1940 Std. mil.)		2.2	0.20		3.3	0.35		1.0	0.19
. AAMR (1970 Std. mil.)		2.0	0.19		3.0	0.33		1.0	0.18
. AAMR (2000 Std. mil.)		2.0	0.18		2.9	0.31		1.0	0.18
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.1	-		9.9	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.23	1	0.4	0.45	0	0.0	0.00
25-29	1	0.2	0.20	1	0.4	0.40	0	0.0	0.00
30-34	15	2.8	0.73	11	4.2	1.26	4	1.5	0.76
35-39	23	4.8	1.01	19	8.1	1.85	4	1.7	0.84
40-44	24	5.4	1.11	13	6.0	1.66	11	4.9	1.47
45-49	34	9.5	1.62	26	14.7	2.88	8	4.4	1.55
50-54	44	15.5	2.33	26	18.7	3.67	18	12.4	2.92
55-59	46	16.9	2.49	34	25.8	4.42	12	8.5	2.46
60-64	38	13.5	2.19	27	20.2	3.89	11	7.4	2.23
65-69	42	15.9	2.45	30	25.2	4.61	12	8.2	2.38
70-74	29	13.1	2.44	19	20.3	4.67	10	7.8	2.48
75-79	9	5.4	1.80	7	10.9	4.11	2	2.0	1.39
80-84	7	6.7	2.52	4	11.4	5.69	3	4.3	2.48
85+	7	7.9	3.00	4	17.6	8.80	3	4.6	2.64
Total deaths	320			222			98		
Connecticut									
. Crude rate		5.4	-		7.8	-		3.2	-
. AAMR (1940 Std. mil.)		4.5	0.26		6.6	0.45		2.6	0.28
. AAMR (1970 Std. mil.)		4.7	0.27		7.0	0.48		2.7	0.29
. AAMR (2000 Std. mil.)		5.4	0.31		8.0	0.55		3.2	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.4	-		11.6	-		3.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.23	0	0.0	0.00	1	0.5	0.46
25-29	8	1.6	0.56	7	2.8	1.05	1	0.4	0.40
30-34	15	2.8	0.73	12	4.5	1.31	3	1.1	0.65
35-39	13	2.7	0.76	11	4.7	1.41	2	0.8	0.59
40-44	6	1.4	0.55	4	1.8	0.92	2	0.9	0.63
45-49	6	1.7	0.68	2	1.1	0.80	4	2.2	1.09
50-54	3	1.1	0.61	1	0.7	0.72	2	1.4	0.97
55-59	3	1.1	0.64	1	0.8	0.76	2	1.4	1.00
60-64	3	1.1	0.61	2	1.5	1.06	1	0.7	0.67
65-69	4	1.5	0.76	3	2.5	1.46	1	0.7	0.69
70-74	6	2.7	1.11	3	3.2	1.85	3	2.4	1.36
75-79	4	2.4	1.20	2	3.1	2.19	2	2.0	1.39
80-84	2	1.9	1.35	0	0.0	0.00	2	2.9	2.02
85+	4	4.5	2.27	3	13.2	7.62	1	1.5	1.52
Total deaths	78			51			27		
Connecticut									
. Crude rate		1.3	-		1.8	-		0.9	-
. AAMR (1940 Std. mil.)		1.1	0.13		1.5	0.21		0.7	0.14
. AAMR (1970 Std. mil.)		1.0	0.11		1.3	0.19		0.7	0.14
. AAMR (2000 Std. mil.)		1.2	0.14		1.7	0.25		0.8	0.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.1	-		5.5	-		2.7	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower