

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Septicemia (038)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	10	1.7	0.53	5	1.6	0.73	5	1.7	0.77
5-9	1	0.2	0.19	0	0.0	0.00	1	0.4	0.38
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.4	0.26	1	0.4	0.36	1	0.4	0.38
20-24	3	0.5	0.27	3	0.9	0.53	0	0.0	0.00
25-29	8	1.1	0.38	5	1.3	0.60	3	0.8	0.47
30-34	7	0.9	0.33	4	1.0	0.50	3	0.8	0.44
35-39	9	1.3	0.42	4	1.1	0.56	5	1.4	0.62
40-44	14	2.1	0.56	10	3.0	0.96	4	1.2	0.58
45-49	17	3.1	0.76	10	3.7	1.18	7	2.5	0.96
50-54	25	5.8	1.17	13	6.2	1.72	12	5.5	1.58
55-59	22	5.5	1.16	15	7.7	1.98	7	3.4	1.27
60-64	40	9.6	1.51	20	10.1	2.26	20	9.1	2.03
65-69	86	21.8	2.35	45	25.4	3.79	41	18.9	2.95
70-74	109	32.6	3.13	60	42.4	5.48	49	25.4	3.64
75-79	186	73.5	5.39	84	85.2	9.29	102	66.1	6.54
80-84	186	116.2	8.52	77	143.2	16.32	109	102.6	9.83
85+	346	256.3	13.78	95	273.6	28.07	251	250.3	15.80
Total deaths	1,071			451			620		
Connecticut									
. Crude rate		12.1	-		10.5	-		13.6	-
. AAMR (1940 Std. mil.)		5.2	0.19		6.1	0.31		4.5	0.23
. AAMR (1970 Std. mil.)		7.9	0.25		9.3	0.44		7.1	0.31
. AAMR (2000 Std. mil.)		11.6	0.36		13.4	0.66		10.5	0.43
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.0	-		4.6	-		3.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is higher
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----HIV infection (042-044)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	7	1.2	0.44	2	0.7	0.46	5	1.7	0.77
5-9	2	0.4	0.26	0	0.0	0.00	2	0.8	0.54
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.2	0.18	0	0.0	0.00	1	0.4	0.38
20-24	13	2.0	0.55	8	2.4	0.86	5	1.6	0.70
25-29	56	7.5	1.01	43	11.5	1.75	13	3.5	0.98
30-34	124	15.6	1.40	103	26.0	2.56	21	5.3	1.15
35-39	105	14.6	1.42	92	25.7	2.68	13	3.6	1.00
40-44	75	11.2	1.29	67	20.3	2.48	8	2.3	0.83
45-49	51	9.4	1.31	47	17.6	2.56	4	1.4	0.72
50-54	41	9.6	1.50	36	17.2	2.87	5	2.3	1.02
55-59	14	3.5	0.93	13	6.7	1.85	1	0.5	0.48
60-64	12	2.9	0.83	11	5.5	1.67	1	0.5	0.45
65-69	8	2.0	0.72	6	3.4	1.38	2	0.9	0.65
70-74	5	1.5	0.67	3	2.1	1.22	2	1.0	0.73
75-79	4	1.6	0.79	1	1.0	1.01	3	1.9	1.12
80-84	1	0.6	0.62	1	1.9	1.86	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	519			433			86		
Connecticut									
. Crude rate		5.9	-		10.1	-		1.9	-
. AAMR (1940 Std. mil.)		5.5	0.24		9.2	0.44		1.8	0.20
. AAMR (1970 Std. mil.)		4.7	0.21		7.8	0.39		1.6	0.18
. AAMR (2000 Std. mil.)		5.7	0.25		9.7	0.47		1.8	0.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.7	-		15.7	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	9	1.5	0.50	5	1.6	0.73	4	1.4	0.69
5-9	20	3.7	0.83	8	2.9	1.03	12	4.6	1.32
10-14	9	1.8	0.60	6	2.3	0.96	3	1.2	0.71
15-19	12	2.2	0.64	8	2.9	1.01	4	1.5	0.75
20-24	40	6.2	0.97	22	6.7	1.42	18	5.6	1.32
25-29	59	7.9	1.03	31	8.3	1.49	28	7.6	1.43
30-34	116	14.6	1.36	57	14.4	1.90	59	14.8	1.93
35-39	179	24.9	1.86	77	21.5	2.45	102	28.2	2.79
40-44	332	49.4	2.71	133	40.4	3.50	199	58.1	4.12
45-49	549	100.9	4.31	263	98.3	6.06	286	103.4	6.11
50-54	764	178.6	6.46	377	180.3	9.29	387	177.1	9.00
55-59	1,277	316.9	8.87	642	329.0	12.99	635	305.5	12.12
60-64	2,076	496.4	10.89	1,120	564.8	16.88	956	434.6	14.06
65-69	2,883	731.9	13.63	1,584	895.3	22.49	1,299	598.7	16.61
70-74	3,305	989.5	17.21	1,802	1273.8	30.01	1,503	780.6	20.14
75-79	3,099	1224.9	22.00	1,595	1617.5	40.50	1,504	974.1	25.12
80-84	2,453	1533.1	30.95	1,215	2259.5	64.82	1,238	1165.4	33.12
85+	2,342	1734.6	35.84	937	2699.0	88.17	1,405	1400.8	37.37
Total deaths	19,524			9,882			9,642		
Connecticut									
. Crude rate		221.1	-		230.8	-		212.0	-
. AAMR (1940 Std. mil.)		124.4	0.99		148.3	1.57		108.7	1.28
. AAMR (1970 Std. mil.)		164.3	1.21		203.1	2.06		140.3	1.51
. AAMR (2000 Std. mil.)		205.6	1.49		261.0	2.73		173.9	1.80
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		134.3	-		165.4	-		112.2	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Oral cavity or pharynx (140.0-149.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.1	0.13	1	0.3	0.25	0	0.0	0.00
35-39	5	0.7	0.31	4	1.1	0.56	1	0.3	0.28
40-44	6	0.9	0.36	5	1.5	0.68	1	0.3	0.29
45-49	13	2.4	0.66	12	4.5	1.29	1	0.4	0.36
50-54	29	6.8	1.26	21	10.0	2.19	8	3.7	1.29
55-59	36	8.9	1.49	24	12.3	2.51	12	5.8	1.67
60-64	59	14.1	1.84	44	22.2	3.35	15	6.8	1.76
65-69	58	14.7	1.93	44	24.9	3.75	14	6.5	1.72
70-74	46	13.8	2.03	24	17.0	3.46	22	11.4	2.44
75-79	44	17.4	2.62	29	29.4	5.46	15	9.7	2.51
80-84	31	19.4	3.48	18	33.5	7.89	13	12.2	3.39
85+	25	18.5	3.70	11	31.7	9.55	14	14.0	3.73
Total deaths	353			237			116		
Connecticut									
. Crude rate		4.0	-		5.5	-		2.6	-
. AAMR (1940 Std. mil.)		2.6	0.15		4.0	0.27		1.4	0.15
. AAMR (1970 Std. mil.)		3.2	0.17		5.0	0.33		1.8	0.17
. AAMR (2000 Std. mil.)		3.8	0.20		5.9	0.40		2.1	0.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.4	-		3.7	-		1.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.15	1	0.3	0.30	0	0.0	0.00
25-29	2	0.3	0.19	2	0.5	0.38	0	0.0	0.00
30-34	8	1.0	0.36	1	0.3	0.25	7	1.8	0.67
35-39	14	1.9	0.52	9	2.5	0.84	5	1.4	0.62
40-44	29	4.3	0.80	12	3.6	1.05	17	5.0	1.20
45-49	41	7.5	1.18	21	7.9	1.71	20	7.2	1.62
50-54	58	13.6	1.78	31	14.8	2.66	27	12.4	2.38
55-59	106	26.3	2.55	57	29.2	3.87	49	23.6	3.37
60-64	210	50.2	3.46	119	60.0	5.50	91	41.4	4.34
65-69	303	76.9	4.42	166	93.8	7.28	137	63.1	5.39
70-74	441	132.0	6.29	247	174.6	11.11	194	100.8	7.23
75-79	428	169.2	8.18	216	219.1	14.90	212	137.3	9.43
80-84	355	221.9	11.78	153	284.5	23.00	202	190.2	13.38
85+	454	336.3	15.78	150	432.1	35.28	304	303.1	17.38
Total deaths	2,450			1,185			1,265		
Connecticut									
. Crude rate		27.7	-		27.7	-		27.8	-
. AAMR (1940 Std. mil.)		13.7	0.31		16.7	0.51		11.5	0.39
. AAMR (1970 Std. mil.)		19.4	0.40		24.1	0.70		16.3	0.49
. AAMR (2000 Std. mil.)		25.8	0.53		32.3	0.98		21.9	0.62
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		14.0	-		17.0	-		11.7	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.3	0.18	2	0.5	0.36	0	0.0	0.00
35-39	5	0.7	0.31	1	0.3	0.28	4	1.1	0.55
40-44	4	0.6	0.30	3	0.9	0.53	1	0.3	0.29
45-49	23	4.2	0.88	13	4.9	1.35	10	3.6	1.14
50-54	31	7.2	1.30	20	9.6	2.14	11	5.0	1.52
55-59	52	12.9	1.79	33	16.9	2.94	19	9.1	2.10
60-64	101	24.1	2.40	56	28.2	3.77	45	20.5	3.05
65-69	161	40.9	3.22	100	56.5	5.65	61	28.1	3.60
70-74	181	54.2	4.03	95	67.2	6.89	86	44.7	4.82
75-79	169	66.8	5.14	61	61.9	7.92	108	70.0	6.73
80-84	131	81.9	7.15	65	120.9	14.99	66	62.1	7.65
85+	143	105.9	8.86	42	121.0	18.67	101	100.7	10.02
Total deaths	1,003			491			512		
Connecticut									
. Crude rate		11.4	-		11.5	-		11.3	-
. AAMR (1940 Std. mil.)		6.0	0.21		7.4	0.35		4.9	0.25
. AAMR (1970 Std. mil.)		8.2	0.27		10.1	0.46		6.8	0.32
. AAMR (2000 Std. mil.)		10.5	0.33		12.8	0.60		8.9	0.40
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.4	-		7.5	-		5.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung and other respiratory cancer (160-165)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.2	0.18	1	0.4	0.36	0	0.0	0.00
20-24	1	0.2	0.15	0	0.0	0.00	1	0.3	0.31
25-29	1	0.1	0.13	1	0.3	0.27	0	0.0	0.00
30-34	7	0.9	0.33	3	0.8	0.44	4	1.0	0.50
35-39	17	2.4	0.57	13	3.6	1.01	4	1.1	0.55
40-44	55	8.2	1.10	32	9.7	1.72	23	6.7	1.40
45-49	160	29.4	2.32	97	36.3	3.68	63	22.8	2.87
50-54	249	58.2	3.69	142	67.9	5.70	107	49.0	4.73
55-59	434	107.7	5.17	243	124.5	7.99	191	91.9	6.65
60-64	670	160.2	6.19	417	210.3	10.30	253	115.0	7.23
65-69	936	237.6	7.77	558	315.4	13.35	378	174.2	8.96
70-74	983	294.3	9.39	603	426.2	17.36	380	197.4	10.12
75-79	824	325.7	11.35	504	511.1	22.77	320	207.3	11.59
80-84	518	323.7	14.22	335	623.0	34.04	183	172.3	12.73
85+	327	242.2	13.39	179	515.6	38.54	148	147.6	12.13
Total deaths	5,183			3,128			2,055		
Connecticut									
. Crude rate		58.7	-		73.1	-		45.2	-
. AAMR (1940 Std. mil.)		35.3	0.53		48.4	0.90		25.6	0.63
. AAMR (1970 Std. mil.)		45.4	0.64		64.5	1.16		32.2	0.74
. AAMR (2000 Std. mil.)		54.0	0.76		79.5	1.47		37.6	0.84
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		40.7	-		60.4	-		25.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.15	0	0.0	0.00	1	0.3	0.31
25-29	1	0.1	0.13	1	0.3	0.27	0	0.0	0.00
30-34	7	0.9	0.33	3	0.8	0.44	4	1.0	0.50
35-39	17	2.4	0.57	13	3.6	1.01	4	1.1	0.55
40-44	52	7.7	1.07	30	9.1	1.66	22	6.4	1.37
45-49	158	29.0	2.31	97	36.3	3.68	61	22.0	2.82
50-54	242	56.6	3.64	138	66.0	5.62	104	47.6	4.67
55-59	422	104.7	5.10	234	119.9	7.84	188	90.4	6.60
60-64	653	156.1	6.11	406	204.8	10.16	247	112.3	7.15
65-69	907	230.3	7.65	536	302.9	13.09	371	171.0	8.88
70-74	957	286.5	9.26	585	413.5	17.10	372	193.2	10.02
75-79	801	316.6	11.19	487	493.9	22.38	314	203.4	11.48
80-84	490	306.2	13.83	313	582.1	32.90	177	166.6	12.52
85+	311	230.3	13.06	172	495.4	37.78	139	138.6	11.75
Total deaths	5,019			3,015			2,004		
Connecticut									
. Crude rate		56.8	-		70.4	-		44.1	-
. AAMR (1940 Std. mil.)		34.2	0.52		46.7	0.89		25.0	0.62
. AAMR (1970 Std. mil.)		44.0	0.63		62.2	1.14		31.4	0.73
. AAMR (2000 Std. mil.)		52.3	0.75		76.5	1.44		36.6	0.83
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		39.3	-		57.9	-		25.0	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.17	-	-	-	1	0.3	0.34
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	5	0.7	0.30	-	-	-	5	1.4	0.60
30-34	14	1.8	0.47	-	-	-	14	3.5	0.94
35-39	42	5.8	0.90	-	-	-	42	11.6	1.79
40-44	80	11.9	1.33	-	-	-	80	23.4	2.61
45-49	90	16.5	1.74	-	-	-	90	32.5	3.43
50-54	91	21.3	2.23	-	-	-	91	41.6	4.36
55-59	144	35.7	2.98	-	-	-	144	69.3	5.77
60-64	217	51.9	3.52	-	-	-	217	98.7	6.70
65-69	227	57.6	3.82	-	-	-	227	104.6	6.94
70-74	247	73.9	4.71	-	-	-	247	128.3	8.16
75-79	231	91.3	6.01	-	-	-	231	149.6	9.84
80-84	176	110.0	8.29	-	-	-	176	165.7	12.49
85+	212	157.0	10.78	-	-	-	212	211.4	14.52
Total deaths	1,777			-			1,777		
Connecticut									
. Crude rate		-	-		-	-		39.1	-
. AAMR (1940 Std. mil.)		-	-		-	-		22.6	0.61
. AAMR (1970 Std. mil.)		-	-		-	-		27.4	0.68
. AAMR (2000 Std. mil.)		-	-		-	-		33.3	0.81
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		23.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	2	0.3	0.19	-	-	-	2	0.5	0.38
30-34	3	0.4	0.22	-	-	-	3	0.8	0.44
35-39	8	1.1	0.39	-	-	-	8	2.2	0.78
40-44	11	1.6	0.49	-	-	-	11	3.2	0.97
45-49	13	2.4	0.66	-	-	-	13	4.7	1.30
50-54	12	2.8	0.81	-	-	-	12	5.5	1.58
55-59	6	1.5	0.61	-	-	-	6	2.9	1.18
60-64	8	1.9	0.68	-	-	-	8	3.6	1.29
65-69	14	3.6	0.95	-	-	-	14	6.5	1.72
70-74	8	2.4	0.85	-	-	-	8	4.2	1.47
75-79	17	6.7	1.63	-	-	-	17	11.0	2.67
80-84	11	6.9	2.07	-	-	-	11	10.4	3.12
85+	13	9.6	2.67	-	-	-	13	13.0	3.59
Total deaths	126			-			126		
Connecticut									
. Crude rate		-	-		-	-		2.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.8	0.18
. AAMR (1970 Std. mil.)		-	-		-	-		2.0	0.19
. AAMR (2000 Std. mil.)		-	-		-	-		2.5	0.23
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is higher
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	1	0.1	0.15	-	-	-	1	0.3	0.29
45-49	1	0.2	0.18	-	-	-	1	0.4	0.36
50-54	1	0.2	0.23	-	-	-	1	0.5	0.46
55-59	9	2.2	0.74	-	-	-	9	4.3	1.44
60-64	15	3.6	0.93	-	-	-	15	6.8	1.76
65-69	26	6.6	1.29	-	-	-	26	12.0	2.35
70-74	23	6.9	1.44	-	-	-	23	11.9	2.49
75-79	15	5.9	1.53	-	-	-	15	9.7	2.51
80-84	16	10.0	2.50	-	-	-	16	15.1	3.77
85+	20	14.8	3.31	-	-	-	20	19.9	4.46
Total deaths	127			-			127		
Connecticut									
. Crude rate		-	-		-	-		2.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.3	0.13
. AAMR (1970 Std. mil.)		-	-		-	-		1.8	0.17
. AAMR (2000 Std. mil.)		-	-		-	-		2.2	0.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	1	0.2	0.15	-	-	-	1	0.3	0.31
25-29	3	0.4	0.23	-	-	-	3	0.8	0.47
30-34	4	0.5	0.25	-	-	-	4	1.0	0.50
35-39	2	0.3	0.20	-	-	-	2	0.6	0.39
40-44	13	1.9	0.54	-	-	-	13	3.8	1.05
45-49	23	4.2	0.88	-	-	-	23	8.3	1.73
50-54	29	6.8	1.26	-	-	-	29	13.3	2.46
55-59	39	9.7	1.55	-	-	-	39	18.8	3.00
60-64	61	14.6	1.87	-	-	-	61	27.7	3.55
65-69	84	21.3	2.33	-	-	-	84	38.7	4.22
70-74	77	23.1	2.63	-	-	-	77	40.0	4.56
75-79	71	28.1	3.33	-	-	-	71	46.0	5.46
80-84	51	31.9	4.46	-	-	-	51	48.0	6.72
85+	48	35.6	5.13	-	-	-	48	47.9	6.91
Total deaths	506			-			506		
Connecticut									
. Crude rate		-	-		-	-		11.1	-
. AAMR (1940 Std. mil.)		-	-		-	-		6.4	0.32
. AAMR (1970 Std. mil.)		-	-		-	-		7.9	0.37
. AAMR (2000 Std. mil.)		-	-		-	-		9.4	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Bladder cancer (188)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.3	0.20	2	0.6	0.40	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	7	1.3	0.49	2	0.7	0.53	5	1.8	0.81
50-54	6	1.4	0.57	3	1.4	0.83	3	1.4	0.79
55-59	13	3.2	0.89	9	4.6	1.54	4	1.9	0.96
60-64	39	9.3	1.49	29	14.6	2.72	10	4.5	1.44
65-69	43	10.9	1.66	33	18.7	3.25	10	4.6	1.46
70-74	83	24.8	2.73	65	45.9	5.70	18	9.3	2.20
75-79	89	35.2	3.73	62	62.9	7.99	27	17.5	3.37
80-84	84	52.5	5.73	53	98.6	13.54	31	29.2	5.24
85+	107	79.2	7.66	63	181.5	22.86	44	43.9	6.61
Total deaths	473			321			152		
Connecticut									
. Crude rate		5.4	-		7.5	-		3.3	-
. AAMR (1940 Std. mil.)		2.4	0.12		4.1	0.24		1.2	0.12
. AAMR (1970 Std. mil.)		3.6	0.17		6.5	0.36		1.9	0.16
. AAMR (2000 Std. mil.)		5.0	0.23		9.3	0.55		2.6	0.21
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.8	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.5	0.29	1	0.3	0.33	2	0.7	0.49
5-9	8	1.5	0.53	3	1.1	0.63	5	1.9	0.85
10-14	4	0.8	0.40	3	1.2	0.68	1	0.4	0.41
15-19	6	1.1	0.45	4	1.4	0.71	2	0.8	0.53
20-24	8	1.2	0.44	6	1.8	0.74	2	0.6	0.44
25-29	8	1.1	0.38	3	0.8	0.46	5	1.4	0.60
30-34	14	1.8	0.47	9	2.3	0.76	5	1.3	0.56
35-39	20	2.8	0.62	9	2.5	0.84	11	3.0	0.92
40-44	18	2.7	0.63	12	3.6	1.05	6	1.8	0.72
45-49	21	3.9	0.84	13	4.9	1.35	8	2.9	1.02
50-54	24	5.6	1.15	12	5.7	1.66	12	5.5	1.58
55-59	36	8.9	1.49	20	10.3	2.29	16	7.7	1.92
60-64	50	12.0	1.69	32	16.1	2.85	18	8.2	1.93
65-69	81	20.6	2.28	48	27.1	3.92	33	15.2	2.65
70-74	101	30.2	3.01	58	41.0	5.38	43	22.3	3.41
75-79	99	39.1	3.93	46	46.7	6.88	53	34.3	4.72
80-84	88	55.0	5.86	44	81.8	12.34	44	41.4	6.24
85+	95	70.4	7.22	33	95.1	16.55	62	61.8	7.85
Total deaths	684			356			328		
Connecticut									
. Crude rate		7.7	-		8.3	-		7.2	-
. AAMR (1940 Std. mil.)		4.6	0.20		5.7	0.32		3.7	0.25
. AAMR (1970 Std. mil.)		5.8	0.23		7.4	0.40		4.7	0.28
. AAMR (2000 Std. mil.)		7.3	0.28		9.4	0.52		6.0	0.34
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.1	-		6.5	-		3.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes mellitus (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.2	0.20	1	0.4	0.39	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.15	0	0.0	0.00	1	0.3	0.31
25-29	4	0.5	0.27	3	0.8	0.46	1	0.3	0.27
30-34	10	1.3	0.40	5	1.3	0.56	5	1.3	0.56
35-39	14	1.9	0.52	10	2.8	0.88	4	1.1	0.55
40-44	20	3.0	0.67	15	4.6	1.18	5	1.5	0.65
45-49	24	4.4	0.90	13	4.9	1.35	11	4.0	1.20
50-54	33	7.7	1.34	22	10.5	2.24	11	5.0	1.52
55-59	47	11.7	1.70	28	14.4	2.71	19	9.1	2.10
60-64	112	26.8	2.53	63	31.8	4.00	49	22.3	3.18
65-69	160	40.6	3.21	93	52.6	5.45	67	30.9	3.77
70-74	195	58.4	4.18	91	64.3	6.74	104	54.0	5.30
75-79	238	94.1	6.10	109	110.5	10.59	129	83.6	7.36
80-84	216	135.0	9.19	72	133.9	15.78	144	135.6	11.30
85+	238	176.3	11.43	53	152.7	20.97	185	184.4	13.56
Total deaths	1,313			578			735		
Connecticut									
. Crude rate		14.9	-		13.5	-		16.2	-
. AAMR (1940 Std. mil.)		7.4	0.23		8.6	0.38		6.3	0.28
. AAMR (1970 Std. mil.)		10.4	0.29		11.8	0.49		9.1	0.36
. AAMR (2000 Std. mil.)		13.9	0.39		15.3	0.66		12.6	0.47
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.6	-		12.3	-		10.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.17	0	0.0	0.00	1	0.3	0.34
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.2	0.20	1	0.4	0.39	0	0.0	0.00
15-19	1	0.2	0.18	0	0.0	0.00	1	0.4	0.38
20-24	1	0.2	0.15	0	0.0	0.00	1	0.3	0.31
25-29	7	0.9	0.36	5	1.3	0.60	2	0.5	0.38
30-34	18	2.3	0.53	12	3.0	0.87	6	1.5	0.62
35-39	33	4.6	0.80	20	5.6	1.25	13	3.6	1.00
40-44	52	7.7	1.07	37	11.2	1.85	15	4.4	1.13
45-49	83	15.3	1.67	49	18.3	2.62	34	12.3	2.11
50-54	105	24.6	2.40	66	31.6	3.89	39	17.8	2.86
55-59	220	54.6	3.68	123	63.0	5.68	97	46.7	4.74
60-64	488	116.7	5.28	284	143.2	8.50	204	92.7	6.49
65-69	797	202.3	7.17	422	238.5	11.61	375	172.8	8.92
70-74	1,020	305.4	9.56	507	358.4	15.92	513	266.4	11.76
75-79	1,157	457.3	13.44	571	579.1	24.23	586	379.5	15.68
80-84	1,022	638.7	19.98	393	730.9	36.87	629	592.1	23.61
85+	1,312	971.7	26.83	375	1080.2	55.78	937	934.2	30.52
Total deaths	6,318			2,865			3,453		
Connecticut									
. Crude rate		71.6	-		66.9	-		75.9	-
. AAMR (1940 Std. mil.)		33.5	0.47		40.1	0.79		28.5	0.58
. AAMR (1970 Std. mil.)		49.0	0.63		58.2	1.09		42.4	0.76
. AAMR (2000 Std. mil.)		66.5	0.84		78.6	1.53		58.4	1.00
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.2	0.20	1	0.4	0.39	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	3	0.4	0.22	1	0.3	0.25	2	0.5	0.36
35-39	1	0.1	0.14	1	0.3	0.28	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	1	0.2	0.18	0	0.0	0.00	1	0.4	0.36
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	1	0.2	0.25	0	0.0	0.00	1	0.5	0.48
60-64	2	0.5	0.34	1	0.5	0.50	1	0.5	0.45
65-69	3	0.8	0.44	1	0.6	0.57	2	0.9	0.65
70-74	2	0.6	0.42	1	0.7	0.71	1	0.5	0.52
75-79	2	0.8	0.56	2	2.0	1.43	0	0.0	0.00
80-84	2	1.2	0.88	2	3.7	2.63	0	0.0	0.00
85+	3	2.2	1.28	2	5.8	4.07	1	1.0	1.00
Total deaths	21			12			9		
Connecticut									
. Crude rate		0.2	-		0.3	-		0.2	-
. AAMR (1940 Std. mil.)		0.2	0.04		-	-		-	-
. AAMR (1970 Std. mil.)		0.2	0.04		-	-		-	-
. AAMR (2000 Std. mil.)		0.2	0.05		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.6	-		0.6	-		0.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Alzheimer's diseases (331.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	1	0.2	0.23	0	0.0	0.00	1	0.5	0.46
55-59	5	1.2	0.55	4	2.1	1.03	1	0.5	0.48
60-64	12	2.9	0.83	6	3.0	1.24	6	2.7	1.11
65-69	30	7.6	1.39	14	7.9	2.11	16	7.4	1.84
70-74	52	15.6	2.16	22	15.6	3.32	30	15.6	2.84
75-79	103	40.7	4.01	51	51.7	7.24	52	33.7	4.67
80-84	120	75.0	6.85	42	78.1	12.05	78	73.4	8.31
85+	256	189.6	11.85	76	218.9	25.11	180	179.5	13.38
Total deaths	579			215			364		
Connecticut									
. Crude rate		6.6	-		5.0	-		8.0	-
. AAMR (1940 Std. mil.)		2.1	0.10		2.4	0.17		2.0	0.12
. AAMR (1970 Std. mil.)		3.9	0.16		4.4	0.30		3.6	0.20
. AAMR (2000 Std. mil.)		6.3	0.26		7.2	0.51		5.9	0.31
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.1	-		2.4	-		2.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	0.7	0.33	0	0.0	0.00	4	1.4	0.69
5-9	1	0.2	0.19	0	0.0	0.00	1	0.4	0.38
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.2	0.18	0	0.0	0.00	1	0.4	0.38
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	9	1.2	0.40	7	1.9	0.71	2	0.5	0.38
30-34	26	3.3	0.64	23	5.8	1.21	3	0.8	0.44
35-39	61	8.5	1.09	52	14.5	2.02	9	2.5	0.83
40-44	158	23.5	1.87	133	40.4	3.50	25	7.3	1.46
45-49	251	46.1	2.91	210	78.5	5.42	41	14.8	2.31
50-54	366	85.6	4.47	279	133.4	7.99	87	39.8	4.27
55-59	655	162.5	6.35	483	247.5	11.26	172	82.7	6.31
60-64	1,249	298.6	8.45	912	459.9	15.23	337	153.2	8.35
65-69	1,841	467.4	10.89	1,213	685.6	19.68	628	289.4	11.55
70-74	2,618	783.8	15.32	1,560	1102.7	27.92	1,058	549.5	16.89
75-79	3,071	1213.8	21.90	1,748	1772.7	42.40	1,323	856.9	23.56
80-84	3,415	2134.3	36.52	1,517	2821.1	72.43	1,898	1786.7	41.01
85+	6,640	4917.9	60.35	1,911	5504.5	125.92	4,729	4714.9	68.56
Total deaths	20,366			10,048			10,318		
Connecticut									
. Crude rate		230.7	-		234.7	-		226.9	-
. AAMR (1940 Std. mil.)		96.9	0.79		137.4	1.45		65.6	0.81
. AAMR (1970 Std. mil.)		150.8	1.09		206.5	2.07		110.4	1.16
. AAMR (2000 Std. mil.)		219.2	1.54		291.0	3.03		170.1	1.68
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		125.6	-		174.7	-		87.0	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.5	0.29	3	1.0	0.56	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.2	0.18	0	0.0	0.00	1	0.4	0.38
20-24	1	0.2	0.15	1	0.3	0.30	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.3	0.18	1	0.3	0.25	1	0.3	0.25
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.1	0.15	1	0.3	0.30	0	0.0	0.00
45-49	3	0.6	0.32	3	1.1	0.65	0	0.0	0.00
50-54	5	1.2	0.52	3	1.4	0.83	2	0.9	0.65
55-59	4	1.0	0.50	4	2.1	1.03	0	0.0	0.00
60-64	22	5.3	1.12	12	6.1	1.75	10	4.5	1.44
65-69	54	13.7	1.87	32	18.1	3.20	22	10.1	2.16
70-74	99	29.6	2.98	51	36.1	5.05	48	24.9	3.60
75-79	173	68.4	5.20	91	92.3	9.67	82	53.1	5.87
80-84	212	132.5	9.10	88	163.7	17.45	124	116.7	10.48
85+	580	429.6	17.84	156	449.3	35.98	424	422.7	20.53
Total deaths	1,160			446			714		
Connecticut									
. Crude rate		13.1	-		10.4	-		15.7	-
. AAMR (1940 Std. mil.)		4.2	0.14		5.1	0.25		3.5	0.16
. AAMR (1970 Std. mil.)		7.7	0.23		9.2	0.43		6.8	0.27
. AAMR (2000 Std. mil.)		12.7	0.38		14.8	0.73		11.5	0.43
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.7	-		6.6	-		5.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.3	0.18	1	0.3	0.25	1	0.3	0.25
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	2	0.3	0.21	2	0.6	0.43	0	0.0	0.00
45-49	4	0.7	0.37	2	0.7	0.53	2	0.7	0.51
50-54	6	1.4	0.57	3	1.4	0.83	3	1.4	0.79
55-59	10	2.5	0.78	8	4.1	1.45	2	1.0	0.68
60-64	12	2.9	0.83	8	4.0	1.43	4	1.8	0.91
65-69	26	6.6	1.29	19	10.7	2.46	7	3.2	1.22
70-74	42	12.6	1.94	16	11.3	2.83	26	13.5	2.65
75-79	51	20.2	2.82	24	24.3	4.97	27	17.5	3.37
80-84	60	37.5	4.84	20	37.2	8.32	40	37.7	5.95
85+	104	77.0	7.55	22	63.4	13.51	82	81.8	9.03
Total deaths	319			125			194		
Connecticut									
. Crude rate		3.6	-		2.9	-		4.3	-
. AAMR (1940 Std. mil.)		1.5	0.10		1.7	0.16		1.3	0.11
. AAMR (1970 Std. mil.)		2.3	0.13		2.6	0.23		2.1	0.16
. AAMR (2000 Std. mil.)		3.4	0.19		3.6	0.34		3.2	0.23
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.8	-		2.1	-		1.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	0.7	0.33	2	0.7	0.46	2	0.7	0.49
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.2	0.20	1	0.4	0.39	0	0.0	0.00
15-19	1	0.2	0.18	1	0.4	0.36	0	0.0	0.00
20-24	2	0.3	0.22	1	0.3	0.30	1	0.3	0.31
25-29	13	1.7	0.48	4	1.1	0.53	9	2.4	0.81
30-34	17	2.1	0.52	11	2.8	0.84	6	1.5	0.62
35-39	18	2.5	0.59	11	3.1	0.93	7	1.9	0.73
40-44	42	6.3	0.96	17	5.2	1.25	25	7.3	1.46
45-49	41	7.5	1.18	24	9.0	1.83	17	6.1	1.49
50-54	65	15.2	1.89	28	13.4	2.53	37	16.9	2.78
55-59	88	21.8	2.33	41	21.0	3.28	47	22.6	3.30
60-64	163	39.0	3.05	79	39.8	4.48	84	38.2	4.17
65-69	274	69.6	4.20	152	85.9	6.97	122	56.2	5.09
70-74	501	150.0	6.70	230	162.6	10.72	271	140.7	8.55
75-79	735	290.5	10.72	368	373.2	19.45	367	237.7	12.41
80-84	978	611.2	19.55	361	671.3	35.33	617	580.8	23.38
85+	1,909	1413.9	32.36	461	1327.9	61.85	1,448	1443.7	37.94
Total deaths	4,852			1,792			3,060		
Connecticut									
. Crude rate		55.0	-		41.9	-		67.3	-
. AAMR (1940 Std. mil.)		20.5	0.34		22.2	0.55		19.2	0.44
. AAMR (1970 Std. mil.)		34.1	0.50		36.6	0.87		32.3	0.62
. AAMR (2000 Std. mil.)		53.0	0.76		55.9	1.37		50.7	0.92
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		28.1	-		31.0	-		26.1	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Atherosclerosis (440)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.17	0	0.0	0.00	1	0.3	0.34
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	5	0.9	0.41	2	0.7	0.53	3	1.1	0.63
50-54	4	0.9	0.47	1	0.5	0.48	3	1.4	0.79
55-59	7	1.7	0.66	7	3.6	1.36	0	0.0	0.00
60-64	13	3.1	0.86	9	4.5	1.51	4	1.8	0.91
65-69	26	6.6	1.29	14	7.9	2.11	12	5.5	1.60
70-74	48	14.4	2.07	28	19.8	3.74	20	10.4	2.32
75-79	61	24.1	3.09	34	34.5	5.91	27	17.5	3.37
80-84	79	49.4	5.55	31	57.6	10.35	48	45.2	6.52
85+	355	262.9	13.95	81	233.3	25.92	274	273.2	16.50
Total deaths	599			207			392		
Connecticut									
. Crude rate		6.8	-		4.8	-		8.6	-
. AAMR (1940 Std. mil.)		2.1	0.10		2.4	0.18		1.8	0.12
. AAMR (1970 Std. mil.)		3.9	0.16		4.3	0.30		3.6	0.19
. AAMR (2000 Std. mil.)		6.6	0.27		6.9	0.50		6.3	0.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.0	-		3.4	-		2.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	13	2.2	0.60	4	1.3	0.65	9	3.1	1.03
5-9	1	0.2	0.19	0	0.0	0.00	1	0.4	0.38
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.2	0.18	0	0.0	0.00	1	0.4	0.38
20-24	7	1.1	0.41	5	1.5	0.68	2	0.6	0.44
25-29	3	0.4	0.23	3	0.8	0.46	0	0.0	0.00
30-34	8	1.0	0.36	6	1.5	0.62	2	0.5	0.36
35-39	15	2.1	0.54	9	2.5	0.84	6	1.7	0.68
40-44	17	2.5	0.61	11	3.3	1.01	6	1.8	0.72
45-49	24	4.4	0.90	14	5.2	1.40	10	3.6	1.14
50-54	26	6.1	1.19	15	7.2	1.85	11	5.0	1.52
55-59	44	10.9	1.65	31	15.9	2.85	13	6.3	1.73
60-64	79	18.9	2.13	56	28.2	3.77	23	10.5	2.18
65-69	121	30.7	2.79	69	39.0	4.69	52	24.0	3.32
70-74	240	71.9	4.64	153	108.2	8.74	87	45.2	4.84
75-79	393	155.3	7.84	204	206.9	14.48	189	122.4	8.90
80-84	590	368.7	15.18	260	483.5	29.99	330	310.6	17.10
85+	1,584	1173.2	29.48	518	1492.1	65.56	1,066	1062.8	32.55
Total deaths	3,166			1,358			1,808		
Connecticut									
. Crude rate		35.9	-		31.7	-		39.8	-
. AAMR (1940 Std. mil.)		12.0	0.25		15.8	0.46		9.5	0.29
. AAMR (1970 Std. mil.)		21.4	0.39		28.1	0.76		17.6	0.44
. AAMR (2000 Std. mil.)		35.0	0.62		45.6	1.29		29.5	0.70
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.7	-		18.2	-		10.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.5	0.29	3	1.0	0.56	0	0.0	0.00
5-9	1	0.2	0.19	0	0.0	0.00	1	0.4	0.38
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.2	0.18	1	0.4	0.36	0	0.0	0.00
20-24	1	0.2	0.15	1	0.3	0.30	0	0.0	0.00
25-29	3	0.4	0.23	2	0.5	0.38	1	0.3	0.27
30-34	4	0.5	0.25	2	0.5	0.36	2	0.5	0.36
35-39	4	0.6	0.28	0	0.0	0.00	4	1.1	0.55
40-44	8	1.2	0.42	4	1.2	0.61	4	1.2	0.58
45-49	15	2.8	0.71	8	3.0	1.06	7	2.5	0.96
50-54	31	7.2	1.30	13	6.2	1.72	18	8.2	1.94
55-59	84	20.8	2.27	43	22.0	3.36	41	19.7	3.08
60-64	173	41.4	3.14	87	43.9	4.70	86	39.1	4.22
65-69	343	87.1	4.70	189	106.8	7.77	154	71.0	5.72
70-74	532	159.3	6.91	282	199.3	11.87	250	129.8	8.21
75-79	604	238.7	9.71	316	320.5	18.03	288	186.5	10.99
80-84	503	314.4	14.02	275	511.4	30.84	228	214.6	14.21
85+	548	405.9	17.34	263	757.6	46.71	285	284.2	16.83
Total deaths	2,858			1,489			1,369		
Connecticut									
. Crude rate		32.4	-		34.8	-		30.1	-
. AAMR (1940 Std. mil.)		14.6	0.30		19.1	0.52		11.9	0.37
. AAMR (1970 Std. mil.)		21.9	0.42		30.2	0.78		17.4	0.49
. AAMR (2000 Std. mil.)		29.9	0.56		43.2	1.17		23.3	0.63
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		19.5	-		26.9	-		14.8	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.17	1	0.3	0.33	0	0.0	0.00
5-9	1	0.2	0.19	0	0.0	0.00	1	0.4	0.38
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.2	0.18	1	0.4	0.36	0	0.0	0.00
20-24	1	0.2	0.15	1	0.3	0.30	0	0.0	0.00
25-29	2	0.3	0.19	2	0.5	0.38	0	0.0	0.00
30-34	2	0.3	0.18	1	0.3	0.25	1	0.3	0.25
35-39	2	0.3	0.20	0	0.0	0.00	2	0.6	0.39
40-44	2	0.3	0.21	0	0.0	0.00	2	0.6	0.41
45-49	5	0.9	0.41	5	1.9	0.84	0	0.0	0.00
50-54	5	1.2	0.52	3	1.4	0.83	2	0.9	0.65
55-59	10	2.5	0.78	3	1.5	0.89	7	3.4	1.27
60-64	4	1.0	0.48	3	1.5	0.87	1	0.5	0.45
65-69	14	3.6	0.95	3	1.7	0.98	11	5.1	1.53
70-74	15	4.5	1.16	3	2.1	1.22	12	6.2	1.80
75-79	14	5.5	1.48	5	5.1	2.27	9	5.8	1.94
80-84	11	6.9	2.07	2	3.7	2.63	9	8.5	2.82
85+	21	15.6	3.39	5	14.4	6.44	16	16.0	3.99
Total deaths	111			38			73		
Connecticut									
. Crude rate		1.3	-		0.9	-		1.6	-
. AAMR (1940 Std. mil.)		0.7	0.08		0.7	0.12		0.8	0.11
. AAMR (1970 Std. mil.)		0.9	0.09		0.8	0.14		1.0	0.13
. AAMR (2000 Std. mil.)		1.2	0.12		1.0	0.17		1.3	0.16
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.4	-		1.3	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis, nephrotic syndrome/nephrosis (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	0.7	0.33	3	1.0	0.56	1	0.3	0.34
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.2	0.20	0	0.0	0.00	1	0.4	0.41
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.15	1	0.3	0.30	0	0.0	0.00
25-29	2	0.3	0.19	1	0.3	0.27	1	0.3	0.27
30-34	3	0.4	0.22	2	0.5	0.36	1	0.3	0.25
35-39	7	1.0	0.37	4	1.1	0.56	3	0.8	0.48
40-44	3	0.4	0.26	3	0.9	0.53	0	0.0	0.00
45-49	13	2.4	0.66	8	3.0	1.06	5	1.8	0.81
50-54	10	2.3	0.74	7	3.3	1.27	3	1.4	0.79
55-59	14	3.5	0.93	10	5.1	1.62	4	1.9	0.96
60-64	64	15.3	1.91	39	19.7	3.15	25	11.4	2.27
65-69	59	15.0	1.95	23	13.0	2.71	36	16.6	2.77
70-74	98	29.3	2.96	50	35.3	5.00	48	24.9	3.60
75-79	124	49.0	4.40	78	79.1	8.96	46	29.8	4.39
80-84	146	91.2	7.55	68	126.5	15.34	78	73.4	8.31
85+	232	171.8	11.28	95	273.6	28.07	137	136.6	11.67
Total deaths	781			392			389		
Connecticut									
. Crude rate		8.8	-		9.2	-		8.6	-
. AAMR (1940 Std. mil.)		3.8	0.16		5.1	0.27		3.0	0.19
. AAMR (1970 Std. mil.)		5.8	0.21		8.1	0.41		4.5	0.25
. AAMR (2000 Std. mil.)		8.4	0.30		11.9	0.63		6.5	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.5	-		5.7	-		3.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----End stage renal disease (ESRD) (581-588,591,593.8,593.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	0.7	0.33	3	1.0	0.56	1	0.3	0.34
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.2	0.20	0	0.0	0.00	1	0.4	0.41
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.15	1	0.3	0.30	0	0.0	0.00
25-29	2	0.3	0.19	1	0.3	0.27	1	0.3	0.27
30-34	2	0.3	0.18	1	0.3	0.25	1	0.3	0.25
35-39	7	1.0	0.37	4	1.1	0.56	3	0.8	0.48
40-44	4	0.6	0.30	4	1.2	0.61	0	0.0	0.00
45-49	13	2.4	0.66	8	3.0	1.06	5	1.8	0.81
50-54	10	2.3	0.74	7	3.3	1.27	3	1.4	0.79
55-59	16	4.0	0.99	10	5.1	1.62	6	2.9	1.18
60-64	68	16.3	1.97	40	20.2	3.19	28	12.7	2.41
65-69	66	16.8	2.06	30	17.0	3.10	36	16.6	2.77
70-74	108	32.3	3.11	55	38.9	5.24	53	27.5	3.78
75-79	138	54.5	4.64	82	83.2	9.18	56	36.3	4.85
80-84	151	94.4	7.68	71	132.0	15.67	80	75.3	8.42
85+	259	191.8	11.92	105	302.4	29.52	154	153.5	12.37
Total deaths	850			422			428		
Connecticut									
. Crude rate		9.6	-		9.9	-		9.4	-
. AAMR (1940 Std. mil.)		4.1	0.16		5.4	0.28		3.2	0.19
. AAMR (1970 Std. mil.)		6.3	0.22		8.7	0.42		5.0	0.26
. AAMR (2000 Std. mil.)		9.1	0.31		12.9	0.65		7.2	0.35
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.1	-		6.5	-		4.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	54	9.0	1.23	33	10.7	1.87	21	7.2	1.58
5-9	25	4.6	0.93	17	6.2	1.50	8	3.1	1.08
10-14	33	6.6	1.15	23	9.0	1.87	10	4.1	1.29
15-19	191	35.1	2.54	148	52.9	4.35	43	16.2	2.47
20-24	255	39.2	2.46	200	60.7	4.29	55	17.1	2.31
25-29	186	25.0	1.83	146	39.0	3.23	40	10.8	1.71
30-34	164	20.7	1.61	131	33.0	2.89	33	8.3	1.45
35-39	125	17.4	1.55	101	28.2	2.81	24	6.6	1.35
40-44	112	16.7	1.58	89	27.0	2.86	23	6.7	1.40
45-49	79	14.5	1.63	56	20.9	2.80	23	8.3	1.73
50-54	66	15.4	1.90	48	23.0	3.31	18	8.2	1.94
55-59	90	22.3	2.35	66	33.8	4.16	24	11.5	2.36
60-64	92	22.0	2.29	67	33.8	4.13	25	11.4	2.27
65-69	120	30.5	2.78	71	40.1	4.76	49	22.6	3.23
70-74	152	45.5	3.69	87	61.5	6.59	65	33.8	4.19
75-79	163	64.4	5.05	84	85.2	9.29	79	51.2	5.76
80-84	180	112.5	8.39	79	146.9	16.53	101	95.1	9.46
85+	336	248.9	13.58	125	360.1	32.20	211	210.4	14.48
Total deaths	2,423			1,571			852		
Connecticut									
. Crude rate		27.4	-		36.7	-		18.7	-
. AAMR (1940 Std. mil.)		21.9	0.50		32.7	0.87		11.6	0.50
. AAMR (1970 Std. mil.)		23.4	0.51		34.4	0.90		13.4	0.52
. AAMR (2000 Std. mil.)		26.5	0.54		38.8	1.02		16.0	0.56
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		33.8	-		49.4	-		18.9	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	0.7	0.33	3	1.0	0.56	1	0.3	0.34
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	0.6	0.35	3	1.2	0.68	0	0.0	0.00
15-19	4	0.7	0.37	4	1.4	0.71	0	0.0	0.00
20-24	2	0.3	0.22	1	0.3	0.30	1	0.3	0.31
25-29	7	0.9	0.36	5	1.3	0.60	2	0.5	0.38
30-34	3	0.4	0.22	3	0.8	0.44	0	0.0	0.00
35-39	5	0.7	0.31	5	1.4	0.63	0	0.0	0.00
40-44	8	1.2	0.42	7	2.1	0.80	1	0.3	0.29
45-49	5	0.9	0.41	4	1.5	0.75	1	0.4	0.36
50-54	4	0.9	0.47	2	1.0	0.68	2	0.9	0.65
55-59	8	2.0	0.70	6	3.1	1.26	2	1.0	0.68
60-64	26	6.2	1.22	17	8.6	2.08	9	4.1	1.36
65-69	29	7.4	1.37	19	10.7	2.46	10	4.6	1.46
70-74	42	12.6	1.94	27	19.1	3.67	15	7.8	2.01
75-79	55	21.7	2.93	22	22.3	4.76	33	21.4	3.72
80-84	81	50.6	5.62	33	61.4	10.68	48	45.2	6.52
85+	211	156.3	10.76	61	175.7	22.50	150	149.6	12.21
Total deaths	497			222			275		
Connecticut									
. Crude rate		5.6	-		5.2	-		6.0	-
. AAMR (1940 Std. mil.)		2.3	0.13		3.1	0.23		1.6	0.13
. AAMR (1970 Std. mil.)		3.6	0.17		4.6	0.32		2.8	0.18
. AAMR (2000 Std. mil.)		5.4	0.24		6.7	0.48		4.5	0.27
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.7	-		3.6	-		1.6	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	11	1.8	0.56	5	1.6	0.73	6	2.1	0.84
5-9	4	0.7	0.37	4	1.5	0.73	0	0.0	0.00
10-14	3	0.6	0.35	3	1.2	0.68	0	0.0	0.00
15-19	7	1.3	0.49	6	2.1	0.87	1	0.4	0.38
20-24	10	1.5	0.49	10	3.0	0.96	0	0.0	0.00
25-29	7	0.9	0.36	6	1.6	0.66	1	0.3	0.27
30-34	13	1.6	0.45	12	3.0	0.87	1	0.3	0.25
35-39	9	1.3	0.42	9	2.5	0.84	0	0.0	0.00
40-44	9	1.3	0.45	8	2.4	0.86	1	0.3	0.29
45-49	6	1.1	0.45	6	2.2	0.92	0	0.0	0.00
50-54	7	1.6	0.62	6	2.9	1.17	1	0.5	0.46
55-59	5	1.2	0.55	4	2.1	1.03	1	0.5	0.48
60-64	4	1.0	0.48	4	2.0	1.01	0	0.0	0.00
65-69	4	1.0	0.51	4	2.3	1.13	0	0.0	0.00
70-74	6	1.8	0.73	3	2.1	1.22	3	1.6	0.90
75-79	3	1.2	0.68	2	2.0	1.43	1	0.6	0.65
80-84	3	1.9	1.08	1	1.9	1.86	2	1.9	1.33
85+	6	4.4	1.81	3	8.6	4.99	3	3.0	1.73
Total deaths	117			96			21		
Connecticut									
. Crude rate		1.3	-		2.2	-		0.5	-
. AAMR (1940 Std. mil.)		1.3	0.12		2.2	0.23		0.4	0.09
. AAMR (1970 Std. mil.)		1.3	0.12		2.1	0.23		0.4	0.10
. AAMR (2000 Std. mil.)		1.3	0.12		2.3	0.24		0.4	0.10
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		3.2	-		0.6	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	5	1.0	0.45	4	1.6	0.78	1	0.4	0.41
15-19	49	9.0	1.28	41	14.6	2.29	8	3.0	1.07
20-24	88	13.5	1.44	83	25.2	2.77	5	1.6	0.70
25-29	94	12.6	1.30	76	20.3	2.33	18	4.9	1.15
30-34	95	12.0	1.23	75	18.9	2.18	20	5.0	1.13
35-39	79	11.0	1.24	63	17.6	2.22	16	4.4	1.11
40-44	74	11.0	1.28	62	18.8	2.39	12	3.5	1.01
45-49	70	12.9	1.54	57	21.3	2.82	13	4.7	1.30
50-54	46	10.8	1.59	31	14.8	2.66	15	6.9	1.77
55-59	45	11.2	1.66	31	15.9	2.85	14	6.7	1.80
60-64	55	13.2	1.77	44	22.2	3.35	11	5.0	1.51
65-69	40	10.2	1.61	34	19.2	3.30	6	2.8	1.13
70-74	36	10.8	1.80	30	21.2	3.87	6	3.1	1.27
75-79	31	12.3	2.20	22	22.3	4.76	9	5.8	1.94
80-84	30	18.7	3.42	24	44.6	9.11	6	5.6	2.31
85+	21	15.6	3.39	16	46.1	11.52	5	5.0	2.23
Total deaths	858			693			165		
Connecticut									
. Crude rate		9.7	-		16.2	-		3.6	-
. AAMR (1940 Std. mil.)		8.8	0.32		14.7	0.58		3.2	0.27
. AAMR (1970 Std. mil.)		8.5	0.30		14.3	0.56		3.1	0.25
. AAMR (2000 Std. mil.)		9.3	0.32		16.0	0.62		3.5	0.28
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.4	-		18.8	-		4.6	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	8	1.3	0.47	6	2.0	0.80	2	0.7	0.49
5-9	4	0.7	0.37	2	0.7	0.51	2	0.8	0.54
10-14	5	1.0	0.45	3	1.2	0.68	2	0.8	0.58
15-19	36	6.6	1.10	30	10.7	1.96	6	2.3	0.92
20-24	47	7.2	1.05	37	11.2	1.85	10	3.1	0.99
25-29	40	5.4	0.85	32	8.6	1.51	8	2.2	0.77
30-34	44	5.5	0.84	27	6.8	1.31	17	4.3	1.04
35-39	31	4.3	0.77	21	5.9	1.28	10	2.8	0.87
40-44	18	2.7	0.63	11	3.3	1.01	7	2.0	0.77
45-49	17	3.1	0.76	14	5.2	1.40	3	1.1	0.63
50-54	9	2.1	0.70	6	2.9	1.17	3	1.4	0.79
55-59	6	1.5	0.61	4	2.1	1.03	2	1.0	0.68
60-64	12	2.9	0.83	9	4.5	1.51	3	1.4	0.79
65-69	9	2.3	0.76	7	4.0	1.50	2	0.9	0.65
70-74	14	4.2	1.12	7	4.9	1.87	7	3.6	1.37
75-79	2	0.8	0.56	1	1.0	1.01	1	0.6	0.65
80-84	4	2.5	1.25	3	5.6	3.22	1	0.9	0.94
85+	3	2.2	1.28	0	0.0	0.00	3	3.0	1.73
Total deaths	309			220			89		
Connecticut									
. Crude rate		3.5	-		5.1	-		2.0	-
. AAMR (1940 Std. mil.)		3.5	0.21		5.2	0.36		1.9	0.21
. AAMR (1970 Std. mil.)		3.3	0.20		4.8	0.34		1.8	0.20
. AAMR (2000 Std. mil.)		3.3	0.19		4.8	0.33		1.9	0.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		9.4	-		14.6	-		4.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide (E960-E969)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	8	1.3	0.47	6	2.0	0.80	2	0.7	0.49
5-9	4	0.7	0.37	2	0.7	0.51	2	0.8	0.54
10-14	5	1.0	0.45	3	1.2	0.68	2	0.8	0.58
15-19	36	6.6	1.10	30	10.7	1.96	6	2.3	0.92
20-24	47	7.2	1.05	37	11.2	1.85	10	3.1	0.99
25-29	40	5.4	0.85	32	8.6	1.51	8	2.2	0.77
30-34	44	5.5	0.84	27	6.8	1.31	17	4.3	1.04
35-39	30	4.2	0.76	21	5.9	1.28	9	2.5	0.83
40-44	18	2.7	0.63	11	3.3	1.01	7	2.0	0.77
45-49	17	3.1	0.76	14	5.2	1.40	3	1.1	0.63
50-54	9	2.1	0.70	6	2.9	1.17	3	1.4	0.79
55-59	6	1.5	0.61	4	2.1	1.03	2	1.0	0.68
60-64	12	2.9	0.83	9	4.5	1.51	3	1.4	0.79
65-69	9	2.3	0.76	7	4.0	1.50	2	0.9	0.65
70-74	14	4.2	1.12	7	4.9	1.87	7	3.6	1.37
75-79	2	0.8	0.56	1	1.0	1.01	1	0.6	0.65
80-84	4	2.5	1.25	3	5.6	3.22	1	0.9	0.94
85+	3	2.2	1.28	0	0.0	0.00	3	3.0	1.73
Total deaths	308			220			88		
Connecticut									
. Crude rate		3.5	-		5.1	-		1.9	-
. AAMR (1940 Std. mil.)		3.5	0.21		5.2	0.36		1.9	0.21
. AAMR (1970 Std. mil.)		3.3	0.20		4.8	0.34		1.7	0.20
. AAMR (2000 Std. mil.)		3.3	0.19		4.8	0.33		1.8	0.20
. HP 2000 (1940 Std. mil.)		5.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		9.3	-		14.3	-		4.2	-
. HP 2000 (1940 Std. mil.)		7.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All injury (E800-E999)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	63	10.5	1.33	40	13.0	2.06	23	7.9	1.65
5-9	30	5.6	1.02	20	7.3	1.62	10	3.8	1.21
10-14	44	8.8	1.32	31	12.1	2.17	13	5.3	1.47
15-19	277	50.8	3.05	220	78.6	5.30	57	21.5	2.85
20-24	394	60.6	3.05	324	98.4	5.46	70	21.8	2.61
25-29	331	44.5	2.45	261	69.8	4.32	70	18.9	2.26
30-34	319	40.2	2.25	242	61.1	3.92	77	19.4	2.21
35-39	252	35.0	2.21	200	55.9	3.95	52	14.4	1.99
40-44	220	32.7	2.21	174	52.8	4.00	46	13.4	1.98
45-49	171	31.4	2.40	131	49.0	4.28	40	14.5	2.29
50-54	123	28.8	2.59	85	40.7	4.41	38	17.4	2.82
55-59	141	35.0	2.95	101	51.8	5.15	40	19.2	3.04
60-64	163	39.0	3.05	123	62.0	5.59	40	18.2	2.88
65-69	170	43.2	3.31	113	63.9	6.01	57	26.3	3.48
70-74	203	60.8	4.27	125	88.4	7.90	78	40.5	4.59
75-79	197	77.9	5.55	108	109.5	10.54	89	57.6	6.11
80-84	214	133.7	9.14	106	197.1	19.15	108	101.7	9.78
85+	361	267.4	14.07	141	406.1	34.20	220	219.3	14.79
Total deaths	3,673			2,545			1,128		
Connecticut									
. Crude rate		41.6	-		59.4	-		24.8	-
. AAMR (1940 Std. mil.)		35.1	0.64		53.9	1.12		17.2	0.61
. AAMR (1970 Std. mil.)		35.9	0.63		54.8	1.12		18.7	0.62
. AAMR (2000 Std. mil.)		40.0	0.67		61.0	1.25		21.8	0.66
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		55.6	-		84.4	-		28.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	16	2.7	0.67	12	3.9	1.13	4	1.4	0.69
5-9	14	2.6	0.70	9	3.3	1.09	5	1.9	0.85
10-14	17	3.4	0.82	11	4.3	1.29	6	2.4	1.00
15-19	149	27.3	2.24	112	40.0	3.78	37	14.0	2.30
20-24	199	30.6	2.17	155	47.1	3.78	44	13.7	2.07
25-29	130	17.5	1.53	98	26.2	2.65	32	8.7	1.53
30-34	101	12.7	1.27	76	19.2	2.20	25	6.3	1.26
35-39	56	7.8	1.04	44	12.3	1.85	12	3.3	0.96
40-44	47	7.0	1.02	33	10.0	1.74	14	4.1	1.09
45-49	36	6.6	1.10	22	8.2	1.75	14	5.1	1.35
50-54	21	4.9	1.07	17	8.1	1.97	4	1.8	0.92
55-59	35	8.7	1.47	25	12.8	2.56	10	4.8	1.52
60-64	25	6.0	1.20	18	9.1	2.14	7	3.2	1.20
65-69	40	10.2	1.61	21	11.9	2.59	19	8.8	2.01
70-74	37	11.1	1.82	17	12.0	2.91	20	10.4	2.32
75-79	51	20.2	2.82	29	29.4	5.46	22	14.2	3.04
80-84	33	20.6	3.59	13	24.2	6.71	20	18.8	4.21
85+	22	16.3	3.47	7	20.2	7.62	15	15.0	3.86
Total deaths	1,029			719			310		
Connecticut									
. Crude rate		11.7	-		16.8	-		6.8	-
. AAMR (1940 Std. mil.)		11.7	0.38		17.1	0.66		6.2	0.39
. AAMR (1970 Std. mil.)		11.3	0.37		16.4	0.64		6.3	0.39
. AAMR (2000 Std. mil.)		11.2	0.35		16.1	0.61		6.4	0.37
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		18.6	-		26.1	-		10.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	0.7	0.33	3	1.0	0.56	1	0.3	0.34
5-9	1	0.2	0.19	1	0.4	0.36	0	0.0	0.00
10-14	3	0.6	0.35	3	1.2	0.68	0	0.0	0.00
15-19	3	0.6	0.32	3	1.1	0.62	0	0.0	0.00
20-24	3	0.5	0.27	3	0.9	0.53	0	0.0	0.00
25-29	7	0.9	0.36	5	1.3	0.60	2	0.5	0.38
30-34	8	1.0	0.36	6	1.5	0.62	2	0.5	0.36
35-39	6	0.8	0.34	6	1.7	0.68	0	0.0	0.00
40-44	6	0.9	0.36	6	1.8	0.74	0	0.0	0.00
45-49	6	1.1	0.45	4	1.5	0.75	2	0.7	0.51
50-54	5	1.2	0.52	2	1.0	0.68	3	1.4	0.79
55-59	9	2.2	0.74	7	3.6	1.36	2	1.0	0.68
60-64	26	6.2	1.22	17	8.6	2.08	9	4.1	1.36
65-69	27	6.9	1.32	17	9.6	2.33	10	4.6	1.46
70-74	37	11.1	1.82	24	17.0	3.46	13	6.8	1.87
75-79	41	16.2	2.53	15	15.2	3.93	26	16.8	3.30
80-84	62	38.7	4.92	24	44.6	9.11	38	35.8	5.80
85+	156	115.5	9.25	47	135.4	19.75	109	108.7	10.41
Total deaths	410			193			217		
Connecticut									
. Crude rate		4.6	-		4.5	-		4.8	-
. AAMR (1940 Std. mil.)		2.1	0.13		2.9	0.23		1.4	0.12
. AAMR (1970 Std. mil.)		3.1	0.16		4.0	0.30		2.3	0.17
. AAMR (2000 Std. mil.)		4.5	0.22		5.6	0.43		3.6	0.25
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.6	-		3.6	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.2	0.19	1	0.4	0.36	0	0.0	0.00
10-14	9	1.8	0.60	7	2.7	1.03	2	0.8	0.58
15-19	61	11.2	1.43	54	19.3	2.62	7	2.6	1.00
20-24	82	12.6	1.39	75	22.8	2.63	7	2.2	0.82
25-29	77	10.4	1.18	65	17.4	2.16	12	3.2	0.94
30-34	66	8.3	1.02	49	12.4	1.77	17	4.3	1.04
35-39	58	8.1	1.06	46	12.9	1.90	12	3.3	0.96
40-44	49	7.3	1.04	39	11.8	1.90	10	2.9	0.92
45-49	40	7.4	1.16	33	12.3	2.15	7	2.5	0.96
50-54	22	5.1	1.10	17	8.1	1.97	5	2.3	1.02
55-59	18	4.5	1.05	16	8.2	2.05	2	1.0	0.68
60-64	27	6.5	1.24	26	13.1	2.57	1	0.5	0.45
65-69	26	6.6	1.29	23	13.0	2.71	3	1.4	0.80
70-74	22	6.6	1.40	19	13.4	3.08	3	1.6	0.90
75-79	11	4.3	1.31	10	10.1	3.21	1	0.6	0.65
80-84	17	10.6	2.58	15	27.9	7.20	2	1.9	1.33
85+	9	6.7	2.22	8	23.0	8.15	1	1.0	1.00
Total deaths	595			503			92		
Connecticut									
. Crude rate		6.7	-		11.7	-		2.0	-
. AAMR (1940 Std. mil.)		6.5	0.28		11.3	0.52		2.0	0.22
. AAMR (1970 Std. mil.)		6.2	0.27		10.8	0.50		1.8	0.20
. AAMR (2000 Std. mil.)		6.5	0.27		11.4	0.52		2.0	0.21
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.6	-		23.7	-		4.1	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	0.4	0.28	2	0.8	0.55	0	0.0	0.00
15-19	25	4.6	0.92	22	7.9	1.68	3	1.1	0.65
20-24	37	5.7	0.94	37	11.2	1.85	0	0.0	0.00
25-29	45	6.1	0.90	37	9.9	1.63	8	2.2	0.77
30-34	36	4.5	0.76	28	7.1	1.33	8	2.0	0.71
35-39	37	5.1	0.85	32	8.9	1.58	5	1.4	0.62
40-44	35	5.2	0.88	29	8.8	1.63	6	1.8	0.72
45-49	28	5.1	0.97	23	8.6	1.79	5	1.8	0.81
50-54	16	3.7	0.94	13	6.2	1.72	3	1.4	0.79
55-59	15	3.7	0.96	13	6.7	1.85	2	1.0	0.68
60-64	22	5.3	1.12	22	11.1	2.37	0	0.0	0.00
65-69	23	5.8	1.22	21	11.9	2.59	2	0.9	0.65
70-74	16	4.8	1.20	16	11.3	2.83	0	0.0	0.00
75-79	10	4.0	1.25	9	9.1	3.04	1	0.6	0.65
80-84	13	8.1	2.25	12	22.3	6.44	1	0.9	0.94
85+	8	5.9	2.09	8	23.0	8.15	0	0.0	0.00
Total deaths	368			324			44		
Connecticut									
. Crude rate		4.2	-		7.6	-		1.0	-
. AAMR (1940 Std. mil.)		3.8	0.21		6.9	0.39		0.9	0.15
. AAMR (1970 Std. mil.)		3.6	0.20		6.7	0.39		0.8	0.13
. AAMR (2000 Std. mil.)		4.0	0.21		7.5	0.43		0.9	0.15
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.8	-		12.0	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.2	0.19	1	0.4	0.36	0	0.0	0.00
10-14	5	1.0	0.45	3	1.2	0.68	2	0.8	0.58
15-19	30	5.5	1.01	26	9.3	1.82	4	1.5	0.75
20-24	38	5.8	0.95	32	9.7	1.72	6	1.9	0.76
25-29	28	3.8	0.71	24	6.4	1.31	4	1.1	0.54
30-34	24	3.0	0.62	17	4.3	1.04	7	1.8	0.67
35-39	20	2.8	0.62	14	3.9	1.05	6	1.7	0.68
40-44	10	1.5	0.47	6	1.8	0.74	4	1.2	0.58
45-49	10	1.8	0.58	8	3.0	1.06	2	0.7	0.51
50-54	6	1.4	0.57	4	1.9	0.96	2	0.9	0.65
55-59	2	0.5	0.35	2	1.0	0.72	0	0.0	0.00
60-64	5	1.2	0.53	4	2.0	1.01	1	0.5	0.45
65-69	3	0.8	0.44	2	1.1	0.80	1	0.5	0.46
70-74	6	1.8	0.73	3	2.1	1.22	3	1.6	0.90
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	4	2.5	1.25	3	5.6	3.22	1	0.9	0.94
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	192			149			43		
Connecticut									
. Crude rate		2.2	-		3.5	-		0.9	-
. AAMR (1940 Std. mil.)		2.3	0.17		3.7	0.31		1.0	0.15
. AAMR (1970 Std. mil.)		2.1	0.16		3.4	0.29		0.9	0.15
. AAMR (2000 Std. mil.)		2.1	0.15		3.3	0.27		0.9	0.14
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.1	-		9.9	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	4	0.7	0.37	2	0.7	0.51	2	0.8	0.53
20-24	7	1.1	0.41	4	1.2	0.61	3	0.9	0.54
25-29	25	3.4	0.67	20	5.3	1.20	5	1.4	0.60
30-34	32	4.0	0.71	25	6.3	1.26	7	1.8	0.67
35-39	33	4.6	0.80	25	7.0	1.40	8	2.2	0.78
40-44	21	3.1	0.68	17	5.2	1.25	4	1.2	0.58
45-49	14	2.6	0.69	9	3.4	1.12	5	1.8	0.81
50-54	6	1.4	0.57	1	0.5	0.48	5	2.3	1.02
55-59	10	2.5	0.78	4	2.1	1.03	6	2.9	1.18
60-64	5	1.2	0.53	2	1.0	0.71	3	1.4	0.79
65-69	5	1.3	0.57	3	1.7	0.98	2	0.9	0.65
70-74	7	2.1	0.79	3	2.1	1.22	4	2.1	1.04
75-79	5	2.0	0.88	2	2.0	1.43	3	1.9	1.12
80-84	2	1.2	0.88	0	0.0	0.00	2	1.9	1.33
85+	6	4.4	1.81	4	11.5	5.76	2	2.0	1.41
Total deaths	182			121			61		
Connecticut									
. Crude rate		2.1	-		2.8	-		1.3	-
. AAMR (1940 Std. mil.)		1.8	0.14		2.5	0.23		1.2	0.16
. AAMR (1970 Std. mil.)		1.6	0.12		2.2	0.20		1.1	0.15
. AAMR (2000 Std. mil.)		1.9	0.15		2.7	0.25		1.3	0.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.1	-		5.5	-		2.7	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower