

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	20	2.0	0.45	11	2.1	0.65	9	1.9	0.62
5-9	27	3.0	0.57	13	2.8	0.77	14	3.1	0.84
10-14	14	1.6	0.44	9	2.1	0.69	5	1.2	0.54
15-19	29	3.3	0.62	16	3.6	0.89	13	3.1	0.85
20-24	58	5.6	0.74	33	6.3	1.09	25	4.9	0.98
25-29	86	7.2	0.78	48	8.0	1.16	38	6.4	1.04
30-34	190	14.5	1.05	98	14.9	1.51	92	14.0	1.46
35-39	316	25.9	1.46	135	22.2	1.91	181	29.5	2.19
40-44	570	51.1	2.14	250	45.6	2.89	320	56.3	3.15
45-49	886	94.2	3.17	425	91.9	4.46	461	96.5	4.49
50-54	1,278	174.7	4.89	619	173.0	6.95	659	176.3	6.87
55-59	2,038	310.0	6.87	1,032	323.7	10.08	1,006	297.1	9.37
60-64	3,307	490.1	8.52	1,765	550.1	13.09	1,542	435.7	11.09
65-69	4,703	727.3	10.60	2,596	892.7	17.52	2,107	592.1	12.90
70-74	5,490	976.5	13.18	2,961	1237.9	22.75	2,529	783.0	15.57
75-79	5,326	1228.5	16.83	2,778	1629.3	30.91	2,548	968.6	19.19
80-84	4,221	1534.8	23.62	2,083	2229.4	48.85	2,138	1177.4	25.46
85+	4,167	1789.5	27.72	1,655	2761.7	67.89	2,512	1452.6	28.98
Total deaths	32,726			16,527			16,199		
Connecticut									
. Crude rate		223.2	-		232.4	-		214.6	-
. AAMR (1940 Std. mil.)		123.2	0.76		146.5	1.20		107.7	0.99
. AAMR (1970 Std. mil.)		163.2	0.93		201.1	1.58		139.5	1.17
. AAMR (2000 Std. mil.)		204.9	1.14		259.3	2.09		173.6	1.39
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		134.3	-		165.4	-		112.2	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Oral cavity or pharynx (140.0-149.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.1	0.08	1	0.2	0.15	0	0.0	0.00
35-39	7	0.6	0.22	6	1.0	0.40	1	0.2	0.16
40-44	8	0.7	0.25	6	1.1	0.45	2	0.4	0.25
45-49	23	2.4	0.51	20	4.3	0.97	3	0.6	0.36
50-54	40	5.5	0.86	31	8.7	1.56	9	2.4	0.80
55-59	53	8.1	1.11	36	11.3	1.88	17	5.0	1.22
60-64	76	11.3	1.29	57	17.8	2.35	19	5.4	1.23
65-69	96	14.8	1.52	65	22.4	2.77	31	8.7	1.56
70-74	75	13.3	1.54	42	17.6	2.71	33	10.2	1.78
75-79	79	18.2	2.05	43	25.2	3.85	36	13.7	2.28
80-84	49	17.8	2.55	28	30.0	5.66	21	11.6	2.52
85+	46	19.8	2.91	21	35.0	7.65	25	14.5	2.89
Total deaths	553			356			197		
Connecticut									
. Crude rate		3.8	-		5.0	-		2.6	-
. AAMR (1940 Std. mil.)		2.4	0.11		3.6	0.20		1.3	0.11
. AAMR (1970 Std. mil.)		3.0	0.13		4.5	0.24		1.7	0.13
. AAMR (2000 Std. mil.)		3.5	0.15		5.4	0.29		2.1	0.15
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.4	-		3.7	-		1.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.10	0	0.0	0.00	1	0.2	0.21
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	3	0.2	0.13	2	0.3	0.22	1	0.2	0.15
35-39	9	0.7	0.25	3	0.5	0.29	6	1.0	0.40
40-44	11	1.0	0.30	6	1.1	0.45	5	0.9	0.39
45-49	37	3.9	0.65	25	5.4	1.08	12	2.5	0.73
50-54	56	7.7	1.02	36	10.1	1.68	20	5.4	1.20
55-59	92	14.0	1.46	51	16.0	2.24	41	12.1	1.89
60-64	163	24.2	1.89	93	29.0	3.01	70	19.8	2.36
65-69	259	40.1	2.49	154	53.0	4.27	105	29.5	2.88
70-74	308	54.8	3.12	158	66.1	5.26	150	46.4	3.79
75-79	290	66.9	3.93	119	69.8	6.40	171	65.0	4.97
80-84	229	83.3	5.50	98	104.9	10.60	131	72.1	6.30
85+	258	110.8	6.90	66	110.1	13.56	192	111.0	8.01
Total deaths	1,716			811			905		
Connecticut									
. Crude rate		11.7	-		11.4	-		12.0	-
. AAMR (1940 Std. mil.)		6.1	0.16		7.3	0.27		5.1	0.20
. AAMR (1970 Std. mil.)		8.4	0.21		9.9	0.35		7.1	0.25
. AAMR (2000 Std. mil.)		10.7	0.26		12.4	0.45		9.3	0.31
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.4	-		7.5	-		5.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung and other respiratory cancer (160-165)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.1	0.12	1	0.2	0.23	0	0.0	0.00
15-19	1	0.1	0.11	1	0.2	0.22	0	0.0	0.00
20-24	1	0.1	0.10	0	0.0	0.00	1	0.2	0.20
25-29	2	0.2	0.12	1	0.2	0.17	1	0.2	0.17
30-34	15	1.1	0.30	8	1.2	0.43	7	1.1	0.40
35-39	31	2.5	0.46	21	3.5	0.75	10	1.6	0.52
40-44	107	9.6	0.93	65	11.9	1.47	42	7.4	1.14
45-49	248	26.4	1.68	140	30.3	2.56	108	22.6	2.18
50-54	393	53.7	2.71	218	60.9	4.13	175	46.8	3.54
55-59	683	103.9	3.98	383	120.1	6.14	300	88.6	5.12
60-64	1,079	159.9	4.87	644	200.7	7.91	435	122.9	5.89
65-69	1,570	242.8	6.13	946	325.3	10.58	624	175.3	7.02
70-74	1,643	292.3	7.21	994	415.6	13.18	649	200.9	7.89
75-79	1,432	330.3	8.73	856	502.1	17.16	576	219.0	9.12
80-84	923	335.6	11.05	575	615.4	25.66	348	191.6	10.27
85+	563	241.8	10.19	319	532.3	29.80	244	141.1	9.03
Total deaths	8,692			5,172			3,520		
Connecticut									
. Crude rate		59.3	-		72.7	-		46.6	-
. AAMR (1940 Std. mil.)		35.0	0.41		47.3	0.69		26.1	0.49
. AAMR (1970 Std. mil.)		45.2	0.50		63.2	0.88		32.8	0.58
. AAMR (2000 Std. mil.)		53.9	0.58		78.1	1.12		38.4	0.66
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		40.7	-		60.4	-		25.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.1	0.10	0	0.0	0.00	1	0.2	0.20
25-29	2	0.2	0.12	1	0.2	0.17	1	0.2	0.17
30-34	13	1.0	0.27	8	1.2	0.43	5	0.8	0.34
35-39	30	2.5	0.45	20	3.3	0.74	10	1.6	0.52
40-44	102	9.1	0.90	61	11.1	1.43	41	7.2	1.13
45-49	244	26.0	1.66	138	29.8	2.54	106	22.2	2.16
50-54	378	51.7	2.66	207	57.8	4.02	171	45.8	3.50
55-59	663	100.9	3.92	369	115.8	6.03	294	86.8	5.06
60-64	1,050	155.6	4.80	625	194.8	7.79	425	120.1	5.82
65-69	1,516	234.4	6.02	907	311.9	10.36	609	171.1	6.93
70-74	1,600	284.6	7.12	962	402.2	12.97	638	197.5	7.82
75-79	1,388	320.1	8.59	825	483.9	16.85	563	214.0	9.02
80-84	881	320.3	10.79	541	579.0	24.89	340	187.2	10.15
85+	542	232.8	10.00	308	514.0	29.29	234	135.3	8.85
Total deaths	8,410			4,972			3,438		
Connecticut									
. Crude rate		57.4	-		69.9	-		45.5	-
. AAMR (1940 Std. mil.)		33.9	0.40		45.4	0.67		25.5	0.49
. AAMR (1970 Std. mil.)		43.7	0.49		60.8	0.87		32.1	0.57
. AAMR (2000 Std. mil.)		52.1	0.57		75.1	1.09		37.5	0.65
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		39.3	-		57.9	-		25.0	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.10	-	-	-	1	0.2	0.21
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	1	0.1	0.10	-	-	-	1	0.2	0.20
25-29	8	0.7	0.24	-	-	-	8	1.4	0.48
30-34	20	1.5	0.34	-	-	-	20	3.0	0.68
35-39	73	6.0	0.70	-	-	-	73	11.9	1.39
40-44	121	10.8	0.99	-	-	-	121	21.3	1.94
45-49	141	15.0	1.26	-	-	-	141	29.5	2.49
50-54	165	22.6	1.76	-	-	-	165	44.1	3.44
55-59	224	34.1	2.28	-	-	-	224	66.2	4.42
60-64	331	49.1	2.70	-	-	-	331	93.5	5.14
65-69	360	55.7	2.93	-	-	-	360	101.2	5.33
70-74	402	71.5	3.57	-	-	-	402	124.5	6.21
75-79	374	86.3	4.46	-	-	-	374	142.2	7.35
80-84	294	106.9	6.23	-	-	-	294	161.9	9.44
85+	367	157.6	8.23	-	-	-	367	212.2	11.08
Total deaths	2,882			-			2,882		
Connecticut									
. Crude rate		-	-		-	-		38.2	-
. AAMR (1940 Std. mil.)		-	-		-	-		21.8	0.46
. AAMR (1970 Std. mil.)		-	-		-	-		26.5	0.52
. AAMR (2000 Std. mil.)		-	-		-	-		32.2	0.61
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		23.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is lower
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	2	0.2	0.12	-	-	-	2	0.3	0.24
30-34	6	0.5	0.19	-	-	-	6	0.9	0.37
35-39	13	1.1	0.30	-	-	-	13	2.1	0.59
40-44	15	1.3	0.35	-	-	-	15	2.6	0.68
45-49	21	2.2	0.49	-	-	-	21	4.4	0.96
50-54	21	2.9	0.63	-	-	-	21	5.6	1.23
55-59	8	1.2	0.43	-	-	-	8	2.4	0.84
60-64	10	1.5	0.47	-	-	-	10	2.8	0.89
65-69	19	2.9	0.67	-	-	-	19	5.3	1.22
70-74	14	2.5	0.67	-	-	-	14	4.3	1.16
75-79	24	5.5	1.13	-	-	-	24	9.1	1.86
80-84	15	5.5	1.41	-	-	-	15	8.3	2.13
85+	18	7.7	1.82	-	-	-	18	10.4	2.45
Total deaths	186			-			186		
Connecticut									
. Crude rate		-	-		-	-		2.5	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.6	0.13
. AAMR (1970 Std. mil.)		-	-		-	-		1.8	0.14
. AAMR (2000 Std. mil.)		-	-		-	-		2.2	0.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is higher
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	2	0.2	0.13	-	-	-	2	0.4	0.25
45-49	1	0.1	0.11	-	-	-	1	0.2	0.21
50-54	2	0.3	0.19	-	-	-	2	0.5	0.38
55-59	15	2.3	0.59	-	-	-	15	4.4	1.14
60-64	24	3.6	0.73	-	-	-	24	6.8	1.38
65-69	34	5.3	0.90	-	-	-	34	9.6	1.64
70-74	35	6.2	1.05	-	-	-	35	10.8	1.83
75-79	24	5.5	1.13	-	-	-	24	9.1	1.86
80-84	24	8.7	1.78	-	-	-	24	13.2	2.70
85+	33	14.2	2.47	-	-	-	33	19.1	3.32
Total deaths	194			-			194		
Connecticut									
. Crude rate		-	-		-	-		2.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	0.10
. AAMR (1970 Std. mil.)		-	-		-	-		1.6	0.13
. AAMR (2000 Std. mil.)		-	-		-	-		2.0	0.15
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	1	0.1	0.10	-	-	-	1	0.2	0.20
25-29	3	0.3	0.15	-	-	-	3	0.5	0.29
30-34	4	0.3	0.15	-	-	-	4	0.6	0.30
35-39	3	0.2	0.14	-	-	-	3	0.5	0.28
40-44	25	2.2	0.45	-	-	-	25	4.4	0.88
45-49	32	3.4	0.60	-	-	-	32	6.7	1.18
50-54	46	6.3	0.93	-	-	-	46	12.3	1.81
55-59	65	9.9	1.23	-	-	-	65	19.2	2.38
60-64	101	15.0	1.49	-	-	-	101	28.5	2.84
65-69	129	19.9	1.76	-	-	-	129	36.2	3.19
70-74	132	23.5	2.04	-	-	-	132	40.9	3.56
75-79	119	27.4	2.52	-	-	-	119	45.2	4.15
80-84	103	37.5	3.69	-	-	-	103	56.7	5.59
85+	107	46.0	4.44	-	-	-	107	61.9	5.98
Total deaths	870			-			870		
Connecticut									
. Crude rate		-	-		-	-		11.5	-
. AAMR (1940 Std. mil.)		-	-		-	-		6.3	0.24
. AAMR (1970 Std. mil.)		-	-		-	-		7.9	0.28
. AAMR (2000 Std. mil.)		-	-		-	-		9.5	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Bladder cancer (188)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	3	0.2	0.14	3	0.5	0.29	0	0.0	0.00
40-44	3	0.3	0.16	2	0.4	0.26	1	0.2	0.18
45-49	11	1.2	0.35	6	1.3	0.53	5	1.0	0.47
50-54	9	1.2	0.41	5	1.4	0.62	4	1.1	0.54
55-59	19	2.9	0.66	15	4.7	1.21	4	1.2	0.59
60-64	59	8.7	1.14	46	14.3	2.11	13	3.7	1.02
65-69	76	11.8	1.35	57	19.6	2.60	19	5.3	1.22
70-74	130	23.1	2.03	96	40.1	4.10	34	10.5	1.81
75-79	163	37.6	2.94	111	65.1	6.18	52	19.8	2.74
80-84	152	55.3	4.48	92	98.5	10.27	60	33.0	4.27
85+	185	79.4	5.84	105	175.2	17.10	80	46.3	5.17
Total deaths	810			538			272		
Connecticut									
. Crude rate		5.5	-		7.6	-		3.6	-
. AAMR (1940 Std. mil.)		2.4	0.09		4.1	0.19		1.2	0.09
. AAMR (1970 Std. mil.)		3.6	0.13		6.4	0.28		1.9	0.12
. AAMR (2000 Std. mil.)		5.1	0.18		9.2	0.41		2.7	0.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.8	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes mellitus (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.1	0.11	0	0.0	0.00	1	0.2	0.22
10-14	1	0.1	0.12	1	0.2	0.23	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.1	0.10	0	0.0	0.00	1	0.2	0.20
25-29	6	0.5	0.21	3	0.5	0.29	3	0.5	0.29
30-34	19	1.4	0.33	9	1.4	0.46	10	1.5	0.48
35-39	22	1.8	0.38	14	2.3	0.62	8	1.3	0.46
40-44	35	3.1	0.53	24	4.4	0.89	11	1.9	0.58
45-49	46	4.9	0.72	28	6.1	1.14	18	3.8	0.89
50-54	56	7.7	1.02	37	10.3	1.70	19	5.1	1.17
55-59	96	14.6	1.49	55	17.3	2.33	41	12.1	1.89
60-64	191	28.3	2.05	103	32.1	3.16	88	24.9	2.65
65-69	253	39.1	2.46	137	47.1	4.03	116	32.6	3.03
70-74	337	59.9	3.27	164	68.6	5.35	173	53.6	4.07
75-79	420	96.9	4.73	189	110.9	8.06	231	87.8	5.78
80-84	360	130.9	6.90	137	146.6	12.53	223	122.8	8.22
85+	431	185.1	8.92	98	163.5	16.52	333	192.6	10.55
Total deaths	2,275			999			1,276		
Connecticut									
. Crude rate		15.5	-		14.0	-		16.9	-
. AAMR (1940 Std. mil.)		7.6	0.18		8.8	0.29		6.6	0.22
. AAMR (1970 Std. mil.)		10.7	0.23		12.1	0.38		9.5	0.28
. AAMR (2000 Std. mil.)		14.3	0.30		15.8	0.52		12.9	0.37
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.6	-		12.3	-		10.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.10	0	0.0	0.00	1	0.2	0.21
5-9	1	0.1	0.11	0	0.0	0.00	1	0.2	0.22
10-14	1	0.1	0.12	1	0.2	0.23	0	0.0	0.00
15-19	2	0.2	0.16	0	0.0	0.00	2	0.5	0.33
20-24	2	0.2	0.14	0	0.0	0.00	2	0.4	0.28
25-29	13	1.1	0.30	9	1.5	0.50	4	0.7	0.34
30-34	38	2.9	0.47	20	3.0	0.68	18	2.7	0.65
35-39	54	4.4	0.60	31	5.1	0.92	23	3.7	0.78
40-44	85	7.6	0.83	59	10.8	1.40	26	4.6	0.90
45-49	136	14.5	1.24	83	17.9	1.97	53	11.1	1.52
50-54	194	26.5	1.90	123	34.4	3.10	71	19.0	2.25
55-59	384	58.4	2.98	221	69.3	4.66	163	48.1	3.77
60-64	796	118.0	4.18	452	140.9	6.63	344	97.2	5.24
65-69	1,251	193.5	5.47	663	228.0	8.85	588	165.2	6.81
70-74	1,699	302.2	7.33	856	357.9	12.23	843	261.0	8.99
75-79	1,970	454.4	10.24	970	568.9	18.27	1,000	380.2	12.02
80-84	1,774	645.0	15.31	715	765.2	28.62	1,059	583.2	17.92
85+	2,270	974.9	20.46	636	1061.3	42.08	1,634	944.9	23.38
Total deaths	10,671			4,839			5,832		
Connecticut									
. Crude rate		72.8	-		68.0	-		77.3	-
. AAMR (1940 Std. mil.)		33.5	0.37		40.1	0.61		28.5	0.45
. AAMR (1970 Std. mil.)		49.0	0.49		58.2	0.84		42.4	0.59
. AAMR (2000 Std. mil.)		66.5	0.65		78.6	1.17		58.4	0.77
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.1	0.11	0	0.0	0.00	1	0.2	0.22
10-14	1	0.1	0.12	1	0.2	0.23	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	3	0.2	0.13	1	0.2	0.15	2	0.3	0.22
35-39	2	0.2	0.12	1	0.2	0.16	1	0.2	0.16
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	2	0.2	0.15	1	0.2	0.22	1	0.2	0.21
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	1	0.2	0.15	0	0.0	0.00	1	0.3	0.30
60-64	4	0.6	0.30	2	0.6	0.44	2	0.6	0.40
65-69	3	0.5	0.27	1	0.3	0.34	2	0.6	0.40
70-74	3	0.5	0.31	2	0.8	0.59	1	0.3	0.31
75-79	2	0.5	0.33	2	1.2	0.83	0	0.0	0.00
80-84	3	1.1	0.63	2	2.1	1.51	1	0.6	0.55
85+	5	2.1	0.96	2	3.3	2.36	3	1.7	1.00
Total deaths	30			15			15		
Connecticut									
. Crude rate		0.2	-		0.2	-		0.2	-
. AAMR (1940 Std. mil.)		0.1	0.03		-	-		-	-
. AAMR (1970 Std. mil.)		0.2	0.03		-	-		-	-
. AAMR (2000 Std. mil.)		0.2	0.03		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.6	-		0.6	-		0.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Alzheimer's diseases (331.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	1	0.1	0.14	0	0.0	0.00	1	0.3	0.27
55-59	8	1.2	0.43	4	1.3	0.63	4	1.2	0.59
60-64	19	2.8	0.65	11	3.4	1.03	8	2.3	0.80
65-69	53	8.2	1.13	27	9.3	1.79	26	7.3	1.43
70-74	78	13.9	1.57	36	15.1	2.51	42	13.0	2.01
75-79	167	38.5	2.98	83	48.7	5.34	84	31.9	3.48
80-84	228	82.9	5.49	90	96.3	10.15	138	76.0	6.47
85+	451	193.7	9.12	122	203.6	18.43	329	190.3	10.49
Total deaths	1,005			373			632		
Connecticut									
. Crude rate		6.9	-		5.2	-		8.4	-
. AAMR (1940 Std. mil.)		2.1	0.08		2.4	0.13		1.9	0.09
. AAMR (1970 Std. mil.)		3.9	0.13		4.4	0.23		3.6	0.15
. AAMR (2000 Std. mil.)		6.4	0.20		7.2	0.38		6.0	0.24
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.1	-		2.4	-		2.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cardiovascular disease (390-459)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	66	6.6	0.81	37	7.2	1.19	29	6.0	1.11
5-9	9	1.0	0.33	3	0.6	0.37	6	1.3	0.55
10-14	10	1.2	0.37	5	1.2	0.52	5	1.2	0.54
15-19	14	1.6	0.43	11	2.4	0.74	3	0.7	0.41
20-24	26	2.5	0.49	12	2.3	0.66	14	2.7	0.73
25-29	74	6.2	0.72	42	7.0	1.09	32	5.4	0.96
30-34	144	11.0	0.91	99	15.1	1.52	45	6.9	1.02
35-39	223	18.3	1.22	165	27.2	2.11	58	9.5	1.24
40-44	448	40.1	1.90	336	61.3	3.35	112	19.7	1.86
45-49	721	76.7	2.86	543	117.4	5.04	178	37.3	2.79
50-54	1,001	136.8	4.32	706	197.3	7.43	295	78.9	4.60
55-59	1,652	251.3	6.18	1,172	367.7	10.74	480	141.8	6.47
60-64	2,939	435.6	8.03	2,006	625.2	13.96	933	263.6	8.63
65-69	4,659	720.5	10.56	2,996	1030.3	18.82	1,663	467.3	11.46
70-74	6,833	1215.4	14.70	3,958	1654.8	26.30	2,875	890.1	16.60
75-79	8,733	2014.3	21.55	4,699	2756.0	40.20	4,034	1533.5	24.14
80-84	9,879	3592.1	36.14	4,245	4543.3	69.73	5,634	3102.6	41.34
85+	19,716	8467.1	60.30	5,502	9181.2	123.78	14,214	8219.6	68.94
Total deaths	57,147			26,537			30,610		
Connecticut									
. Crude rate		389.8	-		373.1	-		405.5	-
. AAMR (1940 Std. mil.)		157.6	0.77		211.5	1.38		115.7	0.84
. AAMR (1970 Std. mil.)		247.1	1.07		321.1	1.98		193.0	1.19
. AAMR (2000 Std. mil.)		363.1	1.53		458.3	2.93		297.3	1.71
. HP 2000 (1940 Std. mil.)		84.4	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		197.2	-		260.2	-		147.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diseases of the heart (390-398,402,404-429)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	58	5.8	0.76	32	6.2	1.10	26	5.4	1.05
5-9	8	0.9	0.31	3	0.6	0.37	5	1.1	0.50
10-14	9	1.1	0.35	4	0.9	0.46	5	1.2	0.54
15-19	9	1.0	0.34	7	1.6	0.59	2	0.5	0.33
20-24	21	2.0	0.44	10	1.9	0.60	11	2.2	0.65
25-29	49	4.1	0.59	36	6.0	1.00	13	2.2	0.61
30-34	108	8.2	0.79	83	12.7	1.39	25	3.8	0.76
35-39	182	14.9	1.10	139	22.9	1.94	43	7.0	1.07
40-44	358	32.1	1.70	282	51.5	3.07	76	13.4	1.53
45-49	608	64.7	2.62	482	104.2	4.75	126	26.4	2.35
50-54	843	115.2	3.97	622	173.8	6.97	221	59.1	3.98
55-59	1,432	217.8	5.76	1,050	329.4	10.17	382	112.8	5.77
60-64	2,548	377.6	7.48	1,789	557.6	13.18	759	214.5	7.78
65-69	3,951	611.0	9.72	2,564	881.7	17.41	1,387	389.7	10.47
70-74	5,586	993.6	13.29	3,316	1386.4	24.08	2,270	702.8	14.75
75-79	6,967	1607.0	19.25	3,802	2229.9	36.16	3,165	1203.2	21.39
80-84	7,651	2781.9	31.80	3,363	3599.3	62.07	4,288	2361.4	36.06
85+	15,180	6519.1	52.91	4,337	7237.1	109.89	10,843	6270.2	60.22
Total deaths	45,568			21,921			23,647		
Connecticut									
. Crude rate		310.8	-		308.2	-		313.3	-
. AAMR (1940 Std. mil.)		128.4	0.70		177.3	1.28		90.2	0.75
. AAMR (1970 Std. mil.)		198.9	0.96		265.7	1.80		150.0	1.05
. AAMR (2000 Std. mil.)		289.2	1.36		375.1	2.64		229.7	1.51
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		157.5	-		213.9	-		112.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	0.6	0.25	1	0.2	0.19	5	1.0	0.46
5-9	1	0.1	0.11	0	0.0	0.00	1	0.2	0.22
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.1	0.11	0	0.0	0.00	1	0.2	0.24
20-24	2	0.2	0.14	0	0.0	0.00	2	0.4	0.28
25-29	18	1.5	0.36	15	2.5	0.65	3	0.5	0.29
30-34	44	3.4	0.51	39	5.9	0.95	5	0.8	0.34
35-39	96	7.9	0.80	82	13.5	1.49	14	2.3	0.61
40-44	257	23.0	1.44	207	37.8	2.63	50	8.8	1.24
45-49	456	48.5	2.27	381	82.4	4.22	75	15.7	1.81
50-54	620	84.7	3.40	480	134.1	6.12	140	37.5	3.17
55-59	1,080	164.3	5.00	818	256.6	8.97	262	77.4	4.78
60-64	1,964	291.1	6.57	1,427	444.8	11.77	537	151.7	6.55
65-69	2,983	461.3	8.45	2,000	687.8	15.38	983	276.2	8.81
70-74	4,236	753.5	11.58	2,553	1067.4	21.12	1,683	521.1	12.70
75-79	5,184	1195.7	16.61	2,916	1710.3	31.67	2,268	862.2	18.10
80-84	5,687	2067.8	27.42	2,521	2698.1	53.74	3,166	1743.5	30.99
85+	11,261	4836.1	45.57	3,208	5353.2	94.51	8,053	4656.9	51.89
Total deaths	33,896			16,648			17,248		
Connecticut									
. Crude rate		231.2	-		234.1	-		228.5	-
. AAMR (1940 Std. mil.)		95.2	0.60		134.9	1.11		64.0	0.62
. AAMR (1970 Std. mil.)		147.9	0.83		201.8	1.57		108.1	0.88
. AAMR (2000 Std. mil.)		214.9	1.17		283.6	2.29		166.9	1.28
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		125.6	-		174.7	-		87.0	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	3	0.2	0.13	2	0.3	0.22	1	0.2	0.15
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	6	0.5	0.22	6	1.1	0.45	0	0.0	0.00
45-49	10	1.1	0.34	6	1.3	0.53	4	0.8	0.42
50-54	12	1.6	0.47	8	2.2	0.79	4	1.1	0.54
55-59	18	2.7	0.65	14	4.4	1.17	4	1.2	0.59
60-64	21	3.1	0.68	13	4.1	1.12	8	2.3	0.80
65-69	47	7.3	1.06	29	10.0	1.85	18	5.1	1.19
70-74	61	10.9	1.39	24	10.0	2.05	37	11.5	1.88
75-79	86	19.8	2.14	40	23.5	3.71	46	17.5	2.58
80-84	110	40.0	3.81	39	41.7	6.68	71	39.1	4.64
85+	181	77.7	5.78	50	83.4	11.80	131	75.8	6.62
Total deaths	555			231			324		
Connecticut									
. Crude rate		3.8	-		3.2	-		4.3	-
. AAMR (1940 Std. mil.)		1.5	0.08		1.9	0.13		1.3	0.09
. AAMR (1970 Std. mil.)		2.4	0.11		2.8	0.19		2.1	0.12
. AAMR (2000 Std. mil.)		3.5	0.15		4.0	0.28		3.2	0.18
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.8	-		2.1	-		1.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Atherosclerosis (440)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.10	0	0.0	0.00	1	0.2	0.21
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.1	0.08	0	0.0	0.00	1	0.2	0.16
40-44	1	0.1	0.09	1	0.2	0.18	0	0.0	0.00
45-49	6	0.6	0.26	3	0.6	0.37	3	0.6	0.36
50-54	6	0.8	0.33	3	0.8	0.48	3	0.8	0.46
55-59	10	1.5	0.48	8	2.5	0.89	2	0.6	0.42
60-64	17	2.5	0.61	12	3.7	1.08	5	1.4	0.63
65-69	36	5.6	0.93	21	7.2	1.58	15	4.2	1.09
70-74	75	13.3	1.54	44	18.4	2.77	31	9.6	1.72
75-79	101	23.3	2.32	58	34.0	4.47	43	16.3	2.49
80-84	132	48.0	4.18	54	57.8	7.86	78	43.0	4.86
85+	582	249.9	10.36	134	223.6	19.32	448	259.1	12.24
Total deaths	968			338			630		
Connecticut									
. Crude rate		6.6	-		4.8	-		8.3	-
. AAMR (1940 Std. mil.)		1.9	0.07		2.3	0.13		1.7	0.09
. AAMR (1970 Std. mil.)		3.7	0.12		4.1	0.22		3.3	0.14
. AAMR (2000 Std. mil.)		6.3	0.20		6.6	0.37		5.9	0.24
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.0	-		3.4	-		2.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Aortic aneurysm (441)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.2	0.14	2	0.4	0.28	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.1	0.11	1	0.2	0.22	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.2	0.12	0	0.0	0.00	2	0.3	0.24
30-34	1	0.1	0.08	0	0.0	0.00	1	0.2	0.15
35-39	4	0.3	0.16	2	0.3	0.23	2	0.3	0.23
40-44	5	0.4	0.20	3	0.5	0.32	2	0.4	0.25
45-49	6	0.6	0.26	6	1.3	0.53	0	0.0	0.00
50-54	5	0.7	0.31	5	1.4	0.62	0	0.0	0.00
55-59	23	3.5	0.73	16	5.0	1.25	7	2.1	0.78
60-64	53	7.9	1.08	43	13.4	2.04	10	2.8	0.89
65-69	119	18.4	1.69	98	33.7	3.40	21	5.9	1.29
70-74	175	31.1	2.35	115	48.1	4.48	60	18.6	2.40
75-79	184	42.4	3.13	111	65.1	6.18	73	27.8	3.25
80-84	200	72.7	5.14	111	118.8	11.28	89	49.0	5.20
85+	190	81.6	5.92	75	125.2	14.45	115	66.5	6.20
Total deaths	970			588			382		
Connecticut									
. Crude rate		6.6	-		8.3	-		5.1	-
. AAMR (1940 Std. mil.)		2.9	0.10		4.7	0.20		1.6	0.10
. AAMR (1970 Std. mil.)		4.4	0.14		7.0	0.29		2.6	0.14
. AAMR (2000 Std. mil.)		6.0	0.19		9.6	0.41		3.7	0.19
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.7	-		5.9	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.3	0.17	3	0.6	0.34	0	0.0	0.00
5-9	1	0.1	0.11	0	0.0	0.00	1	0.2	0.22
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.2	0.16	1	0.2	0.22	1	0.2	0.24
20-24	5	0.5	0.22	2	0.4	0.27	3	0.6	0.34
25-29	7	0.6	0.22	3	0.5	0.29	4	0.7	0.34
30-34	5	0.4	0.17	3	0.5	0.26	2	0.3	0.22
35-39	7	0.6	0.22	2	0.3	0.23	5	0.8	0.36
40-44	14	1.3	0.34	7	1.3	0.48	7	1.2	0.47
45-49	24	2.6	0.52	13	2.8	0.78	11	2.3	0.69
50-54	45	6.2	0.92	17	4.8	1.15	28	7.5	1.42
55-59	130	19.8	1.73	65	20.4	2.53	65	19.2	2.38
60-64	292	43.3	2.53	151	47.1	3.83	141	39.8	3.36
65-69	570	88.1	3.69	326	112.1	6.21	244	68.6	4.39
70-74	890	158.3	5.31	471	196.9	9.07	419	129.7	6.34
75-79	1,047	241.5	7.46	546	320.2	13.70	501	190.5	8.51
80-84	940	341.8	11.15	496	530.9	23.84	444	244.5	11.60
85+	1,044	448.3	13.88	467	779.3	36.06	577	333.7	13.89
Total deaths	5,026			2,573			2,453		
Connecticut									
. Crude rate		34.3	-		36.2	-		32.5	-
. AAMR (1940 Std. mil.)		14.9	0.23		19.3	0.40		12.2	0.29
. AAMR (1970 Std. mil.)		22.6	0.33		30.6	0.61		18.1	0.39
. AAMR (2000 Std. mil.)		31.1	0.44		43.9	0.90		24.6	0.50
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		19.5	-		26.9	-		14.8	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.10	1	0.2	0.19	0	0.0	0.00
5-9	1	0.1	0.11	0	0.0	0.00	1	0.2	0.22
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.2	0.16	1	0.2	0.22	1	0.2	0.24
20-24	4	0.4	0.19	1	0.2	0.19	3	0.6	0.34
25-29	5	0.4	0.19	3	0.5	0.29	2	0.3	0.24
30-34	3	0.2	0.13	2	0.3	0.22	1	0.2	0.15
35-39	3	0.2	0.14	1	0.2	0.16	2	0.3	0.23
40-44	6	0.5	0.22	2	0.4	0.26	4	0.7	0.35
45-49	8	0.9	0.30	6	1.3	0.53	2	0.4	0.30
50-54	8	1.1	0.39	3	0.8	0.48	5	1.3	0.60
55-59	14	2.1	0.57	6	1.9	0.77	8	2.4	0.84
60-64	9	1.3	0.44	8	2.5	0.88	1	0.3	0.28
65-69	23	3.6	0.74	7	2.4	0.91	16	4.5	1.12
70-74	26	4.6	0.91	4	1.7	0.84	22	6.8	1.45
75-79	20	4.6	1.03	6	3.5	1.44	14	5.3	1.42
80-84	20	7.3	1.63	4	4.3	2.14	16	8.8	2.20
85+	44	18.9	2.85	10	16.7	5.28	34	19.7	3.37
Total deaths	197			65			132		
Connecticut									
. Crude rate		1.3	-		0.9	-		1.7	-
. AAMR (1940 Std. mil.)		0.8	0.06		0.7	0.09		0.8	0.09
. AAMR (1970 Std. mil.)		1.0	0.07		0.8	0.10		1.1	0.10
. AAMR (2000 Std. mil.)		1.3	0.09		1.0	0.13		1.4	0.12
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.4	-		1.3	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.1	0.10	0	0.0	0.00	1	0.2	0.20
25-29	14	1.2	0.31	8	1.3	0.47	6	1.0	0.41
30-34	40	3.0	0.48	28	4.3	0.81	12	1.8	0.53
35-39	68	5.6	0.68	47	7.7	1.13	21	3.4	0.75
40-44	94	8.4	0.87	66	12.0	1.48	28	4.9	0.93
45-49	96	10.2	1.04	71	15.4	1.82	25	5.2	1.05
50-54	136	18.6	1.59	101	28.2	2.81	35	9.4	1.58
55-59	140	21.3	1.80	101	31.7	3.15	39	11.5	1.84
60-64	165	24.5	1.90	112	34.9	3.30	53	15.0	2.06
65-69	191	29.5	2.14	108	37.1	3.57	83	23.3	2.56
70-74	175	31.1	2.35	102	42.6	4.22	73	22.6	2.65
75-79	139	32.1	2.72	73	42.8	5.01	66	25.1	3.09
80-84	87	31.6	3.39	45	48.2	7.18	42	23.1	3.57
85+	47	20.2	2.94	24	40.0	8.17	23	13.3	2.77
Total deaths	1,393			886			507		
Connecticut									
. Crude rate		9.5	-		12.5	-		6.7	-
. AAMR (1940 Std. mil.)		6.9	0.20		9.8	0.34		4.3	0.21
. AAMR (1970 Std. mil.)		7.7	0.21		11.0	0.37		4.9	0.23
. AAMR (2000 Std. mil.)		9.1	0.25		13.0	0.45		5.8	0.26
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		9.2	-		12.8	-		5.5	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis, nephrotic syndrome/nephrosis (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	0.4	0.20	3	0.6	0.34	1	0.2	0.21
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.1	0.12	0	0.0	0.00	1	0.2	0.24
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	0.2	0.14	2	0.4	0.27	0	0.0	0.00
25-29	3	0.3	0.15	2	0.3	0.24	1	0.2	0.17
30-34	6	0.5	0.19	4	0.6	0.30	2	0.3	0.22
35-39	12	1.0	0.28	6	1.0	0.40	6	1.0	0.40
40-44	5	0.4	0.20	4	0.7	0.37	1	0.2	0.18
45-49	15	1.6	0.41	10	2.2	0.68	5	1.0	0.47
50-54	21	2.9	0.63	13	3.6	1.01	8	2.1	0.76
55-59	25	3.8	0.76	16	5.0	1.25	9	2.7	0.89
60-64	83	12.3	1.35	49	15.3	2.18	34	9.6	1.65
65-69	108	16.7	1.61	52	17.9	2.48	56	15.7	2.10
70-74	163	29.0	2.27	87	36.4	3.90	76	23.5	2.70
75-79	204	47.1	3.29	123	72.1	6.50	81	30.8	3.42
80-84	271	98.5	5.99	127	135.9	12.06	144	79.3	6.61
85+	399	171.4	8.58	174	290.4	22.01	225	130.1	8.67
Total deaths	1,322			672			650		
Connecticut									
. Crude rate		9.0	-		9.4	-		8.6	-
. AAMR (1940 Std. mil.)		3.7	0.12		5.0	0.21		2.9	0.14
. AAMR (1970 Std. mil.)		5.8	0.16		8.1	0.31		4.5	0.19
. AAMR (2000 Std. mil.)		8.3	0.23		12.1	0.49		6.4	0.25
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.5	-		5.7	-		3.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	82	8.2	0.91	52	10.1	1.40	30	6.2	1.13
5-9	38	4.2	0.68	28	6.0	1.13	10	2.2	0.71
10-14	51	6.0	0.84	36	8.3	1.38	15	3.6	0.93
15-19	278	31.8	1.91	210	46.7	3.22	68	16.0	1.94
20-24	373	36.0	1.87	300	57.1	3.29	73	14.3	1.68
25-29	304	25.6	1.47	239	40.0	2.59	65	11.0	1.37
30-34	312	23.8	1.35	254	38.7	2.43	58	8.8	1.16
35-39	247	20.2	1.29	200	32.9	2.33	47	7.7	1.12
40-44	202	18.1	1.27	162	29.6	2.32	40	7.0	1.11
45-49	145	15.4	1.28	109	23.6	2.26	36	7.5	1.26
50-54	115	15.7	1.47	80	22.4	2.50	35	9.4	1.58
55-59	140	21.3	1.80	104	32.6	3.20	36	10.6	1.77
60-64	159	23.6	1.87	114	35.5	3.33	45	12.7	1.90
65-69	205	31.7	2.21	115	39.5	3.69	90	25.3	2.67
70-74	253	45.0	2.83	142	59.4	4.98	111	34.4	3.26
75-79	267	61.6	3.77	134	78.6	6.79	133	50.6	4.38
80-84	320	116.4	6.50	144	154.1	12.84	176	96.9	7.31
85+	583	250.4	10.37	205	342.1	23.89	378	218.6	11.24
Total deaths	4,074			2,628			1,446		
Connecticut									
. Crude rate		27.8	-		36.9	-		19.2	-
. AAMR (1940 Std. mil.)		21.9	0.39		32.7	0.68		11.5	0.38
. AAMR (1970 Std. mil.)		23.3	0.39		34.1	0.69		13.3	0.40
. AAMR (2000 Std. mil.)		26.6	0.42		38.8	0.78		16.1	0.43
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		33.8	-		49.4	-		18.9	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Motor vehicle accidents (E810-E825)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	27	2.7	0.52	21	4.1	0.89	6	1.2	0.50
5-9	21	2.3	0.50	15	3.2	0.83	6	1.3	0.55
10-14	22	2.6	0.55	14	3.2	0.86	8	1.9	0.68
15-19	217	24.8	1.68	161	35.8	2.82	56	13.2	1.76
20-24	288	27.8	1.64	231	43.9	2.89	57	11.2	1.48
25-29	196	16.5	1.18	148	24.8	2.04	48	8.1	1.17
30-34	171	13.0	1.00	128	19.5	1.73	43	6.5	1.00
35-39	95	7.8	0.80	75	12.3	1.43	20	3.3	0.73
40-44	76	6.8	0.78	55	10.0	1.35	21	3.7	0.81
45-49	59	6.3	0.82	40	8.6	1.37	19	4.0	0.91
50-54	49	6.7	0.96	34	9.5	1.63	15	4.0	1.04
55-59	56	8.5	1.14	41	12.9	2.01	15	4.4	1.14
60-64	52	7.7	1.07	38	11.8	1.92	14	4.0	1.06
65-69	77	11.9	1.36	36	12.4	2.06	41	11.5	1.80
70-74	72	12.8	1.51	32	13.4	2.37	40	12.4	1.96
75-79	81	18.7	2.08	41	24.0	3.76	40	15.2	2.40
80-84	67	24.4	2.98	28	30.0	5.66	39	21.5	3.44
85+	35	15.0	2.54	15	25.0	6.46	20	11.6	2.59
Total deaths	1,661			1,153			508		
Connecticut									
. Crude rate		11.3	-		16.2	-		6.7	-
. AAMR (1940 Std. mil.)		11.2	0.29		16.5	0.50		6.0	0.30
. AAMR (1970 Std. mil.)		10.9	0.28		15.8	0.49		6.1	0.29
. AAMR (2000 Std. mil.)		10.9	0.27		15.8	0.47		6.2	0.28
. HP 2000 (1940 Std. mil.)		10.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		18.8	-		26.7	-		11.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	0.5	0.22	3	0.6	0.34	2	0.4	0.29
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	4	0.5	0.24	4	0.9	0.46	0	0.0	0.00
15-19	8	0.9	0.32	7	1.6	0.59	1	0.2	0.24
20-24	4	0.4	0.19	3	0.6	0.33	1	0.2	0.20
25-29	8	0.7	0.24	6	1.0	0.41	2	0.3	0.24
30-34	7	0.5	0.20	7	1.1	0.40	0	0.0	0.00
35-39	9	0.7	0.25	8	1.3	0.47	1	0.2	0.16
40-44	13	1.2	0.32	10	1.8	0.58	3	0.5	0.30
45-49	13	1.4	0.38	11	2.4	0.72	2	0.4	0.30
50-54	11	1.5	0.45	5	1.4	0.62	6	1.6	0.66
55-59	18	2.7	0.65	14	4.4	1.17	4	1.2	0.59
60-64	41	6.1	0.95	27	8.4	1.62	14	4.0	1.06
65-69	42	6.5	1.00	28	9.6	1.82	14	3.9	1.05
70-74	65	11.6	1.43	40	16.7	2.64	25	7.7	1.55
75-79	91	21.0	2.20	41	24.0	3.76	50	19.0	2.69
80-84	131	47.6	4.16	54	57.8	7.86	77	42.4	4.83
85+	361	155.0	8.16	107	178.6	17.26	254	146.9	9.22
Total deaths	831			375			456		
Connecticut									
. Crude rate		5.7	-		5.3	-		6.0	-
. AAMR (1940 Std. mil.)		2.3	0.10		3.1	0.18		1.6	0.10
. AAMR (1970 Std. mil.)		3.6	0.13		4.6	0.24		2.7	0.14
. AAMR (2000 Std. mil.)		5.4	0.19		6.8	0.36		4.4	0.21
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.7	-		3.6	-		1.6	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	13	1.3	0.36	7	1.4	0.52	6	1.2	0.50
5-9	5	0.5	0.25	5	1.1	0.48	0	0.0	0.00
10-14	5	0.6	0.26	5	1.2	0.52	0	0.0	0.00
15-19	13	1.5	0.41	10	2.2	0.70	3	0.7	0.41
20-24	11	1.1	0.32	11	2.1	0.63	0	0.0	0.00
25-29	10	0.8	0.27	8	1.3	0.47	2	0.3	0.24
30-34	20	1.5	0.34	19	2.9	0.66	1	0.2	0.15
35-39	17	1.4	0.34	15	2.5	0.64	2	0.3	0.23
40-44	17	1.5	0.37	15	2.7	0.71	2	0.4	0.25
45-49	9	1.0	0.32	9	1.9	0.65	0	0.0	0.00
50-54	8	1.1	0.39	6	1.7	0.68	2	0.5	0.38
55-59	6	0.9	0.37	5	1.6	0.70	1	0.3	0.30
60-64	5	0.7	0.33	4	1.2	0.62	1	0.3	0.28
65-69	6	0.9	0.38	5	1.7	0.77	1	0.3	0.28
70-74	11	2.0	0.59	8	3.3	1.18	3	0.9	0.54
75-79	4	0.9	0.46	3	1.8	1.02	1	0.4	0.38
80-84	5	1.8	0.81	1	1.1	1.07	4	2.2	1.10
85+	6	2.6	1.05	3	5.0	2.89	3	1.7	1.00
Total deaths	171			139			32		
Connecticut									
. Crude rate		1.2	-		2.0	-		0.4	-
. AAMR (1940 Std. mil.)		1.1	0.09		1.9	0.16		0.4	0.07
. AAMR (1970 Std. mil.)		1.1	0.09		1.8	0.16		0.4	0.07
. AAMR (2000 Std. mil.)		1.2	0.09		2.0	0.17		0.4	0.07
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		3.2	-		0.6	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	14	1.4	0.37	7	1.4	0.52	7	1.4	0.55
5-9	5	0.5	0.25	3	0.6	0.37	2	0.4	0.32
10-14	3	0.4	0.20	2	0.5	0.33	1	0.2	0.24
15-19	5	0.6	0.26	4	0.9	0.44	1	0.2	0.24
20-24	12	1.2	0.33	8	1.5	0.54	4	0.8	0.39
25-29	3	0.3	0.15	1	0.2	0.17	2	0.3	0.24
30-34	3	0.2	0.13	1	0.2	0.15	2	0.3	0.22
35-39	5	0.4	0.18	3	0.5	0.29	2	0.3	0.23
40-44	9	0.8	0.27	6	1.1	0.45	3	0.5	0.30
45-49	5	0.5	0.24	3	0.6	0.37	2	0.4	0.30
50-54	7	1.0	0.36	3	0.8	0.48	4	1.1	0.54
55-59	9	1.4	0.46	6	1.9	0.77	3	0.9	0.51
60-64	6	0.9	0.36	3	0.9	0.54	3	0.8	0.49
65-69	11	1.7	0.51	6	2.1	0.84	5	1.4	0.63
70-74	13	2.3	0.64	7	2.9	1.11	6	1.9	0.76
75-79	5	1.2	0.52	2	1.2	0.83	3	1.1	0.66
80-84	11	4.0	1.21	5	5.4	2.39	6	3.3	1.35
85+	5	2.1	0.96	4	6.7	3.34	1	0.6	0.58
Total deaths	131			74			57		
Connecticut									
. Crude rate		0.9	-		1.0	-		0.8	-
. AAMR (1940 Std. mil.)		0.8	0.07		0.9	0.12		0.6	0.09
. AAMR (1970 Std. mil.)		0.8	0.08		1.0	0.12		0.7	0.10
. AAMR (2000 Std. mil.)		0.9	0.08		1.1	0.13		0.7	0.10
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.7	-		2.0	-		1.1	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	8	0.9	0.33	7	1.6	0.61	1	0.2	0.24
15-19	72	8.2	0.97	59	13.1	1.71	13	3.1	0.85
20-24	137	13.2	1.13	128	24.3	2.15	9	1.8	0.59
25-29	151	12.7	1.03	122	20.4	1.85	29	4.9	0.91
30-34	153	11.7	0.94	119	18.1	1.66	34	5.2	0.89
35-39	146	12.0	0.99	115	18.9	1.77	31	5.1	0.91
40-44	139	12.5	1.06	116	21.2	1.97	23	4.0	0.84
45-49	111	11.8	1.12	90	19.5	2.05	21	4.4	0.96
50-54	78	10.7	1.21	58	16.2	2.13	20	5.4	1.20
55-59	80	12.2	1.36	56	17.6	2.35	24	7.1	1.45
60-64	77	11.4	1.30	61	19.0	2.43	16	4.5	1.13
65-69	74	11.4	1.33	61	21.0	2.69	13	3.7	1.01
70-74	69	12.3	1.48	60	25.1	3.24	9	2.8	0.93
75-79	55	12.7	1.71	38	22.3	3.62	17	6.5	1.57
80-84	42	15.3	2.36	33	35.3	6.15	9	5.0	1.65
85+	31	13.3	2.39	25	41.7	8.34	6	3.5	1.42
Total deaths	1,423			1,148			275		
Connecticut									
. Crude rate		9.7	-		16.1	-		3.6	-
. AAMR (1940 Std. mil.)		8.8	0.25		14.7	0.45		3.2	0.21
. AAMR (1970 Std. mil.)		8.4	0.23		14.3	0.43		3.1	0.20
. AAMR (2000 Std. mil.)		9.3	0.25		16.0	0.48		3.5	0.21
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.4	-		18.8	-		4.6	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	12	1.2	0.35	8	1.6	0.55	4	0.8	0.41
5-9	7	0.8	0.29	3	0.6	0.37	4	0.9	0.45
10-14	9	1.1	0.35	7	1.6	0.61	2	0.5	0.34
15-19	64	7.3	0.91	54	12.0	1.63	10	2.4	0.74
20-24	86	8.3	0.90	67	12.7	1.56	19	3.7	0.86
25-29	75	6.3	0.73	57	9.5	1.26	18	3.0	0.72
30-34	68	5.2	0.63	49	7.5	1.07	19	2.9	0.66
35-39	51	4.2	0.58	33	5.4	0.95	18	2.9	0.69
40-44	26	2.3	0.46	15	2.7	0.71	11	1.9	0.58
45-49	34	3.6	0.62	28	6.1	1.14	6	1.3	0.51
50-54	15	2.1	0.53	11	3.1	0.93	4	1.1	0.54
55-59	12	1.8	0.53	8	2.5	0.89	4	1.2	0.59
60-64	17	2.5	0.61	13	4.1	1.12	4	1.1	0.57
65-69	10	1.5	0.49	7	2.4	0.91	3	0.8	0.49
70-74	19	3.4	0.78	9	3.8	1.25	10	3.1	0.98
75-79	4	0.9	0.46	2	1.2	0.83	2	0.8	0.54
80-84	7	2.5	0.96	3	3.2	1.85	4	2.2	1.10
85+	4	1.7	0.86	0	0.0	0.00	4	2.3	1.16
Total deaths	520			374			146		
Connecticut									
. Crude rate		3.5	-		5.3	-		1.9	-
. AAMR (1940 Std. mil.)		3.7	0.17		5.5	0.29		1.9	0.17
. AAMR (1970 Std. mil.)		3.4	0.16		5.1	0.28		1.8	0.16
. AAMR (2000 Std. mil.)		3.4	0.15		5.0	0.26		1.9	0.16
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		9.4	-		14.6	-		4.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	0.5	0.22	3	0.6	0.34	2	0.4	0.29
5-9	1	0.1	0.11	1	0.2	0.21	0	0.0	0.00
10-14	4	0.5	0.24	4	0.9	0.46	0	0.0	0.00
15-19	7	0.8	0.30	6	1.3	0.54	1	0.2	0.24
20-24	5	0.5	0.22	5	1.0	0.43	0	0.0	0.00
25-29	9	0.8	0.25	6	1.0	0.41	3	0.5	0.29
30-34	13	1.0	0.27	11	1.7	0.51	2	0.3	0.22
35-39	12	1.0	0.28	11	1.8	0.55	1	0.2	0.16
40-44	11	1.0	0.30	9	1.6	0.55	2	0.4	0.25
45-49	15	1.6	0.41	12	2.6	0.75	3	0.6	0.36
50-54	12	1.6	0.47	6	1.7	0.68	6	1.6	0.66
55-59	18	2.7	0.65	15	4.7	1.21	3	0.9	0.51
60-64	41	6.1	0.95	27	8.4	1.62	14	4.0	1.06
65-69	39	6.0	0.97	26	8.9	1.75	13	3.7	1.01
70-74	57	10.1	1.34	37	15.5	2.54	20	6.2	1.38
75-79	71	16.4	1.94	34	19.9	3.42	37	14.1	2.31
80-84	97	35.3	3.58	38	40.7	6.60	59	32.5	4.23
85+	247	106.1	6.75	79	131.8	14.83	168	97.2	7.50
Total deaths	664			330			334		
Connecticut									
. Crude rate		4.5	-		4.6	-		4.4	-
. AAMR (1940 Std. mil.)		2.1	0.10		3.0	0.18		1.3	0.10
. AAMR (1970 Std. mil.)		3.0	0.12		4.1	0.23		2.1	0.13
. AAMR (2000 Std. mil.)		4.3	0.17		5.7	0.33		3.3	0.18
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.6	-		3.6	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	4	0.4	0.22	2	0.4	0.30	2	0.4	0.32
10-14	14	1.6	0.44	12	2.8	0.80	2	0.5	0.34
15-19	96	11.0	1.12	87	19.3	2.07	9	2.1	0.71
20-24	150	14.5	1.18	133	25.3	2.19	17	3.3	0.81
25-29	131	11.0	0.96	107	17.9	1.73	24	4.1	0.83
30-34	103	7.8	0.77	82	12.5	1.38	21	3.2	0.70
35-39	104	8.5	0.84	81	13.3	1.48	23	3.7	0.78
40-44	78	7.0	0.79	62	11.3	1.44	16	2.8	0.70
45-49	71	7.6	0.90	60	13.0	1.67	11	2.3	0.69
50-54	45	6.2	0.92	36	10.1	1.68	9	2.4	0.80
55-59	37	5.6	0.93	30	9.4	1.72	7	2.1	0.78
60-64	41	6.1	0.95	39	12.2	1.95	2	0.6	0.40
65-69	40	6.2	0.98	36	12.4	2.06	4	1.1	0.56
70-74	43	7.6	1.17	36	15.1	2.51	7	2.2	0.82
75-79	20	4.6	1.03	19	11.1	2.56	1	0.4	0.38
80-84	22	8.0	1.71	20	21.4	4.79	2	1.1	0.78
85+	14	6.0	1.61	13	21.7	6.02	1	0.6	0.58
Total deaths	1,013			855			158		
Connecticut									
. Crude rate		6.9	-		12.0	-		2.1	-
. AAMR (1940 Std. mil.)		6.8	0.22		11.7	0.42		2.1	0.17
. AAMR (1970 Std. mil.)		6.4	0.21		11.2	0.40		1.9	0.16
. AAMR (2000 Std. mil.)		6.7	0.21		11.8	0.41		2.1	0.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.6	-		23.7	-		4.1	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	0.4	0.20	3	0.7	0.40	0	0.0	0.00
15-19	36	4.1	0.69	33	7.3	1.28	3	0.7	0.41
20-24	66	6.4	0.78	64	12.2	1.52	2	0.4	0.28
25-29	73	6.1	0.72	59	9.9	1.29	14	2.4	0.63
30-34	55	4.2	0.57	43	6.6	1.00	12	1.8	0.53
35-39	71	5.8	0.69	59	9.7	1.26	12	2.0	0.56
40-44	59	5.3	0.69	48	8.8	1.26	11	1.9	0.58
45-49	48	5.1	0.74	40	8.6	1.37	8	1.7	0.59
50-54	35	4.8	0.81	29	8.1	1.50	6	1.6	0.66
55-59	33	5.0	0.87	26	8.2	1.60	7	2.1	0.78
60-64	33	4.9	0.85	33	10.3	1.79	0	0.0	0.00
65-69	37	5.7	0.94	34	11.7	2.01	3	0.8	0.49
70-74	34	6.0	1.04	32	13.4	2.37	2	0.6	0.44
75-79	18	4.2	0.98	17	10.0	2.42	1	0.4	0.38
80-84	18	6.5	1.54	17	18.2	4.41	1	0.6	0.55
85+	13	5.6	1.55	13	21.7	6.02	0	0.0	0.00
Total deaths	632			550			82		
Connecticut									
. Crude rate		4.3	-		7.7	-		1.1	-
. AAMR (1940 Std. mil.)		4.0	0.17		7.1	0.31		1.0	0.12
. AAMR (1970 Std. mil.)		3.8	0.16		6.9	0.30		0.9	0.11
. AAMR (2000 Std. mil.)		4.1	0.17		7.7	0.33		1.1	0.12
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.8	-		12.0	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	4	0.4	0.22	2	0.4	0.30	2	0.4	0.32
10-14	9	1.1	0.35	7	1.6	0.61	2	0.5	0.34
15-19	54	6.2	0.84	48	10.7	1.54	6	1.4	0.58
20-24	70	6.8	0.81	57	10.8	1.44	13	2.6	0.71
25-29	53	4.5	0.61	43	7.2	1.10	10	1.7	0.54
30-34	42	3.2	0.49	35	5.3	0.90	7	1.1	0.40
35-39	31	2.5	0.46	21	3.5	0.75	10	1.6	0.52
40-44	14	1.3	0.34	9	1.6	0.55	5	0.9	0.39
45-49	21	2.2	0.49	18	3.9	0.92	3	0.6	0.36
50-54	10	1.4	0.43	7	2.0	0.74	3	0.8	0.46
55-59	3	0.5	0.26	3	0.9	0.54	0	0.0	0.00
60-64	7	1.0	0.39	6	1.9	0.76	1	0.3	0.28
65-69	3	0.5	0.27	2	0.7	0.49	1	0.3	0.28
70-74	8	1.4	0.50	4	1.7	0.84	4	1.2	0.62
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	4	1.5	0.73	3	3.2	1.85	1	0.6	0.55
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	333			265			68		
Connecticut									
. Crude rate		2.3	-		3.7	-		0.9	-
. AAMR (1940 Std. mil.)		2.5	0.14		4.0	0.25		1.0	0.12
. AAMR (1970 Std. mil.)		2.3	0.13		3.7	0.24		0.9	0.11
. AAMR (2000 Std. mil.)		2.2	0.12		3.6	0.22		0.9	0.11
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.1	-		9.9	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	5	0.6	0.26	3	0.7	0.40	2	0.5	0.34
15-19	23	2.6	0.55	13	2.9	0.80	10	2.4	0.74
20-24	53	5.1	0.70	46	8.7	1.29	7	1.4	0.52
25-29	87	7.3	0.79	72	12.1	1.42	15	2.5	0.66
30-34	129	9.8	0.87	105	16.0	1.56	24	3.7	0.75
35-39	122	10.0	0.90	97	16.0	1.62	25	4.1	0.81
40-44	100	9.0	0.90	80	14.6	1.63	20	3.5	0.79
45-49	69	7.3	0.88	54	11.7	1.59	15	3.1	0.81
50-54	34	4.6	0.80	21	5.9	1.28	13	3.5	0.96
55-59	29	4.4	0.82	15	4.7	1.21	14	4.1	1.11
60-64	28	4.1	0.78	17	5.3	1.29	11	3.1	0.94
65-69	23	3.6	0.74	16	5.5	1.38	7	2.0	0.74
70-74	26	4.6	0.91	17	7.1	1.72	9	2.8	0.93
75-79	22	5.1	1.08	6	3.5	1.44	16	6.1	1.52
80-84	16	5.8	1.45	9	9.6	3.21	7	3.9	1.46
85+	17	7.3	1.77	12	20.0	5.78	5	2.9	1.29
Total deaths	783			583			200		
Connecticut									
. Crude rate		5.3	-		8.2	-		2.6	-
. AAMR (1940 Std. mil.)		4.8	0.18		7.4	0.31		2.3	0.18
. AAMR (1970 Std. mil.)		4.4	0.16		6.7	0.28		2.2	0.17
. AAMR (2000 Std. mil.)		5.1	0.18		7.9	0.33		2.5	0.18
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.0	-		6.9	-		3.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989-1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.1	0.12	1	0.2	0.23	0	0.0	0.00
15-19	11	1.3	0.38	5	1.1	0.50	6	1.4	0.58
20-24	31	3.0	0.54	25	4.8	0.95	6	1.2	0.48
25-29	71	6.0	0.71	58	9.7	1.28	13	2.2	0.61
30-34	100	7.6	0.76	87	13.3	1.42	13	2.0	0.55
35-39	103	8.4	0.83	79	13.0	1.46	24	3.9	0.80
40-44	73	6.5	0.77	58	10.6	1.39	15	2.6	0.68
45-49	40	4.3	0.67	30	6.5	1.18	10	2.1	0.66
50-54	9	1.2	0.41	4	1.1	0.56	5	1.3	0.60
55-59	15	2.3	0.59	5	1.6	0.70	10	3.0	0.93
60-64	13	1.9	0.53	7	2.2	0.82	6	1.7	0.69
65-69	9	1.4	0.46	4	1.4	0.69	5	1.4	0.63
70-74	15	2.7	0.69	8	3.3	1.18	7	2.2	0.82
75-79	11	2.5	0.76	2	1.2	0.83	9	3.4	1.14
80-84	3	1.1	0.63	0	0.0	0.00	3	1.7	0.95
85+	10	4.3	1.36	7	11.7	4.41	3	1.7	1.00
Total deaths	515			380			135		
Connecticut									
. Crude rate		3.5	-		5.3	-		1.8	-
. AAMR (1940 Std. mil.)		3.2	0.14		4.8	0.25		1.6	0.14
. AAMR (1970 Std. mil.)		2.8	0.13		4.1	0.22		1.5	0.13
. AAMR (2000 Std. mil.)		3.3	0.15		5.0	0.26		1.7	0.15
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.1	-		5.5	-		2.7	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different