

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

## Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Infectious diseases and parasitic disease (1-139)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	9	4.5	1.50	6	5.8	2.38	3	3.1	1.78
5-9	1	0.6	0.56	0	0.0	0.00	1	1.1	1.15
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	4	2.2	1.11	3	3.2	1.87	1	1.1	1.14
20-24	6	2.8	1.14	6	5.5	2.26	0	0.0	0.00
25-29	24	9.6	1.96	18	14.3	3.38	6	4.8	1.97
30-34	43	16.2	2.47	38	28.6	4.64	5	3.8	1.68
35-39	44	18.3	2.76	36	30.1	5.02	8	6.6	2.35
40-44	24	10.7	2.19	20	18.2	4.07	4	3.5	1.76
45-49	27	14.8	2.85	21	23.4	5.12	6	6.5	2.64
50-54	25	17.6	3.52	20	28.8	6.44	5	6.9	3.08
55-59	17	12.6	3.07	14	21.5	5.75	3	4.3	2.50
60-64	22	15.8	3.38	13	19.7	5.47	9	12.3	4.11
65-69	54	40.9	5.57	27	45.5	8.76	27	37.1	7.15
70-74	46	41.4	6.10	27	57.4	11.05	19	29.6	6.80
75-79	77	91.6	10.44	38	116.2	18.85	39	76.0	12.16
80-84	80	150.4	16.81	34	190.6	32.70	46	130.1	19.18
85+	145	324.9	26.98	36	313.0	52.16	109	329.0	31.51
Total deaths	648			357			291		
Connecticut									
. Crude rate		22.0	-		25.0	-		19.2	-
. AAMR (1940 Std. mil.)		12.8	0.58		18.6	1.03		7.5	0.56
. AAMR (1970 Std. mil.)		15.5	0.63		21.2	1.14		10.5	0.67
. AAMR (2000 Std. mil.)		21.1	0.84		28.0	1.55		15.1	0.89
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		17.3	-		26.9	-		8.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Septicemia (038)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.87	2	1.9	1.38	1	1.0	1.03
5-9	1	0.6	0.56	0	0.0	0.00	1	1.1	1.15
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	0.9	0.66	2	1.8	1.30	0	0.0	0.00
25-29	3	1.2	0.69	2	1.6	1.13	1	0.8	0.80
30-34	2	0.8	0.53	2	1.5	1.06	0	0.0	0.00
35-39	4	1.7	0.83	2	1.7	1.18	2	1.7	1.17
40-44	3	1.3	0.77	2	1.8	1.29	1	0.9	0.88
45-49	8	4.4	1.55	3	3.3	1.93	5	5.4	2.41
50-54	7	4.9	1.86	3	4.3	2.49	4	5.5	2.76
55-59	8	6.0	2.10	6	9.2	3.76	2	2.9	2.04
60-64	15	10.8	2.79	7	10.6	4.02	8	11.0	3.87
65-69	37	28.0	4.61	17	28.7	6.95	20	27.5	6.15
70-74	39	35.1	5.62	23	48.9	10.20	16	24.9	6.24
75-79	70	83.3	9.95	35	107.0	18.09	35	68.2	11.52
80-84	70	131.6	15.73	29	162.6	30.20	41	115.9	18.10
85+	126	282.3	25.15	31	269.5	48.40	95	286.7	29.42
Total deaths	398			166			232		
Connecticut									
. Crude rate		13.5	-		11.6	-		15.3	-
. AAMR (1940 Std. mil.)		5.8	0.34		6.8	0.56		5.1	0.43
. AAMR (1970 Std. mil.)		8.9	0.46		10.2	0.79		8.0	0.57
. AAMR (2000 Std. mil.)		13.0	0.65		14.6	1.18		11.8	0.78
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.9	-		4.6	-		3.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is higher  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----HIV infection (042-044)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.71	2	1.9	1.38	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.55	0	0.0	0.00	1	1.1	1.14
20-24	4	1.9	0.93	4	3.7	1.84	0	0.0	0.00
25-29	19	7.6	1.74	14	11.1	2.98	5	4.0	1.79
30-34	38	14.3	2.32	34	25.6	4.39	4	3.0	1.50
35-39	35	14.6	2.46	30	25.1	4.58	5	4.1	1.85
40-44	16	7.2	1.79	15	13.7	3.53	1	0.9	0.88
45-49	15	8.2	2.13	14	15.6	4.18	1	1.1	1.08
50-54	15	10.6	2.73	14	20.2	5.39	1	1.4	1.38
55-59	6	4.5	1.82	6	9.2	3.76	0	0.0	0.00
60-64	4	2.9	1.44	3	4.6	2.63	1	1.4	1.37
65-69	4	3.0	1.52	3	5.1	2.92	1	1.4	1.38
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	159			139			20		
Connecticut									
. Crude rate		5.4	-		9.7	-		1.3	-
. AAMR (1940 Std. mil.)		5.1	0.41		9.0	0.76		1.3	0.29
. AAMR (1970 Std. mil.)		4.3	0.35		7.7	0.67		1.1	0.25
. AAMR (2000 Std. mil.)		5.2	0.42		9.3	0.80		1.2	0.28
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		9.8	-		17.8	-		2.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.71	1	1.0	0.97	1	1.0	1.03
5-9	10	5.6	1.77	5	5.5	2.45	5	5.7	2.57
10-14	2	1.2	0.85	1	1.2	1.18	1	1.2	1.23
15-19	5	2.8	1.24	2	2.2	1.53	3	3.4	1.98
20-24	10	4.7	1.47	5	4.6	2.06	5	4.7	2.10
25-29	21	8.4	1.83	11	8.8	2.64	10	8.0	2.54
30-34	45	16.9	2.52	22	16.6	3.53	23	17.3	3.61
35-39	61	25.4	3.25	22	18.4	3.92	39	32.3	5.18
40-44	99	44.3	4.45	30	27.3	4.99	69	60.6	7.29
45-49	172	94.4	7.20	82	91.5	10.11	90	97.1	10.24
50-54	248	174.6	11.09	119	171.3	15.70	129	177.7	15.65
55-59	394	293.1	14.76	193	296.6	21.35	201	289.8	20.44
60-64	670	482.2	18.63	359	544.9	28.76	311	425.7	24.14
65-69	952	721.3	23.38	533	899.1	38.94	419	576.4	28.16
70-74	1,043	938.3	29.06	581	1235.6	51.26	462	720.4	33.52
75-79	1,033	1229.0	38.24	559	1709.2	72.29	474	923.1	42.40
80-84	830	1560.1	54.15	418	2343.8	114.64	412	1164.9	57.39
85+	782	1752.1	62.65	304	2642.8	151.57	478	1442.8	65.99
Total deaths	6,379			3,247			3,132		
Connecticut									
. Crude rate		216.8	-		227.7	-		206.6	-
. AAMR (1940 Std. mil.)		121.1	1.68		144.9	2.67		105.9	2.20
. AAMR (1970 Std. mil.)		160.6	2.07		200.2	3.53		136.5	2.59
. AAMR (2000 Std. mil.)		201.9	2.55		258.3	4.71		169.9	3.09
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		134.8	-		166.1	-		112.6	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Oral cavity or pharynx (140.0-149.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.38	1	0.8	0.75	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.4	0.45	1	0.9	0.91	0	0.0	0.00
45-49	4	2.2	1.10	4	4.5	2.23	0	0.0	0.00
50-54	4	2.8	1.41	3	4.3	2.49	1	1.4	1.38
55-59	10	7.4	2.35	6	9.2	3.76	4	5.8	2.88
60-64	16	11.5	2.88	12	18.2	5.26	4	5.5	2.74
65-69	20	15.2	3.39	15	25.3	6.53	5	6.9	3.08
70-74	17	15.3	3.71	9	19.1	6.38	8	12.5	4.41
75-79	11	13.1	3.95	8	24.5	8.65	3	5.8	3.37
80-84	14	26.3	7.03	6	33.6	13.73	8	22.6	8.00
85+	9	20.2	6.72	5	43.5	19.44	4	12.1	6.04
Total deaths	107			70			37		
Connecticut									
. Crude rate		3.6	-		4.9	-		2.4	-
. AAMR (1940 Std. mil.)		2.2	0.23		3.4	0.42		1.2	0.23
. AAMR (1970 Std. mil.)		2.8	0.28		4.4	0.53		1.6	0.28
. AAMR (2000 Std. mil.)		3.3	0.33		5.3	0.66		2.0	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.4	-		3.8	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	4	1.5	0.75	0	0.0	0.00	4	3.0	1.50
35-39	3	1.2	0.72	3	2.5	1.45	0	0.0	0.00
40-44	7	3.1	1.18	2	1.8	1.29	5	4.4	1.96
45-49	12	6.6	1.90	9	10.0	3.35	3	3.2	1.87
50-54	23	16.2	3.38	10	14.4	4.55	13	17.9	4.97
55-59	33	24.5	4.27	16	24.6	6.15	17	24.5	5.94
60-64	66	47.5	5.85	43	65.3	9.95	23	31.5	6.56
65-69	109	82.6	7.91	61	102.9	13.17	48	66.0	9.53
70-74	117	105.3	9.73	61	129.7	16.61	56	87.3	11.67
75-79	139	165.4	14.03	69	211.0	25.40	70	136.3	16.29
80-84	128	240.6	21.27	57	319.6	42.33	71	200.7	23.82
85+	158	354.0	28.16	54	469.4	63.88	104	313.9	30.78
Total deaths	799			385			414		
Connecticut									
. Crude rate		27.2	-		27.0	-		27.3	-
. AAMR (1940 Std. mil.)		13.2	0.53		16.2	0.87		11.1	0.66
. AAMR (1970 Std. mil.)		19.0	0.69		23.7	1.21		15.9	0.83
. AAMR (2000 Std. mil.)		25.4	0.91		32.0	1.70		21.5	1.07
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.8	-		17.1	-		11.4	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Colon cancer (153)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	4	1.5	0.75	0	0.0	0.00	4	3.0	1.50
35-39	2	0.8	0.59	2	1.7	1.18	0	0.0	0.00
40-44	6	2.7	1.09	1	0.9	0.91	5	4.4	1.96
45-49	8	4.4	1.55	5	5.6	2.50	3	3.2	1.87
50-54	18	12.7	2.99	6	8.6	3.53	12	16.5	4.77
55-59	30	22.3	4.07	15	23.0	5.95	15	21.6	5.58
60-64	53	38.1	5.24	33	50.1	8.72	20	27.4	6.12
65-69	88	66.7	7.11	51	86.0	12.05	37	50.9	8.37
70-74	93	83.7	8.68	48	102.1	14.73	45	70.2	10.46
75-79	114	135.6	12.70	57	174.3	23.08	57	111.0	14.70
80-84	106	199.2	19.35	47	263.5	38.44	59	166.8	21.72
85+	127	284.5	25.25	44	382.5	57.67	83	250.5	27.50
Total deaths	649			309			340		
Connecticut									
. Crude rate		22.1	-		21.7	-		22.4	-
. AAMR (1940 Std. mil.)		10.7	0.47		12.9	0.77		9.4	0.61
. AAMR (1970 Std. mil.)		15.4	0.62		19.0	1.08		13.2	0.76
. AAMR (2000 Std. mil.)		20.6	0.82		25.7	1.53		17.8	0.98
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.6	-		14.2	-		10.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Rectal cancer (154)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.42	1	0.8	0.84	0	0.0	0.00
40-44	1	0.4	0.45	1	0.9	0.91	0	0.0	0.00
45-49	3	1.6	0.95	3	3.3	1.93	0	0.0	0.00
50-54	5	3.5	1.57	4	5.8	2.88	1	1.4	1.38
55-59	3	2.2	1.29	1	1.5	1.54	2	2.9	2.04
60-64	10	7.2	2.28	7	10.6	4.02	3	4.1	2.37
65-69	21	15.9	3.47	10	16.9	5.33	11	15.1	4.56
70-74	21	18.9	4.12	11	23.4	7.05	10	15.6	4.93
75-79	22	26.2	5.58	10	30.6	9.67	12	23.4	6.75
80-84	19	35.7	8.19	7	39.3	14.84	12	33.9	9.79
85+	22	49.3	10.51	4	34.8	17.39	18	54.3	12.81
Total deaths	128			59			69		
Connecticut									
. Crude rate		4.4	-		4.1	-		4.6	-
. AAMR (1940 Std. mil.)		2.2	0.22		2.7	0.37		1.7	0.24
. AAMR (1970 Std. mil.)		3.1	0.28		3.6	0.47		2.6	0.32
. AAMR (2000 Std. mil.)		4.1	0.36		4.6	0.62		3.5	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.0	-		2.5	-		1.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.38	1	0.8	0.75	0	0.0	0.00
35-39	2	0.8	0.59	1	0.8	0.84	1	0.8	0.83
40-44	1	0.4	0.45	0	0.0	0.00	1	0.9	0.88
45-49	7	3.8	1.45	4	4.5	2.23	3	3.2	1.87
50-54	14	9.9	2.63	9	13.0	4.32	5	6.9	3.08
55-59	19	14.1	3.24	12	18.4	5.32	7	10.1	3.81
60-64	32	23.0	4.07	19	28.8	6.62	13	17.8	4.94
65-69	58	43.9	5.77	43	72.5	11.06	15	20.6	5.33
70-74	53	47.7	6.55	31	65.9	11.84	22	34.3	7.31
75-79	49	58.3	8.33	25	76.4	15.29	24	46.7	9.54
80-84	48	90.2	13.02	20	112.1	25.08	28	79.2	14.96
85+	52	116.5	16.16	14	121.7	32.53	38	114.7	18.61
Total deaths	336			179			157		
Connecticut									
. Crude rate		11.4	-		12.6	-		10.4	-
. AAMR (1940 Std. mil.)		6.1	0.37		8.2	0.64		4.4	0.42
. AAMR (1970 Std. mil.)		8.3	0.47		11.1	0.83		6.2	0.53
. AAMR (2000 Std. mil.)		10.6	0.58		13.8	1.07		8.2	0.67
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.5	-		7.5	-		5.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Lung and other respiratory cancer (160-165)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.55	1	1.1	1.08	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.8	0.53	1	0.8	0.75	1	0.8	0.75
35-39	4	1.7	0.83	1	0.8	0.84	3	2.5	1.44
40-44	12	5.4	1.55	6	5.5	2.23	6	5.3	2.15
45-49	52	28.5	3.96	32	35.7	6.32	20	21.6	4.83
50-54	85	59.8	6.49	45	64.8	9.66	40	55.1	8.71
55-59	142	105.6	8.86	78	119.9	13.57	64	92.3	11.53
60-64	214	154.0	10.53	131	198.8	17.37	83	113.6	12.47
65-69	296	224.3	13.04	173	291.8	22.19	123	169.2	15.26
70-74	319	287.0	16.07	202	429.6	30.22	117	182.4	16.87
75-79	289	343.8	20.22	183	559.5	41.36	106	206.4	20.05
80-84	175	328.9	24.87	109	611.2	58.54	66	186.6	22.97
85+	102	228.5	22.63	57	495.5	65.63	45	135.8	20.25
Total deaths	1,693			1,019			674		
Connecticut									
. Crude rate		57.5	-		71.5	-		44.5	-
. AAMR (1940 Std. mil.)		34.4	0.91		47.0	1.54		25.4	1.09
. AAMR (1970 Std. mil.)		44.5	1.11		63.2	1.99		31.8	1.28
. AAMR (2000 Std. mil.)		53.1	1.30		78.2	2.53		37.2	1.46
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		41.3	-		61.1	-		26.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.8	0.53	1	0.8	0.75	1	0.8	0.75
35-39	4	1.7	0.83	1	0.8	0.84	3	2.5	1.44
40-44	12	5.4	1.55	6	5.5	2.23	6	5.3	2.15
45-49	51	28.0	3.92	32	35.7	6.32	19	20.5	4.70
50-54	83	58.4	6.41	45	64.8	9.66	38	52.3	8.49
55-59	140	104.1	8.80	76	116.8	13.40	64	92.3	11.53
60-64	213	153.3	10.50	130	197.3	17.30	83	113.6	12.47
65-69	287	217.5	12.84	165	278.3	21.67	122	167.8	15.19
70-74	309	278.0	15.81	197	418.9	29.85	112	174.6	16.50
75-79	277	329.5	19.80	174	532.0	40.33	103	200.6	19.76
80-84	166	312.0	24.22	103	577.5	56.91	63	178.1	22.44
85+	95	212.8	21.84	53	460.7	63.29	42	126.8	19.56
Total deaths	1,639			983			656		
Connecticut									
. Crude rate		55.7	-		68.9	-		43.3	-
. AAMR (1940 Std. mil.)		33.5	0.90		45.5	1.51		24.9	1.08
. AAMR (1970 Std. mil.)		43.2	1.09		60.9	1.95		31.1	1.27
. AAMR (2000 Std. mil.)		51.4	1.28		75.2	2.47		36.2	1.44
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		39.9	-		58.3	-		25.6	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	5	1.9	0.84	2	1.5	1.06	3	2.3	1.30
35-39	3	1.2	0.72	1	0.8	0.84	2	1.7	1.17
40-44	5	2.2	1.00	4	3.6	1.82	1	0.9	0.88
45-49	6	3.3	1.34	3	3.3	1.93	3	3.2	1.87
50-54	7	4.9	1.86	3	4.3	2.49	4	5.5	2.76
55-59	9	6.7	2.23	5	7.7	3.44	4	5.8	2.88
60-64	7	5.0	1.90	3	4.6	2.63	4	5.5	2.74
65-69	14	10.6	2.83	9	15.2	5.06	5	6.9	3.08
70-74	11	9.9	2.98	8	17.0	6.01	3	4.7	2.70
75-79	10	11.9	3.76	4	12.2	6.12	6	11.7	4.77
80-84	5	9.4	4.20	4	22.4	11.21	1	2.8	2.83
85+	5	11.2	5.01	1	8.7	8.69	4	12.1	6.04
Total deaths	87			47			40		
Connecticut									
. Crude rate		3.0	-		3.3	-		2.6	-
. AAMR (1940 Std. mil.)		2.1	0.24		2.4	0.37		1.8	0.31
. AAMR (1970 Std. mil.)		2.4	0.26		2.9	0.42		2.0	0.33
. AAMR (2000 Std. mil.)		2.8	0.31		3.5	0.52		2.4	0.38
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.7	-		1.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	2	0.8	0.57	-	-	-	2	1.6	1.13
30-34	4	1.5	0.75	-	-	-	4	3.0	1.50
35-39	14	5.8	1.56	-	-	-	14	11.6	3.10
40-44	30	13.4	2.45	-	-	-	30	26.3	4.81
45-49	32	17.6	3.10	-	-	-	32	34.5	6.11
50-54	21	14.8	3.23	-	-	-	21	28.9	6.31
55-59	43	32.0	4.88	-	-	-	43	62.0	9.45
60-64	70	50.4	6.02	-	-	-	70	95.8	11.45
65-69	71	53.8	6.38	-	-	-	71	97.7	11.59
70-74	76	68.4	7.84	-	-	-	76	118.5	13.59
75-79	78	92.8	10.51	-	-	-	78	151.9	17.20
80-84	52	97.7	13.55	-	-	-	52	147.0	20.39
85+	77	172.5	19.66	-	-	-	77	232.4	26.49
Total deaths	570			-			570		
Connecticut									
. Crude rate		-	-		-	-		37.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		21.3	1.02
. AAMR (1970 Std. mil.)		-	-		-	-		26.0	1.15
. AAMR (2000 Std. mil.)		-	-		-	-		31.9	1.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		23.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: CT rate is not significantly different



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	2	0.8	0.53	-	-	-	2	1.5	1.06
35-39	3	1.2	0.72	-	-	-	3	2.5	1.44
40-44	4	1.8	0.89	-	-	-	4	3.5	1.76
45-49	5	2.7	1.23	-	-	-	5	5.4	2.41
50-54	1	0.7	0.70	-	-	-	1	1.4	1.38
55-59	1	0.7	0.74	-	-	-	1	1.4	1.44
60-64	2	1.4	1.02	-	-	-	2	2.7	1.94
65-69	4	3.0	1.52	-	-	-	4	5.5	2.75
70-74	2	1.8	1.27	-	-	-	2	3.1	2.21
75-79	5	5.9	2.66	-	-	-	5	9.7	4.35
80-84	3	5.6	3.26	-	-	-	3	8.5	4.90
85+	4	9.0	4.48	-	-	-	4	12.1	6.04
Total deaths	36			-			36		
Connecticut									
. Crude rate		-	-		-	-		2.4	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.5	0.28
. AAMR (1970 Std. mil.)		-	-		-	-		1.7	0.29
. AAMR (2000 Std. mil.)		-	-		-	-		2.1	0.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	1	0.4	0.45	-	-	-	1	0.9	0.88
45-49	1	0.5	0.55	-	-	-	1	1.1	1.08
50-54	0	0.0	0.00	-	-	-	0	0.0	0.00
55-59	2	1.5	1.05	-	-	-	2	2.9	2.04
60-64	7	5.0	1.90	-	-	-	7	9.6	3.62
65-69	6	4.5	1.86	-	-	-	6	8.3	3.37
70-74	6	5.4	2.20	-	-	-	6	9.4	3.82
75-79	6	7.1	2.91	-	-	-	6	11.7	4.77
80-84	6	11.3	4.60	-	-	-	6	17.0	6.93
85+	5	11.2	5.01	-	-	-	5	15.1	6.75
Total deaths	40			-			40		
Connecticut									
. Crude rate		-	-		-	-		2.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.3	0.23
. AAMR (1970 Std. mil.)		-	-		-	-		1.7	0.28
. AAMR (2000 Std. mil.)		-	-		-	-		2.1	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	1	0.4	0.40	-	-	-	1	0.8	0.80
30-34	2	0.8	0.53	-	-	-	2	1.5	1.06
35-39	1	0.4	0.42	-	-	-	1	0.8	0.83
40-44	5	2.2	1.00	-	-	-	5	4.4	1.96
45-49	3	1.6	0.95	-	-	-	3	3.2	1.87
50-54	10	7.0	2.23	-	-	-	10	13.8	4.36
55-59	8	6.0	2.10	-	-	-	8	11.5	4.08
60-64	21	15.1	3.30	-	-	-	21	28.7	6.27
65-69	32	24.2	4.29	-	-	-	32	44.0	7.78
70-74	27	24.3	4.67	-	-	-	27	42.1	8.10
75-79	26	30.9	6.07	-	-	-	26	50.6	9.93
80-84	10	18.8	5.94	-	-	-	10	28.3	8.94
85+	15	33.6	8.68	-	-	-	15	45.3	11.69
Total deaths	161			-			161		
Connecticut									
. Crude rate		-	-		-	-		10.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		6.0	0.53
. AAMR (1970 Std. mil.)		-	-		-	-		7.4	0.61
. AAMR (2000 Std. mil.)		-	-		-	-		8.8	0.71
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Bladder cancer (188)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	1	0.5	0.55	0	0.0	0.00	1	1.1	1.08
50-54	1	0.7	0.70	0	0.0	0.00	1	1.4	1.38
55-59	4	3.0	1.49	3	4.6	2.66	1	1.4	1.44
60-64	13	9.4	2.59	9	13.7	4.55	4	5.5	2.74
65-69	14	10.6	2.83	11	18.6	5.59	3	4.1	2.38
70-74	29	26.1	4.84	20	42.5	9.51	9	14.0	4.68
75-79	27	32.1	6.18	20	61.2	13.67	7	13.6	5.15
80-84	27	50.7	9.77	16	89.7	22.43	11	31.1	9.38
85+	41	91.9	14.35	23	199.9	41.69	18	54.3	12.81
Total deaths	157			102			55		
Connecticut									
. Crude rate		5.3	-		7.2	-		3.6	-
. AAMR (1940 Std. mil.)		2.3	0.20		3.8	0.40		1.3	0.21
. AAMR (1970 Std. mil.)		3.5	0.29		6.2	0.62		2.0	0.28
. AAMR (2000 Std. mil.)		5.0	0.40		9.1	0.94		2.8	0.38
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.8	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Brain & other CNS cancer (191-192)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	2	1.1	0.79	2	2.2	1.55	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.55	0	0.0	0.00	1	1.1	1.14
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	4	1.6	0.80	4	3.2	1.59	0	0.0	0.00
30-34	6	2.3	0.92	2	1.5	1.06	4	3.0	1.50
35-39	8	3.3	1.18	5	4.2	1.87	3	2.5	1.44
40-44	6	2.7	1.09	2	1.8	1.29	4	3.5	1.76
45-49	9	4.9	1.65	8	8.9	3.16	1	1.1	1.08
50-54	10	7.0	2.23	6	8.6	3.53	4	5.5	2.76
55-59	9	6.7	2.23	5	7.7	3.44	4	5.8	2.88
60-64	15	10.8	2.79	8	12.1	4.29	7	9.6	3.62
65-69	18	13.6	3.21	11	18.6	5.59	7	9.6	3.64
70-74	18	16.2	3.82	8	17.0	6.01	10	15.6	4.93
75-79	24	28.6	5.83	11	33.6	10.14	13	25.3	7.02
80-84	10	18.8	5.94	4	22.4	11.21	6	17.0	6.93
85+	3	6.7	3.88	3	26.1	15.06	0	0.0	0.00
Total deaths	143			79			64		
Connecticut									
. Crude rate		4.9	-		5.5	-		4.2	-
. AAMR (1940 Std. mil.)		3.4	0.31		4.3	0.50		2.7	0.37
. AAMR (1970 Std. mil.)		3.9	0.34		4.9	0.56		3.1	0.40
. AAMR (2000 Std. mil.)		4.6	0.39		5.9	0.68		3.7	0.47
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.7	-		4.4	-		3.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.71	1	1.0	0.97	1	1.0	1.03
5-9	4	2.2	1.12	2	2.2	1.55	2	2.3	1.62
10-14	1	0.6	0.60	1	1.2	1.18	0	0.0	0.00
15-19	3	1.7	0.96	1	1.1	1.08	2	2.3	1.61
20-24	3	1.4	0.81	2	1.8	1.30	1	0.9	0.94
25-29	4	1.6	0.80	1	0.8	0.80	3	2.4	1.39
30-34	4	1.5	0.75	4	3.0	1.51	0	0.0	0.00
35-39	9	3.7	1.25	4	3.3	1.67	5	4.1	1.85
40-44	4	1.8	0.89	3	2.7	1.58	1	0.9	0.88
45-49	7	3.8	1.45	3	3.3	1.93	4	4.3	2.16
50-54	9	6.3	2.11	5	7.2	3.22	4	5.5	2.76
55-59	10	7.4	2.35	5	7.7	3.44	5	7.2	3.22
60-64	18	13.0	3.05	12	18.2	5.26	6	8.2	3.35
65-69	26	19.7	3.86	17	28.7	6.95	9	12.4	4.13
70-74	28	25.2	4.76	17	36.2	8.77	11	17.2	5.17
75-79	32	38.1	6.73	17	52.0	12.61	15	29.2	7.54
80-84	27	50.7	9.77	12	67.3	19.42	15	42.4	10.95
85+	36	80.7	13.44	13	113.0	31.34	23	69.4	14.48
Total deaths	227			120			107		
Connecticut									
. Crude rate		7.7	-		8.4	-		7.1	-
. AAMR (1940 Std. mil.)		4.6	0.35		5.8	0.56		3.8	0.45
. AAMR (1970 Std. mil.)		5.9	0.41		7.5	0.70		4.7	0.50
. AAMR (2000 Std. mil.)		7.3	0.49		9.5	0.91		5.9	0.59
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.0	-		6.3	-		3.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Diabetes mellitus (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.8	0.57	1	0.8	0.80	1	0.8	0.80
30-34	1	0.4	0.38	0	0.0	0.00	1	0.8	0.75
35-39	3	1.2	0.72	2	1.7	1.18	1	0.8	0.83
40-44	7	3.1	1.18	5	4.6	2.04	2	1.8	1.24
45-49	7	3.8	1.45	3	3.3	1.93	4	4.3	2.16
50-54	10	7.0	2.23	7	10.1	3.81	3	4.1	2.39
55-59	23	17.1	3.57	12	18.4	5.32	11	15.9	4.78
60-64	34	24.5	4.20	17	25.8	6.26	17	23.3	5.64
65-69	57	43.2	5.72	31	52.3	9.39	26	35.8	7.01
70-74	63	56.7	7.14	29	61.7	11.45	34	53.0	9.09
75-79	87	103.5	11.10	35	107.0	18.09	52	101.3	14.04
80-84	68	127.8	15.50	18	100.9	23.79	50	141.4	19.99
85+	85	190.4	20.66	20	173.9	38.88	65	196.2	24.34
Total deaths	447			180			267		
Connecticut									
. Crude rate		15.2	-		12.6	-		17.6	-
. AAMR (1940 Std. mil.)		7.5	0.40		8.0	0.63		6.9	0.51
. AAMR (1970 Std. mil.)		10.6	0.52		11.1	0.83		10.1	0.65
. AAMR (2000 Std. mil.)		14.2	0.68		14.5	1.12		13.8	0.85
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.7	-		12.5	-		11.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.8	0.57	1	0.8	0.80	1	0.8	0.80
30-34	3	1.1	0.65	2	1.5	1.06	1	0.8	0.75
35-39	8	3.3	1.18	5	4.2	1.87	3	2.5	1.44
40-44	19	8.5	1.95	13	11.8	3.28	6	5.3	2.15
45-49	33	18.1	3.15	14	15.6	4.18	19	20.5	4.70
50-54	35	24.6	4.16	20	28.8	6.44	15	20.7	5.34
55-59	73	54.3	6.36	35	53.8	9.09	38	54.8	8.89
60-64	158	113.7	9.05	89	135.1	14.32	69	94.4	11.37
65-69	256	194.0	12.12	136	229.4	19.67	120	165.1	15.07
70-74	351	315.8	16.86	172	365.8	27.89	179	279.1	20.86
75-79	403	479.4	23.88	192	587.0	42.37	211	410.9	28.29
80-84	348	654.1	35.06	126	706.5	62.94	222	627.7	42.13
85+	441	988.1	47.05	125	1086.7	97.19	316	953.8	53.66
Total deaths	2,130			930			1,200		
Connecticut									
. Crude rate		72.4	-		65.2	-		79.2	-
. AAMR (1940 Std. mil.)		33.7	0.82		38.7	1.33		30.1	1.04
. AAMR (1970 Std. mil.)		49.6	1.10		56.7	1.87		44.6	1.36
. AAMR (2000 Std. mil.)		67.6	1.47		77.0	2.63		61.4	1.79
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): No comparison is available  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.38	0	0.0	0.00	1	0.8	0.75
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	1	0.5	0.55	0	0.0	0.00	1	1.1	1.08
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
65-69	2	1.5	1.07	1	1.7	1.69	1	1.4	1.38
70-74	2	1.8	1.27	1	2.1	2.13	1	1.6	1.56
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	1	2.2	2.24	1	8.7	8.69	0	0.0	0.00
Total deaths	7			3			4		
Connecticut									
. Crude rate		0.2	-		0.2	-		0.3	-
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. AAMR (1970 Std. mil.)		-	-		-	-		-	-
. AAMR (2000 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.5	-		0.6	-		0.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									No comparison is available
. THE U.S. HP 2000 target:									No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Cardiovascular disease (390-459)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	15	7.5	1.94	10	9.7	3.08	5	5.2	2.30
5-9	3	1.7	0.97	1	1.1	1.09	2	2.3	1.62
10-14	1	0.6	0.60	1	1.2	1.18	0	0.0	0.00
15-19	1	0.6	0.55	0	0.0	0.00	1	1.1	1.14
20-24	6	2.8	1.14	3	2.8	1.60	3	2.8	1.63
25-29	19	7.6	1.74	9	7.2	2.39	10	8.0	2.54
30-34	25	9.4	1.88	18	13.6	3.19	7	5.3	1.99
35-39	34	14.2	2.43	25	20.9	4.18	9	7.5	2.49
40-44	80	35.8	4.00	56	51.0	6.82	24	21.1	4.30
45-49	134	73.5	6.35	101	112.8	11.22	33	35.6	6.20
50-54	202	142.2	10.00	137	197.2	16.85	65	89.5	11.11
55-59	342	254.4	13.76	232	356.5	23.40	110	158.6	15.12
60-64	614	441.9	17.83	438	664.8	31.76	176	240.9	18.16
65-69	936	709.2	23.18	582	981.7	40.69	354	486.9	25.88
70-74	1,383	1244.2	33.46	769	1635.4	58.97	614	957.4	38.64
75-79	1,707	2030.8	49.15	945	2889.4	93.99	762	1484.0	53.76
80-84	1,913	3595.7	82.21	847	4749.4	163.19	1,066	3014.0	92.31
85+	3,749	8399.6	137.18	1,007	8754.2	275.87	2,742	8276.5	158.06
Total deaths	11,164			5,181			5,983		
Connecticut									
. Crude rate		379.5	-		363.3	-		394.7	-
. AAMR (1940 Std. mil.)		157.7	1.73		210.7	3.10		117.1	1.91
. AAMR (1970 Std. mil.)		247.4	2.41		320.4	4.47		194.3	2.69
. AAMR (2000 Std. mil.)		362.9	3.45		456.0	6.62		298.5	3.89
. HP 2000 (1940 Std. mil.)		84.4	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		190.8	-		251.6	-		143.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is higher  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Diseases of the heart (390-398,402,404-429)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	14	7.0	1.87	10	9.7	3.08	4	4.1	2.06
5-9	3	1.7	0.97	1	1.1	1.09	2	2.3	1.62
10-14	1	0.6	0.60	1	1.2	1.18	0	0.0	0.00
15-19	1	0.6	0.55	0	0.0	0.00	1	1.1	1.14
20-24	4	1.9	0.93	2	1.8	1.30	2	1.9	1.33
25-29	13	5.2	1.44	8	6.4	2.25	5	4.0	1.79
30-34	16	6.0	1.50	12	9.0	2.61	4	3.0	1.50
35-39	28	11.7	2.20	22	18.4	3.92	6	5.0	2.03
40-44	64	28.6	3.58	53	48.3	6.63	11	9.7	2.91
45-49	116	63.7	5.91	92	102.7	10.71	24	25.9	5.29
50-54	163	114.7	8.99	117	168.4	15.57	46	63.4	9.34
55-59	292	217.2	12.71	208	319.6	22.16	84	121.1	13.21
60-64	532	382.9	16.60	391	593.4	30.01	141	193.0	16.25
65-69	799	605.4	21.42	496	836.7	37.57	303	416.8	23.94
70-74	1,141	1026.5	30.39	657	1397.2	54.51	484	754.7	34.31
75-79	1,364	1622.7	43.94	769	2351.3	84.79	595	1158.7	47.50
80-84	1,491	2802.5	72.58	688	3857.8	147.08	803	2270.4	80.12
85+	2,891	6477.3	120.47	799	6946.0	245.73	2,092	6314.5	138.06
Total deaths	8,933			4,326			4,607		
Connecticut									
. Crude rate		303.6	-		303.3	-		303.9	-
. AAMR (1940 Std. mil.)		128.7	1.58		178.2	2.87		90.5	1.68
. AAMR (1970 Std. mil.)		199.6	2.18		267.8	4.09		150.1	2.36
. AAMR (2000 Std. mil.)		289.6	3.08		377.0	5.99		229.5	3.40
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		152.1	-		206.7	-		108.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.55	0	0.0	0.00	1	1.1	1.14
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	4	1.6	0.80	2	1.6	1.13	2	1.6	1.13
30-34	8	3.0	1.06	8	6.0	2.13	0	0.0	0.00
35-39	17	7.1	1.72	16	13.4	3.35	1	0.8	0.83
40-44	51	22.8	3.19	42	38.3	5.90	9	7.9	2.63
45-49	86	47.2	5.09	68	75.9	9.21	18	19.4	4.58
50-54	121	85.2	7.74	90	129.6	13.66	31	42.7	7.67
55-59	221	164.4	11.06	164	252.0	19.68	57	82.2	10.88
60-64	412	296.5	14.61	311	472.0	26.77	101	138.2	13.76
65-69	598	453.1	18.53	384	647.7	33.05	214	294.4	20.12
70-74	875	787.2	26.61	514	1093.1	48.21	361	562.9	29.63
75-79	1,028	1223.0	38.14	594	1816.2	74.52	434	845.2	40.57
80-84	1,098	2063.8	62.28	506	2837.3	126.13	592	1673.8	68.79
85+	2,171	4864.1	104.39	604	5250.8	213.65	1,567	4729.9	119.48
Total deaths	6,691			3,303			3,388		
Connecticut									
. Crude rate		227.4	-		231.6	-		223.5	-
. AAMR (1940 Std. mil.)		96.0	1.36		136.1	2.50		65.0	1.40
. AAMR (1970 Std. mil.)		149.3	1.88		204.2	3.57		109.3	2.00
. AAMR (2000 Std. mil.)		216.8	2.67		286.9	5.22		168.3	2.91
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		121.6	-		169.0	-		84.1	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.38	0	0.0	0.00	1	0.8	0.75
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	1	0.5	0.55	1	1.1	1.12	0	0.0	0.00
50-54	1	0.7	0.70	1	1.4	1.44	0	0.0	0.00
55-59	2	1.5	1.05	2	3.1	2.17	0	0.0	0.00
60-64	3	2.2	1.25	2	3.0	2.15	1	1.4	1.37
65-69	20	15.2	3.39	11	18.6	5.59	9	12.4	4.13
70-74	28	25.2	4.76	15	31.9	8.24	13	20.3	5.62
75-79	49	58.3	8.33	26	79.5	15.59	23	44.8	9.34
80-84	75	141.0	16.28	31	173.8	31.22	44	124.4	18.75
85+	217	486.2	33.00	56	486.8	65.06	161	486.0	38.30
Total deaths	397			145			252		
Connecticut									
. Crude rate		13.5	-		10.2	-		16.6	-
. AAMR (1940 Std. mil.)		4.0	0.23		4.8	0.42		3.4	0.26
. AAMR (1970 Std. mil.)		7.8	0.40		9.0	0.75		6.9	0.45
. AAMR (2000 Std. mil.)		13.2	0.67		14.9	1.28		12.1	0.77
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.6	-		6.5	-		4.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	1	0.5	0.55	0	0.0	0.00	1	1.1	1.08
50-54	3	2.1	1.22	1	1.4	1.44	2	2.8	1.95
55-59	6	4.5	1.82	5	7.7	3.44	1	1.4	1.44
60-64	6	4.3	1.76	4	6.1	3.04	2	2.7	1.94
65-69	5	3.8	1.69	3	5.1	2.92	2	2.8	1.95
70-74	16	14.4	3.60	5	10.6	4.76	11	17.2	5.17
75-79	14	16.7	4.45	4	12.2	6.12	10	19.5	6.16
80-84	15	28.2	7.28	3	16.8	9.71	12	33.9	9.79
85+	39	87.4	13.99	11	95.6	28.83	28	84.5	15.97
Total deaths	105			36			69		
Connecticut									
. Crude rate		3.6	-		2.5	-		4.6	-
. AAMR (1940 Std. mil.)		1.5	0.17		1.5	0.27		1.5	0.22
. AAMR (1970 Std. mil.)		2.3	0.24		2.3	0.39		2.3	0.30
. AAMR (2000 Std. mil.)		3.5	0.34		3.3	0.58		3.5	0.43
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.8	-		2.1	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	0	0.0	0.00	1	1.0	1.03
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	0.9	0.66	1	0.9	0.92	1	0.9	0.94
25-29	4	1.6	0.80	1	0.8	0.80	3	2.4	1.39
30-34	6	2.3	0.92	5	3.8	1.68	1	0.8	0.75
35-39	5	2.1	0.93	3	2.5	1.45	2	1.7	1.17
40-44	14	6.3	1.67	1	0.9	0.91	13	11.4	3.16
45-49	12	6.6	1.90	7	7.8	2.95	5	5.4	2.41
50-54	25	17.6	3.52	10	14.4	4.55	15	20.7	5.34
55-59	30	22.3	4.07	10	15.4	4.86	20	28.8	6.45
60-64	57	41.0	5.43	31	47.0	8.45	26	35.6	6.98
65-69	86	65.2	7.03	51	86.0	12.05	35	48.1	8.14
70-74	154	138.5	11.16	65	138.2	17.15	89	138.8	14.71
75-79	245	291.5	18.62	121	370.0	33.63	124	241.5	21.69
80-84	323	607.1	33.78	116	650.4	60.39	207	585.3	40.68
85+	652	1460.8	57.21	159	1382.2	109.62	493	1488.1	67.02
Total deaths	1,616			581			1,035		
Connecticut									
. Crude rate		54.9	-		40.7	-		68.3	-
. AAMR (1940 Std. mil.)		20.4	0.59		21.4	0.94		19.8	0.79
. AAMR (1970 Std. mil.)		34.2	0.87		35.8	1.49		33.1	1.10
. AAMR (2000 Std. mil.)		53.3	1.33		55.1	2.38		52.0	1.63
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		27.7	-		30.1	-		25.5	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is higher  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is not significantly different





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Aortic aneurysm (441)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.40	0	0.0	0.00	1	0.8	0.80
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	3	2.1	1.22	3	4.3	2.49	0	0.0	0.00
55-59	5	3.7	1.66	3	4.6	2.66	2	2.9	2.04
60-64	8	5.8	2.04	6	9.1	3.72	2	2.7	1.94
65-69	25	18.9	3.79	21	35.4	7.73	4	5.5	2.75
70-74	30	27.0	4.93	20	42.5	9.51	10	15.6	4.93
75-79	30	35.7	6.52	19	58.1	13.33	11	21.4	6.46
80-84	29	54.5	10.12	18	100.9	23.79	11	31.1	9.38
85+	33	73.9	12.87	13	113.0	31.34	20	60.4	13.50
Total deaths	164			103			61		
Connecticut									
. Crude rate		5.6	-		7.2	-		4.0	-
. AAMR (1940 Std. mil.)		2.5	0.22		4.2	0.43		1.4	0.21
. AAMR (1970 Std. mil.)		3.8	0.30		6.3	0.62		2.1	0.29
. AAMR (2000 Std. mil.)		5.2	0.40		8.5	0.88		3.1	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.7	-		5.9	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.87	0	0.0	0.00	3	3.1	1.78
5-9	1	0.6	0.56	0	0.0	0.00	1	1.1	1.15
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	0.9	0.66	1	0.9	0.92	1	0.9	0.94
25-29	1	0.4	0.40	1	0.8	0.80	0	0.0	0.00
30-34	2	0.8	0.53	1	0.8	0.75	1	0.8	0.75
35-39	6	2.5	1.02	2	1.7	1.18	4	3.3	1.66
40-44	3	1.3	0.77	2	1.8	1.29	1	0.9	0.88
45-49	7	3.8	1.45	3	3.3	1.93	4	4.3	2.16
50-54	10	7.0	2.23	4	5.8	2.88	6	8.3	3.37
55-59	12	8.9	2.58	9	13.8	4.61	3	4.3	2.50
60-64	27	19.4	3.74	22	33.4	7.12	5	6.8	3.06
65-69	40	30.3	4.79	24	40.5	8.26	16	22.0	5.50
70-74	86	77.4	8.34	51	108.5	15.19	35	54.6	9.23
75-79	146	173.7	14.38	69	211.0	25.40	77	150.0	17.09
80-84	216	406.0	27.62	91	510.3	53.49	125	353.4	31.61
85+	530	1187.5	51.58	163	1417.0	110.99	367	1107.8	57.82
Total deaths	1,092			443			649		
Connecticut									
. Crude rate		37.1	-		31.1	-		42.8	-
. AAMR (1940 Std. mil.)		12.4	0.44		15.4	0.77		10.5	0.54
. AAMR (1970 Std. mil.)		22.2	0.69		27.5	1.31		19.2	0.81
. AAMR (2000 Std. mil.)		36.3	1.10		44.6	2.21		32.1	1.27
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		14.1	-		18.6	-		11.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	1	1.0	0.97	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.8	0.57	1	0.8	0.80	1	0.8	0.80
30-34	3	1.1	0.65	1	0.8	0.75	2	1.5	1.06
35-39	2	0.8	0.59	0	0.0	0.00	2	1.7	1.17
40-44	2	0.9	0.63	0	0.0	0.00	2	1.8	1.24
45-49	5	2.7	1.23	3	3.3	1.93	2	2.2	1.53
50-54	11	7.7	2.33	7	10.1	3.81	4	5.5	2.76
55-59	27	20.1	3.87	14	21.5	5.75	13	18.7	5.20
60-64	61	43.9	5.62	28	42.5	8.03	33	45.2	7.86
65-69	126	95.5	8.50	75	126.5	14.61	51	70.2	9.82
70-74	186	167.3	12.27	105	223.3	21.79	81	126.3	14.03
75-79	216	257.0	17.48	115	351.6	32.79	101	196.7	19.57
80-84	159	298.9	23.70	86	482.2	52.00	73	206.4	24.16
85+	171	383.1	29.30	78	678.1	76.78	93	280.7	29.11
Total deaths	972			514			458		
Connecticut									
. Crude rate		33.0	-		36.0	-		30.2	-
. AAMR (1940 Std. mil.)		15.1	0.53		20.2	0.93		12.0	0.65
. AAMR (1970 Std. mil.)		22.5	0.74		31.2	1.38		17.5	0.86
. AAMR (2000 Std. mil.)		30.5	0.98		43.8	2.01		23.4	1.10
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		19.7	-		27.2	-		14.6	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	1	1.0	0.97	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.40	1	0.8	0.80	0	0.0	0.00
30-34	2	0.8	0.53	1	0.8	0.75	1	0.8	0.75
35-39	1	0.4	0.42	0	0.0	0.00	1	0.8	0.83
40-44	1	0.4	0.45	0	0.0	0.00	1	0.9	0.88
45-49	1	0.5	0.55	1	1.1	1.12	0	0.0	0.00
50-54	3	2.1	1.22	3	4.3	2.49	0	0.0	0.00
55-59	5	3.7	1.66	2	3.1	2.17	3	4.3	2.50
60-64	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
65-69	3	2.3	1.31	1	1.7	1.69	2	2.8	1.95
70-74	3	2.7	1.56	1	2.1	2.13	2	3.1	2.21
75-79	6	7.1	2.91	2	6.1	4.32	4	7.8	3.89
80-84	2	3.8	2.66	0	0.0	0.00	2	5.7	4.00
85+	7	15.7	5.93	3	26.1	15.06	4	12.1	6.04
Total deaths	36			16			20		
Connecticut									
. Crude rate		1.2	-		1.1	-		1.3	-
. AAMR (1940 Std. mil.)		0.8	0.15		0.9	0.24		0.7	0.17
. AAMR (1970 Std. mil.)		0.9	0.16		1.1	0.27		0.8	0.20
. AAMR (2000 Std. mil.)		1.2	0.20		1.4	0.36		1.1	0.25
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.4	-		1.3	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	8	3.0	1.06	5	3.8	1.68	3	2.3	1.30
35-39	12	5.0	1.44	8	6.7	2.37	4	3.3	1.66
40-44	17	7.6	1.84	12	10.9	3.16	5	4.4	1.96
45-49	16	8.8	2.20	12	13.4	3.87	4	4.3	2.16
50-54	27	19.0	3.66	18	25.9	6.11	9	12.4	4.13
55-59	25	18.6	3.72	19	29.2	6.70	6	8.7	3.53
60-64	31	22.3	4.01	22	33.4	7.12	9	12.3	4.11
65-69	44	33.3	5.03	29	48.9	9.08	15	20.6	5.33
70-74	43	38.7	5.90	22	46.8	9.97	21	32.7	7.15
75-79	32	38.1	6.73	18	55.0	12.97	14	27.3	7.29
80-84	18	33.8	7.97	6	33.6	13.73	12	33.9	9.79
85+	9	20.2	6.72	6	52.2	21.29	3	9.1	5.23
Total deaths	282			177			105		
Connecticut									
. Crude rate		9.6	-		12.4	-		6.9	-
. AAMR (1940 Std. mil.)		6.7	0.43		9.6	0.75		4.3	0.46
. AAMR (1970 Std. mil.)		7.7	0.47		11.0	0.84		5.0	0.51
. AAMR (2000 Std. mil.)		9.1	0.55		13.1	1.01		6.0	0.60
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.6	-		12.1	-		5.3	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is not significantly different







## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	23	11.5	2.40	14	13.6	3.64	9	9.3	3.09
5-9	6	3.4	1.37	4	4.4	2.19	2	2.3	1.62
10-14	11	6.6	2.00	8	9.4	3.33	3	3.7	2.14
15-19	53	29.4	4.04	43	46.5	7.08	10	11.4	3.61
20-24	94	43.7	4.51	69	63.6	7.66	25	23.5	4.69
25-29	78	31.2	3.53	65	51.7	6.42	13	10.4	2.89
30-34	57	21.4	2.84	49	36.9	5.27	8	6.0	2.13
35-39	41	17.1	2.67	33	27.6	4.81	8	6.6	2.35
40-44	43	19.2	2.93	37	33.7	5.54	6	5.3	2.15
45-49	26	14.3	2.80	16	17.9	4.47	10	10.8	3.41
50-54	21	14.8	3.23	17	24.5	5.94	4	5.5	2.76
55-59	28	20.8	3.94	21	32.3	7.04	7	10.1	3.81
60-64	26	18.7	3.67	16	24.3	6.07	10	13.7	4.33
65-69	35	26.5	4.48	18	30.4	7.16	17	23.4	5.67
70-74	52	46.8	6.49	27	57.4	11.05	25	39.0	7.80
75-79	59	70.2	9.14	26	79.5	15.59	33	64.3	11.19
80-84	63	118.4	14.92	23	129.0	26.89	40	113.1	17.88
85+	118	264.4	24.34	40	347.7	54.98	78	235.4	26.66
Total deaths	834			526			308		
Connecticut									
. Crude rate		28.3	-		36.9	-		20.3	-
. AAMR (1940 Std. mil.)		22.4	0.87		33.1	1.52		12.0	0.86
. AAMR (1970 Std. mil.)		23.9	0.88		34.2	1.54		14.2	0.91
. AAMR (2000 Std. mil.)		27.2	0.95		38.5	1.75		17.1	0.99
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		32.6	-		47.7	-		17.9	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.87	2	1.9	1.38	1	1.0	1.03
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.60	1	1.2	1.18	0	0.0	0.00
15-19	2	1.1	0.78	2	2.2	1.53	0	0.0	0.00
20-24	1	0.5	0.47	1	0.9	0.92	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.38	1	0.8	0.75	0	0.0	0.00
35-39	1	0.4	0.42	1	0.8	0.84	0	0.0	0.00
40-44	3	1.3	0.77	3	2.7	1.58	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	1	0.7	0.70	1	1.4	1.44	0	0.0	0.00
55-59	2	1.5	1.05	1	1.5	1.54	1	1.4	1.44
60-64	8	5.8	2.04	3	4.6	2.63	5	6.8	3.06
65-69	8	6.1	2.14	5	8.4	3.77	3	4.1	2.38
70-74	13	11.7	3.24	7	14.9	5.63	6	9.4	3.82
75-79	22	26.2	5.58	7	21.4	8.09	15	29.2	7.54
80-84	28	52.6	9.95	11	61.7	18.60	17	48.1	11.66
85+	75	168.0	19.40	17	147.8	35.84	58	175.1	22.99
Total deaths	169			63			106		
Connecticut									
. Crude rate		5.7	-		4.4	-		7.0	-
. AAMR (1940 Std. mil.)		2.3	0.22		2.7	0.38		1.8	0.22
. AAMR (1970 Std. mil.)		3.7	0.30		4.0	0.52		3.2	0.34
. AAMR (2000 Std. mil.)		5.6	0.43		5.9	0.77		5.2	0.51
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.5	-		3.8	-		1.6	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.0	1.00	1	1.0	0.97	3	3.1	1.78
5-9	1	0.6	0.56	1	1.1	1.09	0	0.0	0.00
10-14	3	1.8	1.04	3	3.5	2.04	0	0.0	0.00
15-19	2	1.1	0.78	2	2.2	1.53	0	0.0	0.00
20-24	3	1.4	0.81	3	2.8	1.60	0	0.0	0.00
25-29	3	1.2	0.69	3	2.4	1.38	0	0.0	0.00
30-34	4	1.5	0.75	3	2.3	1.30	1	0.8	0.75
35-39	3	1.2	0.72	3	2.5	1.45	0	0.0	0.00
40-44	4	1.8	0.89	4	3.6	1.82	0	0.0	0.00
45-49	1	0.5	0.55	1	1.1	1.12	0	0.0	0.00
50-54	3	2.1	1.22	3	4.3	2.49	0	0.0	0.00
55-59	3	2.2	1.29	3	4.6	2.66	0	0.0	0.00
60-64	2	1.4	1.02	2	3.0	2.15	0	0.0	0.00
65-69	1	0.8	0.76	1	1.7	1.69	0	0.0	0.00
70-74	2	1.8	1.27	2	4.3	3.01	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	1	1.9	1.88	1	5.6	5.61	0	0.0	0.00
85+	2	4.5	3.17	1	8.7	8.69	1	3.0	3.02
Total deaths	42			37			5		
Connecticut									
. Crude rate		1.4	-		2.6	-		0.3	-
. AAMR (1940 Std. mil.)		1.4	0.23		2.5	0.43		-	-
. AAMR (1970 Std. mil.)		1.4	0.23		2.6	0.44		-	-
. AAMR (2000 Std. mil.)		1.4	0.23		2.7	0.45		-	-
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.8	-		3.1	-		0.6	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	2.5	1.12	2	1.9	1.38	3	3.1	1.78
5-9	2	1.1	0.79	2	2.2	1.55	0	0.0	0.00
10-14	1	0.6	0.60	1	1.2	1.18	0	0.0	0.00
15-19	1	0.6	0.55	1	1.1	1.08	0	0.0	0.00
20-24	2	0.9	0.66	2	1.8	1.30	0	0.0	0.00
25-29	1	0.4	0.40	0	0.0	0.00	1	0.8	0.80
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.42	0	0.0	0.00	1	0.8	0.83
40-44	2	0.9	0.63	2	1.8	1.29	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	1	0.7	0.70	0	0.0	0.00	1	1.4	1.38
55-59	2	1.5	1.05	2	3.1	2.17	0	0.0	0.00
60-64	3	2.2	1.25	1	1.5	1.52	2	2.7	1.94
65-69	1	0.8	0.76	0	0.0	0.00	1	1.4	1.38
70-74	5	4.5	2.01	5	10.6	4.76	0	0.0	0.00
75-79	1	1.2	1.19	0	0.0	0.00	1	1.9	1.95
80-84	3	5.6	3.26	0	0.0	0.00	3	8.5	4.90
85+	1	2.2	2.24	0	0.0	0.00	1	3.0	3.02
Total deaths	32			18			14		
Connecticut									
. Crude rate		1.1	-		1.3	-		0.9	-
. AAMR (1940 Std. mil.)		0.9	0.18		1.2	0.30		-	-
. AAMR (1970 Std. mil.)		1.1	0.20		1.4	0.33		-	-
. AAMR (2000 Std. mil.)		1.1	0.19		1.3	0.30		-	-
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.4	-		1.9	-		1.0	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is higher  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	1.2	0.85	2	2.4	1.67	0	0.0	0.00
15-19	16	8.9	2.22	13	14.0	3.90	3	3.4	1.98
20-24	24	11.2	2.28	22	20.3	4.33	2	1.9	1.33
25-29	32	12.8	2.26	25	19.9	3.98	7	5.6	2.12
30-34	28	10.5	1.99	22	16.6	3.53	6	4.5	1.84
35-39	27	11.2	2.16	21	17.6	3.83	6	5.0	2.03
40-44	19	8.5	1.95	18	16.4	3.87	1	0.9	0.88
45-49	19	10.4	2.39	14	15.6	4.18	5	5.4	2.41
50-54	17	12.0	2.90	15	21.6	5.58	2	2.8	1.95
55-59	15	11.2	2.88	9	13.8	4.61	6	8.7	3.53
60-64	15	10.8	2.79	12	18.2	5.26	3	4.1	2.37
65-69	17	12.9	3.12	13	21.9	6.08	4	5.5	2.75
70-74	13	11.7	3.24	10	21.3	6.72	3	4.7	2.70
75-79	9	10.7	3.57	7	21.4	8.09	2	3.9	2.75
80-84	10	18.8	5.94	8	44.9	15.86	2	5.7	4.00
85+	9	20.2	6.72	7	60.9	23.00	2	6.0	4.27
Total deaths	272			218			54		
Connecticut									
. Crude rate		9.2	-		15.3	-		3.6	-
. AAMR (1940 Std. mil.)		8.3	0.53		13.8	0.97		3.1	0.45
. AAMR (1970 Std. mil.)		8.0	0.51		13.6	0.95		3.0	0.43
. AAMR (2000 Std. mil.)		8.9	0.55		15.4	1.07		3.3	0.46
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.5	-		19.1	-		4.4	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All injury (E800-E999)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	28	14.0	2.65	19	18.5	4.24	9	9.3	3.09
5-9	6	3.4	1.37	4	4.4	2.19	2	2.3	1.62
10-14	13	7.8	2.17	10	11.8	3.73	3	3.7	2.14
15-19	78	43.3	4.90	65	70.2	8.71	13	14.8	4.11
20-24	137	63.7	5.45	109	100.5	9.63	28	26.3	4.97
25-29	124	49.5	4.45	102	81.2	8.04	22	17.6	3.76
30-34	103	38.8	3.82	81	61.0	6.78	22	16.5	3.53
35-39	80	33.3	3.73	61	51.0	6.53	19	15.8	3.62
40-44	74	33.1	3.85	62	56.5	7.17	12	10.5	3.04
45-49	51	28.0	3.92	35	39.1	6.60	16	17.3	4.32
50-54	39	27.5	4.40	33	47.5	8.27	6	8.3	3.37
55-59	44	32.7	4.93	31	47.6	8.56	13	18.7	5.20
60-64	43	30.9	4.72	29	44.0	8.17	14	19.2	5.12
65-69	57	43.2	5.72	35	59.0	9.98	22	30.3	6.45
70-74	72	64.8	7.63	41	87.2	13.62	31	48.3	8.68
75-79	69	82.1	9.88	34	104.0	17.83	35	68.2	11.52
80-84	75	141.0	16.28	32	179.4	31.72	43	121.6	18.54
85+	128	286.8	25.35	47	408.6	59.60	81	244.5	27.17
Total deaths	1,221			830			391		
Connecticut									
. Crude rate		41.5	-		58.2	-		25.8	-
. AAMR (1940 Std. mil.)		34.5	1.08		52.9	1.92		16.6	1.02
. AAMR (1970 Std. mil.)		35.3	1.07		53.3	1.92		18.5	1.03
. AAMR (2000 Std. mil.)		39.7	1.15		59.5	2.14		22.1	1.14
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		55.1	-		84.3	-		26.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	8	4.0	1.42	7	6.8	2.57	1	1.0	1.03
5-9	3	1.7	0.97	1	1.1	1.09	2	2.3	1.62
10-14	5	3.0	1.35	2	2.4	1.67	3	3.7	2.14
15-19	42	23.3	3.60	33	35.7	6.21	9	10.3	3.42
20-24	73	34.0	3.97	51	47.0	6.59	22	20.7	4.40
25-29	59	23.6	3.07	49	39.0	5.57	10	8.0	2.54
30-34	33	12.4	2.16	29	21.8	4.05	4	3.0	1.50
35-39	21	8.7	1.91	15	12.5	3.24	6	5.0	2.03
40-44	15	6.7	1.73	11	10.0	3.02	4	3.5	1.76
45-49	15	8.2	2.13	9	10.0	3.35	6	6.5	2.64
50-54	10	7.0	2.23	9	13.0	4.32	1	1.4	1.38
55-59	13	9.7	2.68	8	12.3	4.35	5	7.2	3.22
60-64	7	5.0	1.90	5	7.6	3.39	2	2.7	1.94
65-69	10	7.6	2.40	5	8.4	3.77	5	6.9	3.08
70-74	15	13.5	3.48	5	10.6	4.76	10	15.6	4.93
75-79	16	19.0	4.76	8	24.5	8.65	8	15.6	5.51
80-84	13	24.4	6.78	4	22.4	11.21	9	25.4	8.48
85+	6	13.4	5.49	3	26.1	15.06	3	9.1	5.23
Total deaths	364			254			110		
Connecticut									
. Crude rate		12.4	-		17.8	-		7.3	-
. AAMR (1940 Std. mil.)		12.3	0.68		18.0	1.16		6.6	0.70
. AAMR (1970 Std. mil.)		11.8	0.65		16.9	1.10		6.7	0.69
. AAMR (2000 Std. mil.)		11.8	0.63		16.8	1.08		6.8	0.66
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		18.2	-		25.8	-		10.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.87	2	1.9	1.38	1	1.0	1.03
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.60	1	1.2	1.18	0	0.0	0.00
15-19	1	0.6	0.55	1	1.1	1.08	0	0.0	0.00
20-24	2	0.9	0.66	2	1.8	1.30	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.8	0.53	2	1.5	1.06	0	0.0	0.00
35-39	1	0.4	0.42	1	0.8	0.84	0	0.0	0.00
40-44	3	1.3	0.77	3	2.7	1.58	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	1	0.7	0.70	1	1.4	1.44	0	0.0	0.00
55-59	3	2.2	1.29	2	3.1	2.17	1	1.4	1.44
60-64	9	6.5	2.16	4	6.1	3.04	5	6.8	3.06
65-69	7	5.3	2.00	4	6.7	3.37	3	4.1	2.38
70-74	12	10.8	3.12	7	14.9	5.63	5	7.8	3.49
75-79	13	15.5	4.29	3	9.2	5.30	10	19.5	6.16
80-84	22	41.4	8.82	9	50.5	16.82	13	36.8	10.19
85+	53	118.7	16.31	12	104.3	30.11	41	123.8	19.33
Total deaths	133			54			79		
Connecticut									
. Crude rate		4.5	-		3.8	-		5.2	-
. AAMR (1940 Std. mil.)		2.0	0.21		2.5	0.37		1.4	0.21
. AAMR (1970 Std. mil.)		3.0	0.27		3.5	0.48		2.5	0.30
. AAMR (2000 Std. mil.)		4.4	0.38		4.8	0.68		3.9	0.44
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.5	-		3.6	-		1.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

All firearms (E922.0-.3, E922.8-.9, E955.0-.4, E965.0-.4, E970, E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	1.2	0.85	2	2.4	1.67	0	0.0	0.00
15-19	16	8.9	2.22	14	15.1	4.04	2	2.3	1.61
20-24	30	14.0	2.55	29	26.7	4.97	1	0.9	0.94
25-29	22	8.8	1.87	19	15.1	3.47	3	2.4	1.39
30-34	24	9.0	1.84	16	12.0	3.01	8	6.0	2.13
35-39	22	9.2	1.95	17	14.2	3.45	5	4.1	1.85
40-44	14	6.3	1.67	11	10.0	3.02	3	2.6	1.52
45-49	11	6.0	1.82	10	11.2	3.53	1	1.1	1.08
50-54	7	4.9	1.86	7	10.1	3.81	0	0.0	0.00
55-59	5	3.7	1.66	4	6.1	3.07	1	1.4	1.44
60-64	7	5.0	1.90	7	10.6	4.02	0	0.0	0.00
65-69	8	6.1	2.14	6	10.1	4.13	2	2.8	1.95
70-74	7	6.3	2.38	6	12.8	5.21	1	1.6	1.56
75-79	2	2.4	1.68	2	6.1	4.32	0	0.0	0.00
80-84	8	15.0	5.32	7	39.3	14.84	1	2.8	2.83
85+	3	6.7	3.88	3	26.1	15.06	0	0.0	0.00
Total deaths	188			160			28		
Connecticut									
. Crude rate		6.4	-		11.2	-		1.8	-
. AAMR (1940 Std. mil.)		6.1	0.47		10.7	0.88		1.7	0.33
. AAMR (1970 Std. mil.)		5.7	0.44		10.3	0.84		1.5	0.29
. AAMR (2000 Std. mil.)		6.1	0.45		11.0	0.89		1.7	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		14.5	-		25.3	-		4.4	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	1.2	0.85	2	2.4	1.67	0	0.0	0.00
15-19	9	5.0	1.66	7	7.6	2.86	2	2.3	1.61
20-24	9	4.2	1.40	9	8.3	2.77	0	0.0	0.00
25-29	10	4.0	1.26	8	6.4	2.25	2	1.6	1.13
30-34	14	5.3	1.41	11	8.3	2.50	3	2.3	1.30
35-39	15	6.2	1.61	13	10.9	3.02	2	1.7	1.17
40-44	9	4.0	1.34	8	7.3	2.58	1	0.9	0.88
45-49	8	4.4	1.55	7	7.8	2.95	1	1.1	1.08
50-54	6	4.2	1.72	6	8.6	3.53	0	0.0	0.00
55-59	4	3.0	1.49	3	4.6	2.66	1	1.4	1.44
60-64	7	5.0	1.90	7	10.6	4.02	0	0.0	0.00
65-69	7	5.3	2.00	5	8.4	3.77	2	2.8	1.95
70-74	5	4.5	2.01	5	10.6	4.76	0	0.0	0.00
75-79	2	2.4	1.68	2	6.1	4.32	0	0.0	0.00
80-84	6	11.3	4.60	6	33.6	13.73	0	0.0	0.00
85+	3	6.7	3.88	3	26.1	15.06	0	0.0	0.00
Total deaths	116			102			14		
Connecticut									
. Crude rate		3.9	-		7.2	-		0.9	-
. AAMR (1940 Std. mil.)		3.6	0.35		6.4	0.67		-	-
. AAMR (1970 Std. mil.)		3.4	0.34		6.4	0.66		-	-
. AAMR (2000 Std. mil.)		3.8	0.36		7.3	0.74		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.9	-		12.5	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	5	2.8	1.24	5	5.4	2.42	0	0.0	0.00
20-24	16	7.4	1.86	15	13.8	3.57	1	0.9	0.94
25-29	10	4.0	1.26	9	7.2	2.39	1	0.8	0.80
30-34	7	2.6	1.00	3	2.3	1.30	4	3.0	1.50
35-39	6	2.5	1.02	4	3.3	1.67	2	1.7	1.17
40-44	3	1.3	0.77	1	0.9	0.91	2	1.8	1.24
45-49	2	1.1	0.78	2	2.2	1.58	0	0.0	0.00
50-54	1	0.7	0.70	1	1.4	1.44	0	0.0	0.00
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
65-69	1	0.8	0.76	1	1.7	1.69	0	0.0	0.00
70-74	2	1.8	1.27	1	2.1	2.13	1	1.6	1.56
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	2	3.8	2.66	1	5.6	5.61	1	2.8	2.83
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	55			43			12		
Connecticut									
. Crude rate		1.9	-		3.0	-		0.8	-
. AAMR (1940 Std. mil.)		1.9	0.27		3.1	0.49		-	-
. AAMR (1970 Std. mil.)		1.7	0.24		2.8	0.45		-	-
. AAMR (2000 Std. mil.)		1.7	0.24		2.8	0.43		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.7	-		11.4	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All poisoning (E850-E869, E950-E952, E962, E972, E980-E982)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	3	1.7	0.96	3	3.2	1.87	0	0.0	0.00
20-24	8	3.7	1.32	8	7.4	2.61	0	0.0	0.00
25-29	11	4.4	1.33	9	7.2	2.39	2	1.6	1.13
30-34	17	6.4	1.55	13	9.8	2.71	4	3.0	1.50
35-39	9	3.7	1.25	7	5.9	2.21	2	1.7	1.17
40-44	3	1.3	0.77	3	2.7	1.58	0	0.0	0.00
45-49	13	7.1	1.98	8	8.9	3.16	5	5.4	2.41
50-54	4	2.8	1.41	3	4.3	2.49	1	1.4	1.38
55-59	5	3.7	1.66	2	3.1	2.17	3	4.3	2.50
60-64	6	4.3	1.76	4	6.1	3.04	2	2.7	1.94
65-69	7	5.3	2.00	4	6.7	3.37	3	4.1	2.38
70-74	7	6.3	2.38	5	10.6	4.76	2	3.1	2.21
75-79	2	2.4	1.68	1	3.1	3.06	1	1.9	1.95
80-84	2	3.8	2.66	0	0.0	0.00	2	5.7	4.00
85+	5	11.2	5.01	4	34.8	17.39	1	3.0	3.02
Total deaths	102			74			28		
Connecticut									
. Crude rate		3.5	-		5.2	-		1.8	-
. AAMR (1940 Std. mil.)		3.0	0.31		4.6	0.55		1.4	0.29
. AAMR (1970 Std. mil.)		2.9	0.29		4.4	0.53		1.4	0.28
. AAMR (2000 Std. mil.)		3.2	0.33		5.1	0.61		1.7	0.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.5	-		6.3	-		3.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.5	0.47	1	0.9	0.92	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	6	2.3	0.92	3	2.3	1.30	3	2.3	1.30
35-39	11	4.6	1.38	8	6.7	2.37	3	2.5	1.44
40-44	12	5.4	1.55	8	7.3	2.58	4	3.5	1.76
45-49	13	7.1	1.98	9	10.0	3.35	4	4.3	2.16
50-54	21	14.8	3.23	14	20.2	5.39	7	9.6	3.64
55-59	17	12.6	3.07	14	21.5	5.75	3	4.3	2.50
60-64	14	10.1	2.69	8	12.1	4.29	6	8.2	3.35
65-69	19	14.4	3.30	13	21.9	6.08	6	8.3	3.37
70-74	19	17.1	3.92	12	25.5	7.37	7	10.9	4.13
75-79	3	3.6	2.06	3	9.2	5.30	0	0.0	0.00
80-84	5	9.4	4.20	3	16.8	9.71	2	5.7	4.00
85+	2	4.5	3.17	2	17.4	12.29	0	0.0	0.00
Total deaths	143			98			45		
Connecticut									
. Crude rate		4.9	-		6.9	-		3.0	-
. AAMR (1940 Std. mil.)		3.9	0.34		5.7	0.60		2.4	0.37
. AAMR (1970 Std. mil.)		4.2	0.36		6.2	0.63		2.5	0.38
. AAMR (2000 Std. mil.)		4.8	0.41		7.2	0.75		2.8	0.43
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.2	-		11.4	-		3.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1990

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	5	2.0	0.89	4	3.2	1.59	1	0.8	0.80
30-34	10	3.8	1.19	8	6.0	2.13	2	1.5	1.06
35-39	7	2.9	1.10	7	5.9	2.21	0	0.0	0.00
40-44	1	0.4	0.45	1	0.9	0.91	0	0.0	0.00
45-49	5	2.7	1.23	2	2.2	1.58	3	3.2	1.87
50-54	1	0.7	0.70	0	0.0	0.00	1	1.4	1.38
55-59	1	0.7	0.74	0	0.0	0.00	1	1.4	1.44
60-64	2	1.4	1.02	2	3.0	2.15	0	0.0	0.00
65-69	3	2.3	1.31	2	3.4	2.39	1	1.4	1.38
70-74	5	4.5	2.01	2	4.3	3.01	3	4.7	2.70
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	2	3.8	2.66	0	0.0	0.00	2	5.7	4.00
85+	4	9.0	4.48	3	26.1	15.06	1	3.0	3.02
Total deaths	46			31			15		
Connecticut									
. Crude rate		1.6	-		2.2	-		1.0	-
. AAMR (1940 Std. mil.)		1.2	0.19		1.7	0.32		0.7	0.20
. AAMR (1970 Std. mil.)		1.1	0.17		1.6	0.29		0.7	0.20
. AAMR (2000 Std. mil.)		1.4	0.21		2.1	0.40		0.9	0.24
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.4	-		4.5	-		2.6	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower