

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

## Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Infectious diseases and parasitic disease (1-139)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	13	3.2	0.90	6	2.9	1.19	7	3.6	1.35
5-9	2	0.5	0.39	0	0.0	0.00	2	1.1	0.79
10-14	1	0.3	0.29	0	0.0	0.00	1	0.6	0.60
15-19	3	0.9	0.51	0	0.0	0.00	3	1.8	1.06
20-24	15	3.7	0.95	8	3.9	1.36	7	3.5	1.32
25-29	46	9.8	1.45	32	13.6	2.40	14	6.0	1.61
30-34	124	23.5	2.11	100	38.0	3.80	24	9.1	1.86
35-39	128	25.9	2.29	109	44.4	4.25	19	7.7	1.76
40-44	93	20.6	2.14	77	34.7	3.96	16	7.0	1.74
45-49	78	20.4	2.31	66	35.1	4.32	12	6.2	1.79
50-54	38	13.1	2.12	31	21.8	3.91	7	4.7	1.78
55-59	43	16.7	2.54	33	26.4	4.59	10	7.5	2.38
60-64	55	20.6	2.77	36	28.3	4.71	19	13.6	3.11
65-69	63	24.6	3.10	32	27.8	4.91	31	22.0	3.95
70-74	107	47.1	4.56	50	51.7	7.31	57	43.8	5.79
75-79	137	77.8	6.65	67	96.2	11.75	70	65.7	7.86
80-84	125	111.9	10.01	51	133.6	18.71	74	100.7	11.71
85+	261	275.4	17.05	74	304.6	35.41	187	265.3	19.40
Total deaths	1,332			772			560		
Connecticut									
. Crude rate		22.7	-		27.2	-		18.6	-
. AAMR (1940 Std. mil.)		14.1	0.44		20.3	0.77		8.2	0.45
. AAMR (1970 Std. mil.)		15.8	0.45		21.8	0.80		10.5	0.49
. AAMR (2000 Std. mil.)		21.3	0.59		29.0	1.08		14.4	0.62
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		18.7	-		29.1	-		8.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----HIV infection (042-044)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	1.2	0.56	2	1.0	0.68	3	1.5	0.89
5-9	2	0.5	0.39	0	0.0	0.00	2	1.1	0.79
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	12	2.9	0.85	6	2.9	1.18	6	3.0	1.22
25-29	41	8.8	1.37	30	12.7	2.33	11	4.7	1.43
30-34	112	21.3	2.01	91	34.6	3.62	21	8.0	1.74
35-39	113	22.9	2.15	96	39.1	3.99	17	6.9	1.66
40-44	83	18.4	2.02	71	32.0	3.80	12	5.2	1.51
45-49	52	13.6	1.89	46	24.5	3.61	6	3.1	1.26
50-54	20	6.9	1.54	18	12.6	2.98	2	1.3	0.95
55-59	16	6.2	1.55	15	12.0	3.10	1	0.8	0.75
60-64	15	5.6	1.45	13	10.2	2.83	2	1.4	1.01
65-69	3	1.2	0.68	3	2.6	1.50	0	0.0	0.00
70-74	4	1.8	0.88	2	2.1	1.46	2	1.5	1.09
75-79	4	2.3	1.14	1	1.4	1.44	3	2.8	1.63
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	482			394			88		
Connecticut									
. Crude rate		8.2	-		13.9	-		2.9	-
. AAMR (1940 Std. mil.)		7.5	0.34		12.3	0.62		2.8	0.30
. AAMR (1970 Std. mil.)		6.4	0.30		10.5	0.54		2.4	0.26
. AAMR (2000 Std. mil.)		7.8	0.36		13.0	0.66		2.8	0.30
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.3	-		20.1	-		2.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	7	1.7	0.66	5	2.4	1.08	2	1.0	0.72
5-9	7	1.9	0.72	2	1.1	0.75	5	2.8	1.25
10-14	10	2.9	0.93	7	4.0	1.52	3	1.8	1.04
15-19	11	3.3	0.98	7	4.0	1.52	4	2.4	1.22
20-24	30	7.4	1.34	20	9.7	2.16	10	5.0	1.58
25-29	26	5.6	1.09	14	5.9	1.59	12	5.2	1.49
30-34	67	12.7	1.55	36	13.7	2.28	31	11.8	2.11
35-39	131	26.5	2.32	56	22.8	3.04	75	30.2	3.49
40-44	233	51.6	3.38	110	49.6	4.73	123	53.5	4.83
45-49	371	97.2	5.04	178	94.7	7.10	193	99.5	7.16
50-54	525	180.4	7.87	255	179.1	11.22	270	181.5	11.05
55-59	795	308.3	10.94	417	333.3	16.32	378	284.8	14.65
60-64	1,265	473.1	13.30	689	540.8	20.60	576	411.6	17.15
65-69	1,910	745.5	17.06	1,059	919.8	28.27	851	603.2	20.68
70-74	2,213	975.0	20.73	1,184	1224.4	35.58	1,029	789.8	24.62
75-79	2,172	1233.2	26.46	1,129	1621.2	48.25	1,043	979.5	30.33
80-84	1,731	1550.2	37.26	873	2286.9	77.40	858	1167.5	39.86
85+	1,669	1761.1	43.11	672	2766.5	106.72	997	1414.5	44.80
Total deaths	13,173			6,713			6,460		
Connecticut									
. Crude rate		224.8	-		236.1	-		214.2	-
. AAMR (1940 Std. mil.)		123.6	1.20		148.3	1.92		106.6	1.55
. AAMR (1970 Std. mil.)		163.5	1.47		203.3	2.50		138.3	1.84
. AAMR (2000 Std. mil.)		205.2	1.80		261.8	3.30		172.2	2.18
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		134.4	-		164.9	-		112.5	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Oral cavity or pharynx (140.0-149.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	3	0.6	0.35	3	1.2	0.70	0	0.0	0.00
40-44	3	0.7	0.38	3	1.4	0.78	0	0.0	0.00
45-49	11	2.9	0.87	8	4.3	1.51	3	1.5	0.89
50-54	18	6.2	1.46	15	10.5	2.72	3	2.0	1.16
55-59	22	8.5	1.82	16	12.8	3.20	6	4.5	1.85
60-64	28	10.5	1.98	22	17.3	3.68	6	4.3	1.75
65-69	39	15.2	2.44	22	19.1	4.07	17	12.1	2.92
70-74	25	11.0	2.20	11	11.4	3.43	14	10.7	2.87
75-79	36	20.4	3.41	17	24.4	5.92	19	17.8	4.09
80-84	20	17.9	4.00	14	36.7	9.80	6	8.2	3.33
85+	24	25.3	5.17	9	37.1	12.35	15	21.3	5.49
Total deaths	229			140			89		
Connecticut									
. Crude rate		3.9	-		4.9	-		3.0	-
. AAMR (1940 Std. mil.)		2.4	0.18		3.6	0.32		1.4	0.18
. AAMR (1970 Std. mil.)		3.0	0.21		4.4	0.38		1.9	0.21
. AAMR (2000 Std. mil.)		3.7	0.24		5.4	0.47		2.3	0.25
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.3	-		3.6	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.25	1	0.5	0.48	0	0.0	0.00
25-29	2	0.4	0.30	2	0.8	0.60	0	0.0	0.00
30-34	3	0.6	0.33	1	0.4	0.38	2	0.8	0.54
35-39	11	2.2	0.67	5	2.0	0.91	6	2.4	0.99
40-44	17	3.8	0.91	9	4.1	1.35	8	3.5	1.23
45-49	28	7.3	1.39	17	9.0	2.19	11	5.7	1.71
50-54	40	13.7	2.17	21	14.8	3.22	19	12.8	2.93
55-59	73	28.3	3.31	46	36.8	5.42	27	20.3	3.92
60-64	119	44.5	4.08	64	50.2	6.28	55	39.3	5.30
65-69	184	71.8	5.29	105	91.2	8.90	79	56.0	6.30
70-74	272	119.8	7.27	147	152.0	12.54	125	95.9	8.58
75-79	286	162.4	9.60	154	221.1	17.82	132	124.0	10.79
80-84	240	214.9	13.87	112	293.4	27.72	128	174.2	15.39
85+	275	290.2	17.50	91	374.6	39.27	184	261.1	19.25
Total deaths	1,551			775			776		
Connecticut									
. Crude rate		26.5	-		27.3	-		25.7	-
. AAMR (1940 Std. mil.)		12.9	0.37		16.3	0.62		10.5	0.45
. AAMR (1970 Std. mil.)		18.2	0.48		23.2	0.84		14.8	0.57
. AAMR (2000 Std. mil.)		24.0	0.61		30.9	1.15		19.7	0.72
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.4	-		16.4	-		11.3	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: CT rate is not significantly different



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Rectal cancer (154)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.4	0.29	1	0.4	0.41	1	0.4	0.40
40-44	3	0.7	0.38	1	0.5	0.45	2	0.9	0.62
45-49	7	1.8	0.69	5	2.7	1.19	2	1.0	0.73
50-54	8	2.7	0.97	4	2.8	1.40	4	2.7	1.34
55-59	10	3.9	1.23	6	4.8	1.96	4	3.0	1.51
60-64	24	9.0	1.83	11	8.6	2.60	13	9.3	2.58
65-69	29	11.3	2.10	19	16.5	3.79	10	7.1	2.24
70-74	41	18.1	2.82	23	23.8	4.96	18	13.8	3.26
75-79	47	26.7	3.89	27	38.8	7.46	20	18.8	4.20
80-84	34	30.4	5.22	17	44.5	10.80	17	23.1	5.61
85+	43	45.4	6.92	14	57.6	15.40	29	41.1	7.64
Total deaths	248			128			120		
Connecticut									
. Crude rate		4.2	-		4.5	-		4.0	-
. AAMR (1940 Std. mil.)		2.1	0.15		2.7	0.25		1.7	0.19
. AAMR (1970 Std. mil.)		2.9	0.19		3.8	0.34		2.4	0.23
. AAMR (2000 Std. mil.)		3.9	0.25		5.1	0.46		3.1	0.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.5	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.4	0.27	1	0.4	0.38	1	0.4	0.38
35-39	4	0.8	0.40	1	0.4	0.41	3	1.2	0.70
40-44	4	0.9	0.44	2	0.9	0.64	2	0.9	0.62
45-49	14	3.7	0.98	10	5.3	1.68	4	2.1	1.03
50-54	18	6.2	1.46	11	7.7	2.33	7	4.7	1.78
55-59	38	14.7	2.39	20	16.0	3.57	18	13.6	3.20
60-64	63	23.6	2.97	31	24.3	4.37	32	22.9	4.04
65-69	95	37.1	3.80	47	40.8	5.95	48	34.0	4.91
70-74	124	54.6	4.91	64	66.2	8.27	60	46.1	5.95
75-79	121	68.7	6.25	44	63.2	9.53	77	72.3	8.24
80-84	98	87.8	8.87	37	96.9	15.93	61	83.0	10.63
85+	108	114.0	10.97	26	107.0	20.99	82	116.3	12.85
Total deaths	689			294			395		
Connecticut									
. Crude rate		11.8	-		10.3	-		13.1	-
. AAMR (1940 Std. mil.)		6.0	0.25		6.5	0.40		5.5	0.33
. AAMR (1970 Std. mil.)		8.3	0.33		8.9	0.52		7.7	0.41
. AAMR (2000 Std. mil.)		10.6	0.41		11.3	0.68		10.1	0.51
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.3	-		7.5	-		5.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.2	0.21	0	0.0	0.00	1	0.4	0.43
30-34	7	1.3	0.50	4	1.5	0.76	3	1.1	0.66
35-39	12	2.4	0.70	9	3.7	1.22	3	1.2	0.70
40-44	44	9.7	1.47	25	11.3	2.26	19	8.3	1.90
45-49	100	26.2	2.62	51	27.1	3.80	49	25.3	3.61
50-54	159	54.6	4.33	77	54.1	6.16	82	55.1	6.09
55-59	250	97.0	6.13	146	116.7	9.66	104	78.4	7.68
60-64	412	154.1	7.59	246	193.1	12.31	166	118.6	9.21
65-69	623	243.2	9.74	388	337.0	17.11	235	166.6	10.87
70-74	649	285.9	11.22	382	395.0	20.21	267	204.9	12.54
75-79	576	327.0	13.63	333	478.2	26.20	243	228.2	14.64
80-84	379	339.4	17.43	240	628.7	40.58	139	189.1	16.04
85+	229	241.6	15.97	131	539.3	47.12	98	139.0	14.05
Total deaths	3,441			2,032			1,409		
Connecticut									
. Crude rate		58.7	-		71.5	-		46.7	-
. AAMR (1940 Std. mil.)		34.4	0.64		46.0	1.07		26.0	0.78
. AAMR (1970 Std. mil.)		44.4	0.78		61.7	1.38		32.7	0.92
. AAMR (2000 Std. mil.)		53.1	0.91		76.4	1.74		38.5	1.05
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		39.5	-		57.4	-		25.8	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	3	0.7	0.43	2	1.0	0.68	1	0.5	0.50
25-29	1	0.2	0.21	0	0.0	0.00	1	0.4	0.43
30-34	5	0.9	0.42	3	1.1	0.66	2	0.8	0.54
35-39	10	2.0	0.64	7	2.8	1.08	3	1.2	0.70
40-44	12	2.7	0.77	7	3.2	1.19	5	2.2	0.97
45-49	8	2.1	0.74	5	2.7	1.19	3	1.5	0.89
50-54	11	3.8	1.14	6	4.2	1.72	5	3.4	1.50
55-59	13	5.0	1.40	7	5.6	2.11	6	4.5	1.85
60-64	18	6.7	1.59	11	8.6	2.60	7	5.0	1.89
65-69	28	10.9	2.07	13	11.3	3.13	15	10.6	2.75
70-74	29	12.8	2.37	21	21.7	4.74	8	6.1	2.17
75-79	19	10.8	2.47	11	15.8	4.76	8	7.5	2.66
80-84	12	10.7	3.10	7	18.3	6.93	5	6.8	3.04
85+	14	14.8	3.95	7	28.8	10.89	7	9.9	3.75
Total deaths	183			107			76		
Connecticut									
. Crude rate		3.1	-		3.8	-		2.5	-
. AAMR (1940 Std. mil.)		2.1	0.17		2.7	0.27		1.6	0.21
. AAMR (1970 Std. mil.)		2.4	0.18		3.2	0.31		1.8	0.22
. AAMR (2000 Std. mil.)		2.9	0.22		4.0	0.39		2.2	0.25
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.7	-		1.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	2	0.4	0.30	-	-	-	2	0.9	0.61
30-34	5	0.9	0.42	-	-	-	5	1.9	0.85
35-39	33	6.7	1.16	-	-	-	33	13.3	2.32
40-44	45	10.0	1.49	-	-	-	45	19.6	2.92
45-49	57	14.9	1.98	-	-	-	57	29.4	3.89
50-54	67	23.0	2.81	-	-	-	67	45.0	5.50
55-59	95	36.8	3.78	-	-	-	95	71.6	7.34
60-64	117	43.8	4.05	-	-	-	117	83.6	7.73
65-69	149	58.2	4.76	-	-	-	149	105.6	8.65
70-74	152	67.0	5.43	-	-	-	152	116.7	9.46
75-79	145	82.3	6.84	-	-	-	145	136.2	11.31
80-84	119	106.6	9.77	-	-	-	119	161.9	14.84
85+	128	135.1	11.94	-	-	-	128	181.6	16.05
Total deaths	1,114			-			1,114		
Connecticut									
. Crude rate		-	-		-	-		36.9	-
. AAMR (1940 Std. mil.)		-	-		-	-		21.3	0.73
. AAMR (1970 Std. mil.)		-	-		-	-		25.8	0.82
. AAMR (2000 Std. mil.)		-	-		-	-		31.2	0.96
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		22.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is lower
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is not significantly different



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	2	0.4	0.30	-	-	-	2	0.9	0.61
30-34	2	0.4	0.27	-	-	-	2	0.8	0.54
35-39	5	1.0	0.45	-	-	-	5	2.0	0.90
40-44	5	1.1	0.50	-	-	-	5	2.2	0.97
45-49	9	2.4	0.79	-	-	-	9	4.6	1.55
50-54	15	5.2	1.33	-	-	-	15	10.1	2.60
55-59	3	1.2	0.67	-	-	-	3	2.3	1.31
60-64	2	0.7	0.53	-	-	-	2	1.4	1.01
65-69	7	2.7	1.03	-	-	-	7	5.0	1.88
70-74	7	3.1	1.17	-	-	-	7	5.4	2.03
75-79	7	4.0	1.50	-	-	-	7	6.6	2.48
80-84	7	6.3	2.37	-	-	-	7	9.5	3.60
85+	8	8.4	2.98	-	-	-	8	11.4	4.01
Total deaths	79			-			79		
Connecticut									
. Crude rate		-	-		-	-		2.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.8	0.23
. AAMR (1970 Std. mil.)		-	-		-	-		2.0	0.23
. AAMR (2000 Std. mil.)		-	-		-	-		2.4	0.28
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is higher
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is higher

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	0	0.0	0.00	-	-	-	0	0.0	0.00
45-49	0	0.0	0.00	-	-	-	0	0.0	0.00
50-54	1	0.3	0.34	-	-	-	1	0.7	0.67
55-59	7	2.7	1.03	-	-	-	7	5.3	1.99
60-64	7	2.6	0.99	-	-	-	7	5.0	1.89
65-69	17	6.6	1.61	-	-	-	17	12.1	2.92
70-74	15	6.6	1.71	-	-	-	15	11.5	2.97
75-79	7	4.0	1.50	-	-	-	7	6.6	2.48
80-84	9	8.1	2.69	-	-	-	9	12.2	4.08
85+	13	13.7	3.80	-	-	-	13	18.4	5.12
Total deaths	76			-			76		
Connecticut									
. Crude rate		-	-		-	-		2.5	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	0.16
. AAMR (1970 Std. mil.)		-	-		-	-		1.6	0.20
. AAMR (2000 Std. mil.)		-	-		-	-		2.0	0.23
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	1	0.2	0.25	-	-	-	1	0.5	0.50
25-29	1	0.2	0.21	-	-	-	1	0.4	0.43
30-34	2	0.4	0.27	-	-	-	2	0.8	0.54
35-39	1	0.2	0.20	-	-	-	1	0.4	0.40
40-44	8	1.8	0.63	-	-	-	8	3.5	1.23
45-49	14	3.7	0.98	-	-	-	14	7.2	1.93
50-54	15	5.2	1.33	-	-	-	15	10.1	2.60
55-59	25	9.7	1.94	-	-	-	25	18.8	3.77
60-64	41	15.3	2.39	-	-	-	41	29.3	4.58
65-69	49	19.1	2.73	-	-	-	49	34.7	4.96
70-74	53	23.3	3.21	-	-	-	53	40.7	5.59
75-79	50	28.4	4.01	-	-	-	50	47.0	6.64
80-84	48	43.0	6.20	-	-	-	48	65.3	9.43
85+	41	43.3	6.76	-	-	-	41	58.2	9.08
Total deaths	349			-			349		
Connecticut									
. Crude rate		-	-		-	-		11.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		6.2	0.38
. AAMR (1970 Std. mil.)		-	-		-	-		7.8	0.44
. AAMR (2000 Std. mil.)		-	-		-	-		9.5	0.52
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Bladder cancer (188)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.2	0.20	1	0.4	0.41	0	0.0	0.00
40-44	1	0.2	0.22	0	0.0	0.00	1	0.4	0.44
45-49	5	1.3	0.59	2	1.1	0.75	3	1.5	0.89
50-54	3	1.0	0.60	0	0.0	0.00	3	2.0	1.16
55-59	6	2.3	0.95	5	4.0	1.79	1	0.8	0.75
60-64	26	9.7	1.91	20	15.7	3.51	6	4.3	1.75
65-69	34	13.3	2.28	26	22.6	4.43	8	5.7	2.00
70-74	47	20.7	3.02	38	39.3	6.37	9	6.9	2.30
75-79	68	38.6	4.68	45	64.6	9.63	23	21.6	4.50
80-84	68	60.9	7.38	37	96.9	15.93	31	42.2	7.58
85+	67	70.7	8.64	30	123.5	22.55	37	52.5	8.63
Total deaths	326			204			122		
Connecticut									
. Crude rate		5.6	-		7.2	-		4.0	-
. AAMR (1940 Std. mil.)		2.4	0.15		3.9	0.29		1.4	0.15
. AAMR (1970 Std. mil.)		3.6	0.21		6.0	0.42		2.1	0.20
. AAMR (2000 Std. mil.)		5.0	0.28		8.3	0.60		3.0	0.28
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.7	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.25	0	0.0	0.00	1	0.5	0.51
5-9	4	1.1	0.54	1	0.5	0.53	3	1.7	0.97
10-14	4	1.2	0.59	3	1.7	0.99	1	0.6	0.60
15-19	2	0.6	0.42	2	1.2	0.81	0	0.0	0.00
20-24	5	1.2	0.55	4	1.9	0.97	1	0.5	0.50
25-29	4	0.9	0.43	2	0.8	0.60	2	0.9	0.61
30-34	9	1.7	0.57	4	1.5	0.76	5	1.9	0.85
35-39	9	1.8	0.61	7	2.8	1.08	2	0.8	0.57
40-44	15	3.3	0.86	12	5.4	1.56	3	1.3	0.75
45-49	18	4.7	1.11	11	5.9	1.76	7	3.6	1.36
50-54	16	5.5	1.37	6	4.2	1.72	10	6.7	2.13
55-59	24	9.3	1.90	13	10.4	2.88	11	8.3	2.50
60-64	31	11.6	2.08	19	14.9	3.42	12	8.6	2.48
65-69	49	19.1	2.73	28	24.3	4.60	21	14.9	3.25
70-74	78	34.4	3.89	45	46.5	6.94	33	25.3	4.41
75-79	64	36.3	4.54	30	43.1	7.87	34	31.9	5.48
80-84	57	51.0	6.76	29	76.0	14.11	28	38.1	7.20
85+	71	74.9	8.89	27	111.2	21.39	44	62.4	9.41
Total deaths	461			243			218		
Connecticut									
. Crude rate		7.9	-		8.5	-		7.2	-
. AAMR (1940 Std. mil.)		4.5	0.24		5.7	0.39		3.6	0.30
. AAMR (1970 Std. mil.)		5.8	0.28		7.4	0.48		4.6	0.34
. AAMR (2000 Std. mil.)		7.3	0.34		9.5	0.63		5.8	0.40
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.9	-		6.2	-		4.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Diabetes mellitus (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.3	0.27	0	0.0	0.00	1	0.6	0.56
10-14	1	0.3	0.29	1	0.6	0.57	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	3	0.6	0.37	1	0.4	0.42	2	0.9	0.61
30-34	8	1.5	0.54	4	1.5	0.76	4	1.5	0.76
35-39	12	2.4	0.70	7	2.8	1.08	5	2.0	0.90
40-44	12	2.7	0.77	9	4.1	1.35	3	1.3	0.75
45-49	15	3.9	1.01	11	5.9	1.76	4	2.1	1.03
50-54	20	6.9	1.54	14	9.8	2.63	6	4.0	1.65
55-59	36	14.0	2.33	23	18.4	3.83	13	9.8	2.72
60-64	69	25.8	3.11	38	29.8	4.84	31	22.2	3.98
65-69	88	34.3	3.66	49	42.6	6.08	39	27.6	4.43
70-74	138	60.8	5.18	66	68.3	8.40	72	55.3	6.51
75-79	168	95.4	7.36	74	106.3	12.35	94	88.3	9.10
80-84	144	129.0	10.75	58	151.9	19.95	86	117.0	12.62
85+	167	176.2	13.64	39	160.6	25.71	128	181.6	16.05
Total deaths	882			394			488		
Connecticut									
. Crude rate		15.1	-		13.9	-		16.2	-
. AAMR (1940 Std. mil.)		7.3	0.28		8.6	0.46		6.1	0.34
. AAMR (1970 Std. mil.)		10.2	0.36		11.8	0.60		8.9	0.43
. AAMR (2000 Std. mil.)		13.7	0.46		15.5	0.81		12.2	0.56
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.9	-		12.7	-		11.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.3	0.27	0	0.0	0.00	1	0.6	0.56
10-14	1	0.3	0.29	1	0.6	0.57	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.2	0.19	0	0.0	0.00	1	0.4	0.38
35-39	1	0.2	0.20	1	0.4	0.41	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	1	0.4	0.37	0	0.0	0.00	1	0.7	0.71
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	1	0.4	0.44	1	1.0	1.03	0	0.0	0.00
75-79	2	1.1	0.80	2	2.9	2.03	0	0.0	0.00
80-84	2	1.8	1.27	2	5.2	3.70	0	0.0	0.00
85+	1	1.1	1.06	1	4.1	4.12	0	0.0	0.00
Total deaths	11			8			3		
Connecticut									
. Crude rate		0.2	-		0.3	-		0.1	-
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. AAMR (1970 Std. mil.)		-	-		-	-		-	-
. AAMR (2000 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.5	-		0.6	-		0.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									No comparison is available
. THE U.S. HP 2000 target:									No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Alzheimer's diseases (331.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	1	0.3	0.34	0	0.0	0.00	1	0.7	0.67
55-59	2	0.8	0.55	1	0.8	0.80	1	0.8	0.75
60-64	7	2.6	0.99	5	3.9	1.75	2	1.4	1.01
65-69	23	9.0	1.87	13	11.3	3.13	10	7.1	2.24
70-74	36	15.9	2.64	13	13.4	3.73	23	17.7	3.68
75-79	54	30.7	4.17	30	43.1	7.87	24	22.5	4.60
80-84	80	71.6	8.01	32	83.8	14.82	48	65.3	9.43
85+	168	177.3	13.68	46	189.4	27.92	122	173.1	15.67
Total deaths	371			140			231		
Connecticut									
. Crude rate		6.3	-		4.9	-		7.7	-
. AAMR (1940 Std. mil.)		2.0	0.12		2.3	0.20		1.8	0.14
. AAMR (1970 Std. mil.)		3.6	0.19		4.1	0.35		3.3	0.23
. AAMR (2000 Std. mil.)		5.8	0.30		6.6	0.58		5.4	0.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.1	-		2.2	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available







## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.5	0.35	1	0.5	0.48	1	0.5	0.51
5-9	1	0.3	0.27	0	0.0	0.00	1	0.6	0.56
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	0.5	0.35	0	0.0	0.00	2	1.0	0.71
25-29	8	1.7	0.60	7	3.0	1.12	1	0.4	0.43
30-34	15	2.8	0.74	14	5.3	1.42	1	0.4	0.38
35-39	46	9.3	1.37	37	15.1	2.48	9	3.6	1.21
40-44	107	23.7	2.29	85	38.3	4.16	22	9.6	2.04
45-49	184	48.2	3.55	162	86.2	6.77	22	11.3	2.42
50-54	253	86.9	5.46	199	139.8	9.91	54	36.3	4.94
55-59	406	157.5	7.82	303	242.2	13.91	103	77.6	7.65
60-64	767	286.9	10.36	563	441.9	18.62	204	145.8	10.21
65-69	1,210	472.3	13.58	821	713.1	24.89	389	275.7	13.98
70-74	1,658	730.4	17.94	1,006	1040.3	32.80	652	500.4	19.60
75-79	2,009	1140.7	25.45	1,113	1598.2	47.91	896	841.4	28.11
80-84	2,273	2035.5	42.70	1,009	2643.2	83.21	1,264	1719.9	48.38
85+	4,502	4750.3	70.80	1,303	5364.1	148.60	3,199	4538.7	80.25
Total deaths	13,443			6,623			6,820		
Connecticut									
. Crude rate		229.5	-		233.0	-		226.1	-
. AAMR (1940 Std. mil.)		93.8	0.95		133.5	1.75		62.5	0.96
. AAMR (1970 Std. mil.)		145.3	1.30		199.2	2.46		105.5	1.37
. AAMR (2000 Std. mil.)		211.0	1.83		279.8	3.57		162.9	1.99
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		117.8	-		163.5	-		81.6	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.2	0.19	1	0.4	0.38	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.2	0.22	1	0.5	0.45	0	0.0	0.00
45-49	2	0.5	0.37	1	0.5	0.53	1	0.5	0.52
50-54	2	0.7	0.49	2	1.4	0.99	0	0.0	0.00
55-59	6	2.3	0.95	4	3.2	1.60	2	1.5	1.07
60-64	17	6.4	1.54	10	7.8	2.48	7	5.0	1.89
65-69	39	15.2	2.44	21	18.2	3.98	18	12.8	3.01
70-74	77	33.9	3.87	28	29.0	5.47	49	37.6	5.37
75-79	111	63.0	5.98	52	74.7	10.35	59	55.4	7.21
80-84	165	147.8	11.50	66	172.9	21.28	99	134.7	13.54
85+	420	443.2	21.62	112	461.1	43.57	308	437.0	24.90
Total deaths	841			298			543		
Connecticut									
. Crude rate		14.4	-		10.5	-		18.0	-
. AAMR (1940 Std. mil.)		4.3	0.17		4.8	0.30		4.0	0.21
. AAMR (1970 Std. mil.)		8.1	0.28		8.8	0.51		7.6	0.34
. AAMR (2000 Std. mil.)		13.3	0.46		14.4	0.87		12.6	0.54
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.7	-		6.4	-		5.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.4	0.27	1	0.4	0.38	1	0.4	0.38
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	5	1.3	0.59	3	1.6	0.92	2	1.0	0.73
50-54	5	1.7	0.77	3	2.1	1.22	2	1.3	0.95
55-59	5	1.9	0.87	4	3.2	1.60	1	0.8	0.75
60-64	9	3.4	1.12	6	4.7	1.92	3	2.1	1.24
65-69	22	8.6	1.83	14	12.2	3.25	8	5.7	2.00
70-74	19	8.4	1.92	8	8.3	2.92	11	8.4	2.55
75-79	39	22.1	3.55	20	28.7	6.42	19	17.8	4.09
80-84	53	47.5	6.52	21	55.0	12.00	32	43.5	7.70
85+	67	70.7	8.64	19	78.2	17.94	48	68.1	9.83
Total deaths	226			99			127		
Connecticut									
. Crude rate		3.9	-		3.5	-		4.2	-
. AAMR (1940 Std. mil.)		1.6	0.12		1.9	0.21		1.3	0.14
. AAMR (1970 Std. mil.)		2.4	0.17		3.0	0.30		2.0	0.19
. AAMR (2000 Std. mil.)		3.6	0.24		4.3	0.44		3.1	0.28
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.8	-		2.2	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.5	0.35	1	0.5	0.48	1	0.5	0.51
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.30	1	0.6	0.58	0	0.0	0.00
20-24	1	0.2	0.25	1	0.5	0.48	0	0.0	0.00
25-29	5	1.1	0.48	0	0.0	0.00	5	2.2	0.96
30-34	8	1.5	0.54	5	1.9	0.85	3	1.1	0.66
35-39	15	3.0	0.78	8	3.3	1.15	7	2.8	1.07
40-44	25	5.5	1.11	19	8.6	1.97	6	2.6	1.07
45-49	33	8.6	1.50	13	6.9	1.92	20	10.3	2.31
50-54	46	15.8	2.33	23	16.2	3.37	23	15.5	3.22
55-59	55	21.3	2.88	29	23.2	4.30	26	19.6	3.84
60-64	92	34.4	3.59	46	36.1	5.32	46	32.9	4.85
65-69	158	61.7	4.91	88	76.4	8.15	70	49.6	5.93
70-74	340	149.8	8.12	180	186.1	13.87	160	122.8	9.71
75-79	491	278.8	12.58	245	351.8	22.48	246	231.0	14.73
80-84	638	571.3	22.62	235	615.6	40.16	403	548.4	27.32
85+	1,337	1410.7	38.58	337	1387.3	75.57	1,000	1418.8	44.87
Total deaths	3,247			1,231			2,016		
Connecticut									
. Crude rate		55.4	-		43.3	-		66.8	-
. AAMR (1940 Std. mil.)		19.6	0.41		22.0	0.67		17.8	0.52
. AAMR (1970 Std. mil.)		32.9	0.60		36.4	1.04		30.4	0.73
. AAMR (2000 Std. mil.)		51.4	0.90		55.8	1.65		48.3	1.08
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		26.8	-		29.3	-		24.6	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is higher  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Atherosclerosis (440)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.2	0.22	1	0.5	0.45	0	0.0	0.00
45-49	1	0.3	0.26	0	0.0	0.00	1	0.5	0.52
50-54	3	1.0	0.60	1	0.7	0.70	2	1.3	0.95
55-59	5	1.9	0.87	3	2.4	1.38	2	1.5	1.07
60-64	8	3.0	1.06	5	3.9	1.75	3	2.1	1.24
65-69	18	7.0	1.66	12	10.4	3.01	6	4.3	1.74
70-74	28	12.3	2.33	14	14.5	3.87	14	10.7	2.87
75-79	42	23.8	3.68	24	34.5	7.03	18	16.9	3.98
80-84	58	51.9	6.82	23	60.3	12.56	35	47.6	8.05
85+	238	251.1	16.28	53	218.2	29.97	185	262.5	19.30
Total deaths	402			136			266		
Connecticut									
. Crude rate		6.9	-		4.8	-		8.8	-
. AAMR (1940 Std. mil.)		2.0	0.12		2.2	0.21		1.8	0.14
. AAMR (1970 Std. mil.)		3.8	0.19		4.0	0.35		3.5	0.23
. AAMR (2000 Std. mil.)		6.4	0.32		6.6	0.58		6.1	0.38
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.6	-		3.0	-		2.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	8	2.0	0.70	5	2.4	1.08	3	1.5	0.89
5-9	1	0.3	0.27	1	0.5	0.53	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.30	1	0.6	0.58	0	0.0	0.00
20-24	3	0.7	0.43	3	1.4	0.84	0	0.0	0.00
25-29	1	0.2	0.21	1	0.4	0.42	0	0.0	0.00
30-34	5	0.9	0.42	5	1.9	0.85	0	0.0	0.00
35-39	9	1.8	0.61	5	2.0	0.91	4	1.6	0.81
40-44	8	1.8	0.63	6	2.7	1.10	2	0.9	0.62
45-49	21	5.5	1.20	17	9.0	2.19	4	2.1	1.03
50-54	15	5.2	1.33	10	7.0	2.22	5	3.4	1.50
55-59	26	10.1	1.98	16	12.8	3.20	10	7.5	2.38
60-64	47	17.6	2.56	25	19.6	3.92	22	15.7	3.35
65-69	70	27.3	3.27	38	33.0	5.35	32	22.7	4.01
70-74	151	66.5	5.41	97	100.3	10.18	54	41.4	5.64
75-79	275	156.1	9.42	151	216.8	17.65	124	116.4	10.46
80-84	392	351.0	17.73	179	468.9	35.05	213	289.8	19.86
85+	1,107	1168.1	35.11	378	1556.1	80.04	729	1034.3	38.31
Total deaths	2,140			938			1,202		
Connecticut									
. Crude rate		36.5	-		33.0	-		39.9	-
. AAMR (1940 Std. mil.)		11.5	0.30		15.6	0.55		8.9	0.33
. AAMR (1970 Std. mil.)		20.7	0.46		28.0	0.92		16.6	0.51
. AAMR (2000 Std. mil.)		34.1	0.74		46.0	1.56		28.1	0.82
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.3	-		17.6	-		10.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.25	1	0.5	0.48	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.6	0.42	1	0.6	0.58	1	0.6	0.61
20-24	1	0.2	0.25	0	0.0	0.00	1	0.5	0.50
25-29	3	0.6	0.37	1	0.4	0.42	2	0.9	0.61
30-34	1	0.2	0.19	1	0.4	0.38	0	0.0	0.00
35-39	2	0.4	0.29	1	0.4	0.41	1	0.4	0.40
40-44	8	1.8	0.63	4	1.8	0.90	4	1.7	0.87
45-49	8	2.1	0.74	5	2.7	1.19	3	1.5	0.89
50-54	15	5.2	1.33	6	4.2	1.72	9	6.1	2.02
55-59	42	16.3	2.51	22	17.6	3.75	20	15.1	3.37
60-64	103	38.5	3.80	58	45.5	5.98	45	32.2	4.79
65-69	251	98.0	6.18	139	120.7	10.24	112	79.4	7.50
70-74	357	157.3	8.32	188	194.4	14.18	169	129.7	9.98
75-79	423	240.2	11.68	221	317.4	21.35	202	189.7	13.35
80-84	371	332.2	17.25	205	537.0	37.51	166	225.9	17.53
85+	486	512.8	23.26	215	885.1	60.36	271	384.5	23.36
Total deaths	2,074			1,068			1,006		
Connecticut									
. Crude rate		35.4	-		37.6	-		33.4	-
. AAMR (1940 Std. mil.)		14.8	0.37		19.6	0.63		11.9	0.45
. AAMR (1970 Std. mil.)		22.8	0.51		31.4	0.96		18.0	0.60
. AAMR (2000 Std. mil.)		31.8	0.70		45.6	1.45		24.7	0.79
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		20.1	-		27.0	-		15.4	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.6	0.42	1	0.6	0.58	1	0.6	0.61
20-24	1	0.2	0.25	0	0.0	0.00	1	0.5	0.50
25-29	2	0.4	0.30	1	0.4	0.42	1	0.4	0.43
30-34	1	0.2	0.19	1	0.4	0.38	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	3	0.7	0.38	1	0.5	0.45	2	0.9	0.62
45-49	3	0.8	0.45	2	1.1	0.75	1	0.5	0.52
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	3	1.2	0.67	2	1.6	1.13	1	0.8	0.75
60-64	4	1.5	0.75	3	2.4	1.36	1	0.7	0.71
65-69	12	4.7	1.35	4	3.5	1.74	8	5.7	2.00
70-74	11	4.8	1.46	2	2.1	1.46	9	6.9	2.30
75-79	4	2.3	1.14	1	1.4	1.44	3	2.8	1.63
80-84	7	6.3	2.37	1	2.6	2.62	6	8.2	3.33
85+	21	22.2	4.84	2	8.2	5.82	19	27.0	6.18
Total deaths	74			21			53		
Connecticut									
. Crude rate		1.3	-		0.7	-		1.8	-
. AAMR (1940 Std. mil.)		0.7	0.09		0.6	0.13		0.7	0.13
. AAMR (1970 Std. mil.)		0.9	0.11		0.7	0.15		1.0	0.15
. AAMR (2000 Std. mil.)		1.2	0.14		0.8	0.18		1.3	0.19
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.4	-		1.3	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	7	1.5	0.57	4	1.7	0.85	3	1.3	0.74
30-34	13	2.5	0.68	8	3.0	1.07	5	1.9	0.85
35-39	32	6.5	1.15	20	8.1	1.82	12	4.8	1.40
40-44	40	8.9	1.40	31	14.0	2.51	9	3.9	1.31
45-49	39	10.2	1.64	30	16.0	2.91	9	4.6	1.55
50-54	44	15.1	2.28	36	25.3	4.21	8	5.4	1.90
55-59	54	20.9	2.85	38	30.4	4.93	16	12.1	3.01
60-64	62	23.2	2.95	39	30.6	4.90	23	16.4	3.43
65-69	72	28.1	3.31	37	32.1	5.28	35	24.8	4.19
70-74	72	31.7	3.74	43	44.5	6.78	29	22.3	4.13
75-79	53	30.1	4.13	30	43.1	7.87	23	21.6	4.50
80-84	40	35.8	5.66	19	49.8	11.42	21	28.6	6.24
85+	12	12.7	3.66	5	20.6	9.21	7	9.9	3.75
Total deaths	540			340			200		
Connecticut									
. Crude rate		9.2	-		12.0	-		6.6	-
. AAMR (1940 Std. mil.)		6.6	0.30		9.4	0.53		4.2	0.32
. AAMR (1970 Std. mil.)		7.4	0.33		10.5	0.58		4.8	0.35
. AAMR (2000 Std. mil.)		8.7	0.38		12.4	0.68		5.6	0.41
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.2	-		11.7	-		5.1	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is higher

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Nephritis, nephrotic syndrome/nephrosis (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.25	0	0.0	0.00	1	0.5	0.51
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.25	1	0.5	0.48	0	0.0	0.00
25-29	2	0.4	0.30	2	0.8	0.60	0	0.0	0.00
30-34	1	0.2	0.19	0	0.0	0.00	1	0.4	0.38
35-39	6	1.2	0.50	3	1.2	0.70	3	1.2	0.70
40-44	3	0.7	0.38	3	1.4	0.78	0	0.0	0.00
45-49	4	1.0	0.52	2	1.1	0.75	2	1.0	0.73
50-54	12	4.1	1.19	7	4.9	1.86	5	3.4	1.50
55-59	11	4.3	1.29	7	5.6	2.11	4	3.0	1.51
60-64	29	10.8	2.01	19	14.9	3.42	10	7.1	2.26
65-69	40	15.6	2.47	22	19.1	4.07	18	12.8	3.01
70-74	80	35.2	3.94	43	44.5	6.78	37	28.4	4.67
75-79	89	50.5	5.36	51	73.2	10.25	38	35.7	5.79
80-84	119	106.6	9.77	56	146.7	19.60	63	85.7	10.80
85+	168	177.3	13.68	76	312.9	35.89	92	130.5	13.61
Total deaths	566			292			274		
Connecticut									
. Crude rate		9.7	-		10.3	-		9.1	-
. AAMR (1940 Std. mil.)		3.9	0.19		5.4	0.34		3.0	0.23
. AAMR (1970 Std. mil.)		6.1	0.26		8.7	0.51		4.6	0.30
. AAMR (2000 Std. mil.)		8.9	0.37		13.0	0.79		6.7	0.41
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.4	-		5.5	-		3.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----End stage renal disease (ESRD) (581-588,591,593.8,593.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.25	0	0.0	0.00	1	0.5	0.51
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.25	1	0.5	0.48	0	0.0	0.00
25-29	2	0.4	0.30	2	0.8	0.60	0	0.0	0.00
30-34	1	0.2	0.19	0	0.0	0.00	1	0.4	0.38
35-39	6	1.2	0.50	3	1.2	0.70	3	1.2	0.70
40-44	3	0.7	0.38	3	1.4	0.78	0	0.0	0.00
45-49	4	1.0	0.52	2	1.1	0.75	2	1.0	0.73
50-54	12	4.1	1.19	7	4.9	1.86	5	3.4	1.50
55-59	13	5.0	1.40	7	5.6	2.11	6	4.5	1.85
60-64	34	12.7	2.18	22	17.3	3.68	12	8.6	2.48
65-69	50	19.5	2.76	30	26.1	4.76	20	14.2	3.17
70-74	85	37.4	4.06	44	45.5	6.86	41	31.5	4.91
75-79	98	55.6	5.62	55	79.0	10.65	43	40.4	6.16
80-84	133	119.1	10.33	64	167.7	20.96	69	93.9	11.30
85+	189	199.4	14.51	83	341.7	37.51	106	150.4	14.61
Total deaths	632			323			309		
Connecticut									
. Crude rate		10.8	-		11.4	-		10.2	-
. AAMR (1940 Std. mil.)		4.4	0.20		6.0	0.36		3.4	0.24
. AAMR (1970 Std. mil.)		6.8	0.28		9.6	0.54		5.2	0.32
. AAMR (2000 Std. mil.)		9.9	0.39		14.4	0.83		7.6	0.44
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.0	-		6.3	-		4.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	28	7.0	1.32	18	8.7	2.05	10	5.1	1.62
5-9	13	3.5	0.98	9	4.8	1.59	4	2.2	1.12
10-14	13	3.8	1.06	11	6.3	1.90	2	1.2	0.85
15-19	97	28.8	2.92	71	40.9	4.85	26	15.9	3.12
20-24	134	32.9	2.84	113	54.5	5.13	21	10.5	2.29
25-29	102	21.8	2.16	77	32.7	3.73	25	10.8	2.15
30-34	118	22.4	2.06	95	36.1	3.70	23	8.7	1.82
35-39	95	19.2	1.97	75	30.5	3.52	20	8.1	1.80
40-44	81	17.9	1.99	63	28.4	3.58	18	7.8	1.85
45-49	57	14.9	1.98	39	20.8	3.32	18	9.3	2.19
50-54	44	15.1	2.28	31	21.8	3.91	13	8.7	2.42
55-59	54	20.9	2.85	38	30.4	4.93	16	12.1	3.01
60-64	68	25.4	3.08	47	36.9	5.38	21	15.0	3.27
65-69	84	32.8	3.58	54	46.9	6.38	30	21.3	3.88
70-74	103	45.4	4.47	59	61.0	7.94	44	33.8	5.09
75-79	101	57.3	5.71	50	71.8	10.15	51	47.9	6.71
80-84	123	110.1	9.93	54	141.5	19.25	69	93.9	11.30
85+	236	249.0	16.21	86	354.0	38.18	150	212.8	17.38
Total deaths	1,551			990			561		
Connecticut									
. Crude rate		26.5	-		34.8	-		18.6	-
. AAMR (1940 Std. mil.)		20.4	0.59		30.3	1.03		10.9	0.59
. AAMR (1970 Std. mil.)		21.8	0.60		31.9	1.05		12.7	0.61
. AAMR (2000 Std. mil.)		25.3	0.65		36.7	1.21		15.5	0.67
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		31.0	-		45.4	-		17.1	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.6	0.42	2	1.2	0.81	0	0.0	0.00
20-24	2	0.5	0.35	2	1.0	0.68	0	0.0	0.00
25-29	4	0.9	0.43	3	1.3	0.74	1	0.4	0.43
30-34	4	0.8	0.38	4	1.5	0.76	0	0.0	0.00
35-39	1	0.2	0.20	1	0.4	0.41	0	0.0	0.00
40-44	6	1.3	0.54	5	2.3	1.01	1	0.4	0.44
45-49	6	1.6	0.64	4	2.1	1.06	2	1.0	0.73
50-54	4	1.4	0.69	1	0.7	0.70	3	2.0	1.16
55-59	8	3.1	1.10	7	5.6	2.11	1	0.8	0.75
60-64	17	6.4	1.54	12	9.4	2.72	5	3.6	1.60
65-69	20	7.8	1.75	14	12.2	3.25	6	4.3	1.74
70-74	26	11.5	2.25	17	17.6	4.26	9	6.9	2.30
75-79	34	19.3	3.31	19	27.3	6.26	15	14.1	3.64
80-84	60	53.7	6.94	24	62.9	12.83	36	49.0	8.16
85+	146	154.1	12.75	48	197.6	28.52	98	139.0	14.05
Total deaths	340			163			177		
Connecticut									
. Crude rate		5.8	-		5.7	-		5.9	-
. AAMR (1940 Std. mil.)		2.3	0.15		3.3	0.28		1.5	0.15
. AAMR (1970 Std. mil.)		3.6	0.20		4.9	0.39		2.6	0.21
. AAMR (2000 Std. mil.)		5.4	0.29		7.3	0.60		4.2	0.32
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.5	-		3.7	-		1.6	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.5	0.35	1	0.5	0.48	1	0.5	0.51
5-9	2	0.5	0.39	2	1.1	0.75	0	0.0	0.00
10-14	1	0.3	0.29	1	0.6	0.57	0	0.0	0.00
15-19	6	1.8	0.73	5	2.9	1.29	1	0.6	0.61
20-24	4	1.0	0.49	4	1.9	0.97	0	0.0	0.00
25-29	5	1.1	0.48	4	1.7	0.85	1	0.4	0.43
30-34	7	1.3	0.50	7	2.7	1.01	0	0.0	0.00
35-39	5	1.0	0.45	4	1.6	0.81	1	0.4	0.40
40-44	8	1.8	0.63	8	3.6	1.28	0	0.0	0.00
45-49	2	0.5	0.37	2	1.1	0.75	0	0.0	0.00
50-54	4	1.4	0.69	3	2.1	1.22	1	0.7	0.67
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	3	1.1	0.65	2	1.6	1.11	1	0.7	0.71
65-69	4	1.6	0.78	3	2.6	1.50	1	0.7	0.71
70-74	6	2.6	1.08	5	5.2	2.31	1	0.8	0.77
75-79	2	1.1	0.80	2	2.9	2.03	0	0.0	0.00
80-84	2	1.8	1.27	0	0.0	0.00	2	2.7	1.92
85+	4	4.2	2.11	2	8.2	5.82	2	2.8	2.01
Total deaths	67			55			12		
Connecticut									
. Crude rate		1.1	-		1.9	-		0.4	-
. AAMR (1940 Std. mil.)		1.0	0.14		1.8	0.25		-	-
. AAMR (1970 Std. mil.)		1.0	0.13		1.8	0.25		-	-
. AAMR (2000 Std. mil.)		1.1	0.14		1.9	0.27		-	-
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.7	-		2.9	-		0.5	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	7	1.7	0.66	4	1.9	0.97	3	1.5	0.89
5-9	2	0.5	0.39	1	0.5	0.53	1	0.6	0.56
10-14	2	0.6	0.41	1	0.6	0.57	1	0.6	0.60
15-19	2	0.6	0.42	1	0.6	0.58	1	0.6	0.61
20-24	4	1.0	0.49	2	1.0	0.68	2	1.0	0.71
25-29	1	0.2	0.21	0	0.0	0.00	1	0.4	0.43
30-34	1	0.2	0.19	0	0.0	0.00	1	0.4	0.38
35-39	2	0.4	0.29	2	0.8	0.58	0	0.0	0.00
40-44	3	0.7	0.38	1	0.5	0.45	2	0.9	0.62
45-49	2	0.5	0.37	0	0.0	0.00	2	1.0	0.73
50-54	3	1.0	0.60	2	1.4	0.99	1	0.7	0.67
55-59	4	1.6	0.78	2	1.6	1.13	2	1.5	1.07
60-64	2	0.7	0.53	1	0.8	0.78	1	0.7	0.71
65-69	4	1.6	0.78	3	2.6	1.50	1	0.7	0.71
70-74	8	3.5	1.25	2	2.1	1.46	6	4.6	1.88
75-79	4	2.3	1.14	2	2.9	2.03	2	1.9	1.33
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	2	2.1	1.49	2	8.2	5.82	0	0.0	0.00
Total deaths	53			26			27		
Connecticut									
. Crude rate		0.9	-		0.9	-		0.9	-
. AAMR (1940 Std. mil.)		0.8	0.12		0.8	0.17		0.8	0.17
. AAMR (1970 Std. mil.)		0.9	0.13		0.9	0.19		0.9	0.18
. AAMR (2000 Std. mil.)		0.9	0.12		1.0	0.20		0.9	0.17
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.4	-		1.8	-		1.1	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	5	1.5	0.66	4	2.3	1.15	1	0.6	0.60
15-19	31	9.2	1.65	26	15.0	2.94	5	3.1	1.37
20-24	62	15.2	1.93	59	28.5	3.71	3	1.5	0.87
25-29	68	14.5	1.76	55	23.4	3.15	13	5.6	1.55
30-34	69	13.1	1.58	52	19.8	2.74	17	6.5	1.57
35-39	63	12.8	1.61	50	20.3	2.88	13	5.2	1.45
40-44	62	13.7	1.74	50	22.6	3.19	12	5.2	1.51
45-49	49	12.8	1.83	41	21.8	3.41	8	4.1	1.46
50-54	25	8.6	1.72	17	11.9	2.90	8	5.4	1.90
55-59	29	11.2	2.09	24	19.2	3.92	5	3.8	1.68
60-64	35	13.1	2.21	27	21.2	4.08	8	5.7	2.02
65-69	31	12.1	2.17	27	23.5	4.51	4	2.8	1.42
70-74	24	10.6	2.16	23	23.8	4.96	1	0.8	0.77
75-79	26	14.8	2.90	19	27.3	6.26	7	6.6	2.48
80-84	20	17.9	4.00	18	47.2	11.11	2	2.7	1.92
85+	12	12.7	3.66	9	37.1	12.35	3	4.3	2.46
Total deaths	611			501			110		
Connecticut									
. Crude rate		10.4	-		17.6	-		3.6	-
. AAMR (1940 Std. mil.)		9.5	0.41		16.1	0.75		3.3	0.33
. AAMR (1970 Std. mil.)		9.1	0.38		15.6	0.72		3.1	0.31
. AAMR (2000 Std. mil.)		10.0	0.41		17.3	0.79		3.5	0.34
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.3	-		18.9	-		4.4	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	1.0	0.50	1	0.5	0.48	3	1.5	0.89
5-9	2	0.5	0.39	1	0.5	0.53	1	0.6	0.56
10-14	3	0.9	0.51	2	1.1	0.81	1	0.6	0.60
15-19	26	7.7	1.51	22	12.7	2.70	4	2.4	1.22
20-24	36	8.8	1.47	29	14.0	2.60	7	3.5	1.32
25-29	35	7.5	1.26	27	11.5	2.21	8	3.4	1.22
30-34	31	5.9	1.06	25	9.5	1.90	6	2.3	0.93
35-39	21	4.3	0.93	13	5.3	1.47	8	3.2	1.14
40-44	7	1.6	0.59	5	2.3	1.01	2	0.9	0.62
45-49	8	2.1	0.74	8	4.3	1.51	0	0.0	0.00
50-54	8	2.7	0.97	6	4.2	1.72	2	1.3	0.95
55-59	4	1.6	0.78	2	1.6	1.13	2	1.5	1.07
60-64	6	2.2	0.92	6	4.7	1.92	0	0.0	0.00
65-69	1	0.4	0.39	1	0.9	0.87	0	0.0	0.00
70-74	5	2.2	0.99	4	4.1	2.07	1	0.8	0.77
75-79	3	1.7	0.98	1	1.4	1.44	2	1.9	1.33
80-84	3	2.7	1.55	2	5.2	3.70	1	1.4	1.36
85+	1	1.1	1.06	0	0.0	0.00	1	1.4	1.42
Total deaths	204			155			49		
Connecticut									
. Crude rate		3.5	-		5.5	-		1.6	-
. AAMR (1940 Std. mil.)		3.7	0.27		5.7	0.47		1.7	0.26
. AAMR (1970 Std. mil.)		3.4	0.25		5.2	0.44		1.6	0.24
. AAMR (2000 Std. mil.)		3.4	0.24		5.2	0.42		1.6	0.23
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.8	-		17.4	-		4.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Homicide (E960-E969)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	1.0	0.50	1	0.5	0.48	3	1.5	0.89
5-9	2	0.5	0.39	1	0.5	0.53	1	0.6	0.56
10-14	3	0.9	0.51	2	1.1	0.81	1	0.6	0.60
15-19	26	7.7	1.51	22	12.7	2.70	4	2.4	1.22
20-24	36	8.8	1.47	29	14.0	2.60	7	3.5	1.32
25-29	34	7.3	1.25	26	11.0	2.17	8	3.4	1.22
30-34	31	5.9	1.06	25	9.5	1.90	6	2.3	0.93
35-39	21	4.3	0.93	13	5.3	1.47	8	3.2	1.14
40-44	7	1.6	0.59	5	2.3	1.01	2	0.9	0.62
45-49	8	2.1	0.74	8	4.3	1.51	0	0.0	0.00
50-54	8	2.7	0.97	6	4.2	1.72	2	1.3	0.95
55-59	4	1.6	0.78	2	1.6	1.13	2	1.5	1.07
60-64	6	2.2	0.92	6	4.7	1.92	0	0.0	0.00
65-69	1	0.4	0.39	1	0.9	0.87	0	0.0	0.00
70-74	5	2.2	0.99	4	4.1	2.07	1	0.8	0.77
75-79	3	1.7	0.98	1	1.4	1.44	2	1.9	1.33
80-84	3	2.7	1.55	2	5.2	3.70	1	1.4	1.36
85+	1	1.1	1.06	0	0.0	0.00	1	1.4	1.42
Total deaths	203			154			49		
Connecticut									
. Crude rate		3.5	-		5.4	-		1.6	-
. AAMR (1940 Std. mil.)		3.7	0.27		5.7	0.47		1.7	0.26
. AAMR (1970 Std. mil.)		3.4	0.25		5.2	0.44		1.6	0.24
. AAMR (2000 Std. mil.)		3.4	0.24		5.1	0.42		1.6	0.23
. HP 2000 (1940 Std. mil.)		5.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.7	-		17.1	-		4.5	-
. HP 2000 (1940 Std. mil.)		7.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	10	2.5	0.79	8	3.9	1.37	2	1.0	0.72
5-9	5	1.4	0.61	4	2.1	1.06	1	0.6	0.56
10-14	5	1.5	0.66	4	2.3	1.15	1	0.6	0.60
15-19	70	20.8	2.48	48	27.7	3.99	22	13.5	2.87
20-24	100	24.5	2.45	86	41.5	4.47	14	7.0	1.87
25-29	60	12.8	1.66	41	17.4	2.72	19	8.2	1.87
30-34	60	11.4	1.47	43	16.3	2.49	17	6.5	1.57
35-39	34	6.9	1.18	25	10.2	2.03	9	3.6	1.21
40-44	27	6.0	1.15	18	8.1	1.91	9	3.9	1.31
45-49	23	6.0	1.26	14	7.5	1.99	9	4.6	1.55
50-54	17	5.8	1.42	13	9.1	2.53	4	2.7	1.34
55-59	22	8.5	1.82	14	11.2	2.99	8	6.0	2.13
60-64	21	7.9	1.71	13	10.2	2.83	8	5.7	2.02
65-69	23	9.0	1.87	12	10.4	3.01	11	7.8	2.35
70-74	26	11.5	2.25	11	11.4	3.43	15	11.5	2.97
75-79	28	15.9	3.00	11	15.8	4.76	17	16.0	3.87
80-84	23	20.6	4.29	10	26.2	8.28	13	17.7	4.91
85+	15	15.8	4.09	8	32.9	11.64	7	9.9	3.75
Total deaths	569			383			186		
Connecticut									
. Crude rate		9.7	-		13.5	-		6.2	-
. AAMR (1940 Std. mil.)		9.6	0.43		13.7	0.73		5.4	0.45
. AAMR (1970 Std. mil.)		9.3	0.42		13.2	0.71		5.5	0.44
. AAMR (2000 Std. mil.)		9.4	0.40		13.3	0.69		5.7	0.43
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		16.7	-		23.6	-		9.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.3	0.27	1	0.5	0.53	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.6	0.42	2	1.2	0.81	0	0.0	0.00
20-24	2	0.5	0.35	2	1.0	0.68	0	0.0	0.00
25-29	5	1.1	0.48	3	1.3	0.74	2	0.9	0.61
30-34	8	1.5	0.54	6	2.3	0.93	2	0.8	0.54
35-39	3	0.6	0.35	3	1.2	0.70	0	0.0	0.00
40-44	6	1.3	0.54	5	2.3	1.01	1	0.4	0.44
45-49	8	2.1	0.74	5	2.7	1.19	3	1.5	0.89
50-54	5	1.7	0.77	1	0.7	0.70	4	2.7	1.34
55-59	7	2.7	1.03	7	5.6	2.11	0	0.0	0.00
60-64	17	6.4	1.54	12	9.4	2.72	5	3.6	1.60
65-69	19	7.4	1.70	13	11.3	3.13	6	4.3	1.74
70-74	23	10.1	2.11	15	15.5	4.01	8	6.1	2.17
75-79	31	17.6	3.16	18	25.8	6.09	13	12.2	3.39
80-84	43	38.5	5.87	17	44.5	10.80	26	35.4	6.94
85+	103	108.7	10.71	36	148.2	24.70	67	95.1	11.61
Total deaths	283			146			137		
Connecticut									
. Crude rate		4.8	-		5.1	-		4.5	-
. AAMR (1940 Std. mil.)		2.2	0.16		3.1	0.28		1.4	0.16
. AAMR (1970 Std. mil.)		3.1	0.19		4.4	0.37		2.2	0.20
. AAMR (2000 Std. mil.)		4.5	0.27		6.3	0.54		3.3	0.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.4	-		3.6	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	2	0.5	0.39	1	0.5	0.53	1	0.6	0.56
10-14	4	1.2	0.59	3	1.7	0.99	1	0.6	0.60
15-19	38	11.3	1.83	38	21.9	3.55	0	0.0	0.00
20-24	69	16.9	2.04	61	29.4	3.77	8	4.0	1.41
25-29	64	13.7	1.71	55	23.4	3.15	9	3.9	1.29
30-34	41	7.8	1.22	37	14.1	2.31	4	1.5	0.76
35-39	38	7.7	1.25	29	11.8	2.19	9	3.6	1.21
40-44	31	6.9	1.23	26	11.7	2.30	5	2.2	0.97
45-49	26	6.8	1.34	23	12.2	2.55	3	1.5	0.89
50-54	15	5.2	1.33	11	7.7	2.33	4	2.7	1.34
55-59	17	6.6	1.60	16	12.8	3.20	1	0.8	0.75
60-64	15	5.6	1.45	14	11.0	2.94	1	0.7	0.71
65-69	16	6.2	1.56	15	13.0	3.36	1	0.7	0.71
70-74	15	6.6	1.71	14	14.5	3.87	1	0.8	0.77
75-79	13	7.4	2.05	12	17.2	4.97	1	0.9	0.94
80-84	9	8.1	2.69	9	23.6	7.86	0	0.0	0.00
85+	6	6.3	2.58	5	20.6	9.21	1	1.4	1.42
Total deaths	419			369			50		
Connecticut									
. Crude rate		7.2	-		13.0	-		1.7	-
. AAMR (1940 Std. mil.)		7.1	0.37		12.7	0.69		1.7	0.24
. AAMR (1970 Std. mil.)		6.7	0.34		12.2	0.66		1.5	0.22
. AAMR (2000 Std. mil.)		6.9	0.34		12.6	0.67		1.6	0.23
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.1	-		26.4	-		4.2	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.3	0.29	1	0.6	0.57	0	0.0	0.00
15-19	15	4.5	1.15	15	8.6	2.23	0	0.0	0.00
20-24	33	8.1	1.41	32	15.4	2.73	1	0.5	0.50
25-29	41	8.8	1.37	35	14.9	2.51	6	2.6	1.05
30-34	21	4.0	0.87	19	7.2	1.66	2	0.8	0.54
35-39	22	4.5	0.95	18	7.3	1.73	4	1.6	0.81
40-44	27	6.0	1.15	22	9.9	2.12	5	2.2	0.97
45-49	19	5.0	1.14	16	8.5	2.13	3	1.5	0.89
50-54	10	3.4	1.09	8	5.6	1.99	2	1.3	0.95
55-59	15	5.8	1.50	14	11.2	2.99	1	0.8	0.75
60-64	11	4.1	1.24	11	8.6	2.60	0	0.0	0.00
65-69	15	5.9	1.51	14	12.2	3.25	1	0.7	0.71
70-74	13	5.7	1.59	13	13.4	3.73	0	0.0	0.00
75-79	12	6.8	1.97	11	15.8	4.76	1	0.9	0.94
80-84	7	6.3	2.37	7	18.3	6.93	0	0.0	0.00
85+	5	5.3	2.36	5	20.6	9.21	0	0.0	0.00
Total deaths	267			241			26		
Connecticut									
. Crude rate		4.6	-		8.5	-		0.9	-
. AAMR (1940 Std. mil.)		4.2	0.27		7.8	0.53		0.8	0.16
. AAMR (1970 Std. mil.)		4.0	0.26		7.7	0.51		0.7	0.14
. AAMR (2000 Std. mil.)		4.4	0.27		8.3	0.55		0.8	0.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.8	-		12.0	-		1.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	2	0.5	0.39	1	0.5	0.53	1	0.6	0.56
10-14	3	0.9	0.51	2	1.1	0.81	1	0.6	0.60
15-19	21	6.2	1.36	21	12.1	2.64	0	0.0	0.00
20-24	29	7.1	1.32	23	11.1	2.31	6	3.0	1.22
25-29	22	4.7	1.00	19	8.1	1.85	3	1.3	0.74
30-34	19	3.6	0.83	18	6.8	1.61	1	0.4	0.38
35-39	16	3.2	0.81	11	4.5	1.35	5	2.0	0.90
40-44	3	0.7	0.38	3	1.4	0.78	0	0.0	0.00
45-49	7	1.8	0.69	7	3.7	1.41	0	0.0	0.00
50-54	5	1.7	0.77	3	2.1	1.22	2	1.3	0.95
55-59	2	0.8	0.55	2	1.6	1.13	0	0.0	0.00
60-64	3	1.1	0.65	3	2.4	1.36	0	0.0	0.00
65-69	1	0.4	0.39	1	0.9	0.87	0	0.0	0.00
70-74	1	0.4	0.44	1	1.0	1.03	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	2	1.8	1.27	2	5.2	3.70	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	136			117			19		
Connecticut									
. Crude rate		2.3	-		4.1	-		0.6	-
. AAMR (1940 Std. mil.)		2.6	0.23		4.5	0.43		0.7	0.17
. AAMR (1970 Std. mil.)		2.4	0.22		4.1	0.40		0.6	0.16
. AAMR (2000 Std. mil.)		2.3	0.20		3.9	0.37		0.6	0.15
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.5	-		12.6	-		2.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.25	1	0.5	0.48	0	0.0	0.00
25-29	7	1.5	0.57	5	2.1	0.95	2	0.9	0.61
30-34	22	4.2	0.89	16	6.1	1.52	6	2.3	0.93
35-39	33	6.7	1.16	23	9.4	1.95	10	4.0	1.27
40-44	34	7.5	1.29	27	12.2	2.34	7	3.0	1.15
45-49	31	8.1	1.46	25	13.3	2.66	6	3.1	1.26
50-54	27	9.3	1.79	19	13.3	3.06	8	5.4	1.90
55-59	37	14.4	2.36	29	23.2	4.30	8	6.0	2.13
60-64	28	10.5	1.98	20	15.7	3.51	8	5.7	2.02
65-69	35	13.7	2.31	22	19.1	4.07	13	9.2	2.56
70-74	30	13.2	2.41	17	17.6	4.26	13	10.0	2.77
75-79	17	9.7	2.34	15	21.5	5.56	2	1.9	1.33
80-84	9	8.1	2.69	6	15.7	6.42	3	4.1	2.36
85+	3	3.2	1.83	2	8.2	5.82	1	1.4	1.42
Total deaths	314			227			87		
Connecticut									
. Crude rate		5.4	-		8.0	-		2.9	-
. AAMR (1940 Std. mil.)		4.3	0.25		6.6	0.45		2.2	0.25
. AAMR (1970 Std. mil.)		4.4	0.26		6.8	0.46		2.3	0.25
. AAMR (2000 Std. mil.)		5.2	0.30		8.1	0.54		2.6	0.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.8	-		10.9	-		3.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991-1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.3	0.29	1	0.6	0.57	0	0.0	0.00
15-19	6	1.8	0.73	3	1.7	1.00	3	1.8	1.06
20-24	17	4.2	1.01	15	7.2	1.87	2	1.0	0.71
25-29	33	7.1	1.23	25	10.6	2.12	8	3.4	1.22
30-34	46	8.7	1.29	40	15.2	2.40	6	2.3	0.93
35-39	52	10.5	1.46	40	16.3	2.57	12	4.8	1.40
40-44	34	7.5	1.29	25	11.3	2.26	9	3.9	1.31
45-49	22	5.8	1.23	17	9.0	2.19	5	2.6	1.15
50-54	3	1.0	0.60	0	0.0	0.00	3	2.0	1.16
55-59	10	3.9	1.23	4	3.2	1.60	6	4.5	1.85
60-64	4	1.5	0.75	1	0.8	0.78	3	2.1	1.24
65-69	1	0.4	0.39	0	0.0	0.00	1	0.7	0.71
70-74	3	1.3	0.76	1	1.0	1.03	2	1.5	1.09
75-79	3	1.7	0.98	0	0.0	0.00	3	2.8	1.63
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	4	4.2	2.11	3	12.4	7.13	1	1.4	1.42
Total deaths	239			175			64		
Connecticut									
. Crude rate		4.1	-		6.2	-		2.1	-
. AAMR (1940 Std. mil.)		3.8	0.25		5.7	0.43		2.0	0.26
. AAMR (1970 Std. mil.)		3.3	0.22		4.8	0.37		1.8	0.24
. AAMR (2000 Std. mil.)		3.9	0.25		5.7	0.44		2.1	0.26
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.9	-		5.0	-		2.7	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: CT rate is higher