

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Septicemia (038)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.87	1	1.0	0.97	2	2.1	1.46
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	1.1	0.76	0	0.0	0.00	2	2.2	1.54
25-29	1	0.5	0.46	0	0.0	0.00	1	0.9	0.93
30-34	2	0.8	0.55	0	0.0	0.00	2	1.6	1.10
35-39	5	2.0	0.88	2	1.6	1.12	3	2.4	1.37
40-44	8	3.6	1.27	5	4.6	2.05	3	2.7	1.53
45-49	4	2.0	1.01	4	4.1	2.05	0	0.0	0.00
50-54	4	2.6	1.28	1	1.3	1.31	3	3.8	2.17
55-59	5	3.9	1.76	4	6.5	3.24	1	1.5	1.54
60-64	12	9.6	2.76	6	10.0	4.09	6	9.2	3.74
65-69	29	23.0	4.28	17	29.9	7.26	12	17.4	5.01
70-74	53	46.4	6.37	28	57.0	10.78	25	38.4	7.67
75-79	54	59.2	8.06	30	82.4	15.04	24	43.8	8.94
80-84	75	128.8	14.87	34	169.0	28.99	41	107.5	16.80
85+	127	255.1	22.64	24	185.8	37.93	103	279.4	27.53
Total deaths	384			156			228		
Connecticut									
. Crude rate		13.2	-		11.0	-		15.2	-
. AAMR (1940 Std. mil.)		5.2	0.32		5.9	0.50		4.7	0.41
. AAMR (1970 Std. mil.)		8.0	0.42		8.9	0.72		7.2	0.52
. AAMR (2000 Std. mil.)		11.7	0.60		12.6	1.04		10.8	0.72
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.8	-		4.5	-		3.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is higher
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Oral cavity or pharynx (140.0-149.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.40	1	0.8	0.79	0	0.0	0.00
40-44	2	0.9	0.64	1	0.9	0.92	1	0.9	0.89
45-49	3	1.5	0.87	3	3.1	1.77	0	0.0	0.00
50-54	4	2.6	1.28	3	3.9	2.26	1	1.3	1.25
55-59	7	5.5	2.09	5	8.1	3.62	2	3.1	2.17
60-64	14	11.2	2.98	10	16.7	5.27	4	6.1	3.06
65-69	21	16.7	3.64	15	26.4	6.82	6	8.7	3.55
70-74	19	16.6	3.82	13	26.5	7.35	6	9.2	3.76
75-79	18	19.7	4.65	8	22.0	7.77	10	18.3	5.77
80-84	8	13.7	4.86	4	19.9	9.94	4	10.5	5.25
85+	8	16.1	5.68	5	38.7	17.31	3	8.1	4.70
Total deaths	105			68			37		
Connecticut									
. Crude rate		3.6	-		4.8	-		2.5	-
. AAMR (1940 Std. mil.)		2.1	0.23		3.2	0.41		1.2	0.23
. AAMR (1970 Std. mil.)		2.7	0.27		4.1	0.51		1.6	0.28
. AAMR (2000 Std. mil.)		3.2	0.31		4.9	0.61		1.9	0.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.5	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	3	1.2	0.67	1	0.8	0.78	2	1.6	1.10
35-39	3	1.2	0.69	2	1.6	1.12	1	0.8	0.79
40-44	7	3.2	1.19	2	1.8	1.30	5	4.4	1.98
45-49	13	6.5	1.82	8	8.2	2.89	5	5.0	2.22
50-54	23	14.7	3.07	12	15.7	4.52	11	13.8	4.16
55-59	35	27.6	4.67	19	30.8	7.06	16	24.6	6.14
60-64	77	61.4	7.00	41	68.4	10.68	36	55.0	9.17
65-69	86	68.3	7.37	46	81.0	11.94	40	57.9	9.16
70-74	115	100.7	9.39	68	138.5	16.80	47	72.1	10.52
75-79	127	139.3	12.36	76	208.7	23.94	51	93.1	13.04
80-84	115	197.5	18.41	44	218.7	32.98	71	186.2	22.10
85+	171	343.5	26.27	54	418.1	56.89	117	317.3	29.34
Total deaths	775			373			402		
Connecticut									
. Crude rate		26.6	-		26.4	-		26.9	-
. AAMR (1940 Std. mil.)		12.7	0.52		15.4	0.85		10.7	0.66
. AAMR (1970 Std. mil.)		17.9	0.67		22.0	1.15		15.1	0.82
. AAMR (2000 Std. mil.)		23.7	0.86		29.2	1.56		20.1	1.02
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.0	-		15.8	-		11.1	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	0	0.0	0.00	1	1.0	1.03
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.40	1	0.8	0.79	0	0.0	0.00
40-44	3	1.4	0.78	1	0.9	0.92	2	1.8	1.25
45-49	7	3.5	1.33	5	5.1	2.29	2	2.0	1.40
50-54	15	9.6	2.48	10	13.1	4.13	5	6.3	2.80
55-59	20	15.8	3.53	8	13.0	4.58	12	18.4	5.32
60-64	31	24.7	4.44	22	36.7	7.82	9	13.8	4.59
65-69	46	36.5	5.39	30	52.8	9.64	16	23.2	5.79
70-74	66	57.8	7.11	31	63.2	11.34	35	53.7	9.08
75-79	65	71.3	8.84	34	93.4	16.01	31	56.6	10.16
80-84	46	79.0	11.65	18	89.5	21.09	28	73.4	13.88
85+	54	108.5	14.76	15	116.1	29.99	39	105.8	16.94
Total deaths	355			175			180		
Connecticut									
. Crude rate		12.2	-		12.4	-		12.0	-
. AAMR (1940 Std. mil.)		6.3	0.37		7.7	0.62		5.1	0.46
. AAMR (1970 Std. mil.)		8.5	0.47		10.4	0.80		7.1	0.57
. AAMR (2000 Std. mil.)		10.9	0.58		13.1	1.01		9.2	0.69
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.3	-		7.4	-		5.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung and other respiratory cancer (160-165)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	5	1.9	0.87	2	1.6	1.10	3	2.3	1.35
35-39	7	2.8	1.05	4	3.2	1.59	3	2.4	1.37
40-44	29	13.1	2.43	18	16.5	3.89	11	9.7	2.94
45-49	38	19.1	3.11	19	19.4	4.46	19	18.9	4.33
50-54	69	44.1	5.31	41	53.5	8.36	28	35.1	6.63
55-59	113	89.1	8.38	55	89.1	12.02	58	89.1	11.69
60-64	193	154.0	11.08	103	171.8	16.93	90	137.6	14.50
65-69	307	243.9	13.92	181	318.7	23.69	126	182.4	16.25
70-74	322	281.9	15.71	197	401.4	28.60	125	191.9	17.16
75-79	297	325.7	18.90	171	469.6	35.91	126	230.0	20.49
80-84	202	346.8	24.40	113	561.8	52.85	89	233.4	24.75
85+	132	265.1	23.08	76	588.4	67.50	56	151.9	20.30
Total deaths	1,714			980			734		
Connecticut									
. Crude rate		58.9	-		69.3	-		49.0	-
. AAMR (1940 Std. mil.)		33.3	0.88		43.0	1.45		26.4	1.10
. AAMR (1970 Std. mil.)		43.4	1.08		58.2	1.88		33.6	1.31
. AAMR (2000 Std. mil.)		52.3	1.27		73.0	2.38		39.4	1.48
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		40.8	-		58.1	-		27.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	3	1.2	0.67	2	1.6	1.10	1	0.8	0.78
35-39	6	2.4	0.97	3	2.4	1.38	3	2.4	1.37
40-44	28	12.6	2.38	17	15.6	3.78	11	9.7	2.94
45-49	38	19.1	3.11	19	19.4	4.46	19	18.9	4.33
50-54	67	42.8	5.23	40	52.2	8.26	27	33.8	6.51
55-59	109	85.9	8.23	53	85.9	11.80	56	86.0	11.49
60-64	186	148.4	10.88	98	163.5	16.51	88	134.5	14.34
65-69	293	232.8	13.60	172	302.8	23.09	121	175.2	15.92
70-74	312	273.1	15.46	189	385.1	28.01	123	188.8	17.02
75-79	289	316.9	18.64	167	458.6	35.49	122	222.7	20.16
80-84	196	336.5	24.04	108	536.9	51.66	88	230.8	24.61
85+	128	257.1	22.72	73	565.2	66.15	55	149.2	20.11
Total deaths	1,655			941			714		
Connecticut									
. Crude rate		56.9	-		66.6	-		47.7	-
. AAMR (1940 Std. mil.)		32.1	0.87		41.3	1.42		25.6	1.09
. AAMR (1970 Std. mil.)		41.9	1.06		55.9	1.84		32.6	1.29
. AAMR (2000 Std. mil.)		50.5	1.25		70.1	2.34		38.4	1.46
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		39.2	-		55.7	-		26.5	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	3	1.2	0.67	1	0.8	0.78	2	1.6	1.10
35-39	2	0.8	0.56	1	0.8	0.79	1	0.8	0.79
40-44	5	2.3	1.01	3	2.7	1.59	2	1.8	1.25
45-49	5	2.5	1.13	2	2.0	1.45	3	3.0	1.72
50-54	7	4.5	1.69	5	6.5	2.92	2	2.5	1.77
55-59	10	7.9	2.49	5	8.1	3.62	5	7.7	3.43
60-64	7	5.6	2.11	3	5.0	2.89	4	6.1	3.06
65-69	10	7.9	2.51	7	12.3	4.66	3	4.3	2.51
70-74	13	11.4	3.16	9	18.3	6.11	4	6.1	3.07
75-79	15	16.4	4.25	11	30.2	9.11	4	7.3	3.65
80-84	9	15.5	5.15	4	19.9	9.94	5	13.1	5.87
85+	2	4.0	2.84	0	0.0	0.00	2	5.4	3.84
Total deaths	88			51			37		
Connecticut									
. Crude rate		3.0	-		3.6	-		2.5	-
. AAMR (1940 Std. mil.)		2.0	0.23		2.5	0.37		1.6	0.29
. AAMR (1970 Std. mil.)		2.4	0.26		3.0	0.43		1.8	0.32
. AAMR (2000 Std. mil.)		2.8	0.30		3.7	0.52		2.2	0.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.5	-		1.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	1	0.5	0.53	-	-	-	1	1.1	1.09
25-29	2	0.9	0.65	-	-	-	2	1.9	1.31
30-34	3	1.2	0.67	-	-	-	3	2.3	1.35
35-39	11	4.4	1.31	-	-	-	11	8.7	2.62
40-44	19	8.6	1.96	-	-	-	19	16.8	3.86
45-49	30	15.1	2.76	-	-	-	30	29.8	5.44
50-54	41	26.2	4.09	-	-	-	41	51.4	8.02
55-59	34	26.8	4.60	-	-	-	34	52.2	8.95
60-64	61	48.7	6.23	-	-	-	61	93.3	11.94
65-69	67	53.2	6.50	-	-	-	67	97.0	11.85
70-74	74	64.8	7.53	-	-	-	74	113.6	13.20
75-79	78	85.5	9.68	-	-	-	78	142.4	16.12
80-84	62	106.5	13.52	-	-	-	62	162.6	20.65
85+	84	168.7	18.41	-	-	-	84	227.8	24.86
Total deaths	567			-			567		
Connecticut									
. Crude rate		-	-		-	-		37.9	-
. AAMR (1940 Std. mil.)		-	-		-	-		20.8	1.01
. AAMR (1970 Std. mil.)		-	-		-	-		25.5	1.15
. AAMR (2000 Std. mil.)		-	-		-	-		31.2	1.34
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		21.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is lower
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	2	0.8	0.55	-	-	-	2	1.6	1.10
35-39	2	0.8	0.56	-	-	-	2	1.6	1.12
40-44	3	1.4	0.78	-	-	-	3	2.7	1.53
45-49	4	2.0	1.01	-	-	-	4	4.0	1.98
50-54	3	1.9	1.11	-	-	-	3	3.8	2.17
55-59	2	1.6	1.12	-	-	-	2	3.1	2.17
60-64	2	1.6	1.13	-	-	-	2	3.1	2.16
65-69	1	0.8	0.79	-	-	-	1	1.4	1.45
70-74	1	0.9	0.88	-	-	-	1	1.5	1.53
75-79	3	3.3	1.90	-	-	-	3	5.5	3.16
80-84	3	5.2	2.97	-	-	-	3	7.9	4.54
85+	2	4.0	2.84	-	-	-	2	5.4	3.84
Total deaths	28			-			28		
Connecticut									
. Crude rate		-	-		-	-		1.9	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.3	0.27
. AAMR (1970 Std. mil.)		-	-		-	-		1.4	0.27
. AAMR (2000 Std. mil.)		-	-		-	-		1.7	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	1	0.5	0.45	-	-	-	1	0.9	0.89
45-49	0	0.0	0.00	-	-	-	0	0.0	0.00
50-54	1	0.6	0.64	-	-	-	1	1.3	1.25
55-59	3	2.4	1.37	-	-	-	3	4.6	2.66
60-64	5	4.0	1.78	-	-	-	5	7.6	3.42
65-69	1	0.8	0.79	-	-	-	1	1.4	1.45
70-74	6	5.3	2.14	-	-	-	6	9.2	3.76
75-79	6	6.6	2.69	-	-	-	6	11.0	4.47
80-84	4	6.9	3.43	-	-	-	4	10.5	5.25
85+	7	14.1	5.31	-	-	-	7	19.0	7.18
Total deaths	34			-			34		
Connecticut									
. Crude rate		-	-		-	-		2.3	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.1	0.22
. AAMR (1970 Std. mil.)		-	-		-	-		1.4	0.27
. AAMR (2000 Std. mil.)		-	-		-	-		1.8	0.31
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	1	0.4	0.40	-	-	-	1	0.8	0.79
40-44	8	3.6	1.27	-	-	-	8	7.1	2.51
45-49	4	2.0	1.01	-	-	-	4	4.0	1.98
50-54	9	5.8	1.92	-	-	-	9	11.3	3.76
55-59	15	11.8	3.05	-	-	-	15	23.0	5.95
60-64	19	15.2	3.48	-	-	-	19	29.0	6.66
65-69	22	17.5	3.73	-	-	-	22	31.8	6.79
70-74	26	22.8	4.46	-	-	-	26	39.9	7.83
75-79	22	24.1	5.14	-	-	-	22	40.2	8.56
80-84	29	49.8	9.25	-	-	-	29	76.1	14.13
85+	36	72.3	12.05	-	-	-	36	97.6	16.27
Total deaths	191			-			191		
Connecticut									
. Crude rate		-	-		-	-		12.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		6.3	0.55
. AAMR (1970 Std. mil.)		-	-		-	-		8.2	0.64
. AAMR (2000 Std. mil.)		-	-		-	-		10.2	0.75
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Bladder cancer (188)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	2	0.9	0.64	2	1.8	1.30	0	0.0	0.00
45-49	4	2.0	1.01	4	4.1	2.05	0	0.0	0.00
50-54	2	1.3	0.90	2	2.6	1.85	0	0.0	0.00
55-59	2	1.6	1.12	2	3.2	2.29	0	0.0	0.00
60-64	9	7.2	2.39	7	11.7	4.41	2	3.1	2.16
65-69	15	11.9	3.08	11	19.4	5.84	4	5.8	2.90
70-74	28	24.5	4.63	17	34.6	8.40	11	16.9	5.09
75-79	41	45.0	7.02	25	68.7	13.73	16	29.2	7.30
80-84	24	41.2	8.41	15	74.6	19.25	9	23.6	7.87
85+	40	80.3	12.70	26	201.3	39.48	14	38.0	10.15
Total deaths	167			111			56		
Connecticut									
. Crude rate		5.7	-		7.9	-		3.7	-
. AAMR (1940 Std. mil.)		2.4	0.21		4.1	0.42		1.2	0.18
. AAMR (1970 Std. mil.)		3.6	0.29		6.4	0.61		1.9	0.26
. AAMR (2000 Std. mil.)		5.0	0.39		9.3	0.91		2.7	0.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.7	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Brain & other CNS cancer (191-192)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.87	1	1.0	0.97	2	2.1	1.46
5-9	1	0.5	0.53	1	1.0	1.03	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	1.2	0.86	0	0.0	0.00	2	2.5	1.78
20-24	1	0.5	0.53	0	0.0	0.00	1	1.1	1.09
25-29	1	0.5	0.46	1	0.9	0.92	0	0.0	0.00
30-34	3	1.2	0.67	1	0.8	0.78	2	1.6	1.10
35-39	3	1.2	0.69	1	0.8	0.79	2	1.6	1.12
40-44	4	1.8	0.90	4	3.7	1.83	0	0.0	0.00
45-49	6	3.0	1.23	2	2.0	1.45	4	4.0	1.98
50-54	12	7.7	2.22	8	10.4	3.69	4	5.0	2.51
55-59	13	10.3	2.84	9	14.6	4.86	4	6.1	3.07
60-64	15	12.0	3.09	12	20.0	5.78	3	4.6	2.65
65-69	13	10.3	2.86	6	10.6	4.31	7	10.1	3.83
70-74	26	22.8	4.46	12	24.4	7.06	14	21.5	5.74
75-79	23	25.2	5.26	8	22.0	7.77	15	27.4	7.07
80-84	8	13.7	4.86	6	29.8	12.18	2	5.2	3.71
85+	7	14.1	5.31	3	23.2	13.41	4	10.8	5.42
Total deaths	141			75			66		
Connecticut									
. Crude rate		4.8	-		5.3	-		4.4	-
. AAMR (1940 Std. mil.)		3.3	0.31		3.9	0.48		2.8	0.40
. AAMR (1970 Std. mil.)		3.9	0.34		4.7	0.56		3.3	0.43
. AAMR (2000 Std. mil.)		4.5	0.38		5.5	0.65		3.7	0.47
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.5	-		4.3	-		3.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	2	1.1	0.75	0	0.0	0.00	2	2.2	1.53
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	3	1.8	1.06	1	1.2	1.18	2	2.5	1.78
20-24	1	0.5	0.53	1	1.0	1.05	0	0.0	0.00
25-29	2	0.9	0.65	2	1.8	1.30	0	0.0	0.00
30-34	3	1.2	0.67	2	1.6	1.10	1	0.8	0.78
35-39	4	1.6	0.79	3	2.4	1.38	1	0.8	0.79
40-44	12	5.4	1.56	9	8.2	2.75	3	2.7	1.53
45-49	6	3.0	1.23	2	2.0	1.45	4	4.0	1.98
50-54	10	6.4	2.02	6	7.8	3.20	4	5.0	2.51
55-59	13	10.3	2.84	5	8.1	3.62	8	12.3	4.34
60-64	20	16.0	3.57	11	18.3	5.53	9	13.8	4.59
65-69	26	20.7	4.05	15	26.4	6.82	11	15.9	4.80
70-74	35	30.6	5.18	19	38.7	8.88	16	24.6	6.14
75-79	39	42.8	6.85	26	71.4	14.00	13	23.7	6.58
80-84	26	44.6	8.76	14	69.6	18.60	12	31.5	9.09
85+	40	80.3	12.70	14	108.4	28.97	26	70.5	13.83
Total deaths	242			130			112		
Connecticut									
. Crude rate		8.3	-		9.2	-		7.5	-
. AAMR (1940 Std. mil.)		4.7	0.35		5.8	0.55		3.9	0.45
. AAMR (1970 Std. mil.)		6.0	0.41		7.6	0.68		4.9	0.52
. AAMR (2000 Std. mil.)		7.6	0.49		10.0	0.90		6.0	0.58
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.8	-		6.3	-		4.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.61	0	0.0	0.00	1	1.3	1.26
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	3	1.4	0.80	3	2.8	1.59	0	0.0	0.00
30-34	12	4.7	1.35	4	3.1	1.55	8	6.2	2.21
35-39	10	4.0	1.25	3	2.4	1.38	7	5.5	2.09
40-44	16	7.2	1.80	11	10.1	3.04	5	4.4	1.98
45-49	31	15.6	2.80	20	20.5	4.58	11	10.9	3.29
50-54	46	29.4	4.34	29	37.9	7.03	17	21.3	5.17
55-59	88	69.4	7.40	56	90.8	12.13	32	49.1	8.69
60-64	156	124.4	9.96	83	138.4	15.20	73	111.6	13.06
65-69	231	183.5	12.07	121	213.1	19.37	110	159.2	15.18
70-74	351	307.3	16.40	171	348.4	26.64	180	276.3	20.59
75-79	404	443.0	22.04	196	538.2	38.44	208	379.7	26.33
80-84	410	704.0	34.77	175	870.0	65.77	235	616.4	40.21
85+	470	944.0	43.54	132	1022.0	88.95	338	916.7	49.86
Total deaths	2,229			1,004			1,225		
Connecticut									
. Crude rate		76.6	-		71.0	-		81.8	-
. AAMR (1940 Std. mil.)		34.5	0.84		40.7	1.37		29.8	1.05
. AAMR (1970 Std. mil.)		50.1	1.10		58.9	1.87		43.8	1.35
. AAMR (2000 Std. mil.)		67.7	1.44		79.3	2.57		59.9	1.73
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cardiovascular disease (390-459)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	10	5.0	1.58	5	4.9	2.17	5	5.1	2.30
5-9	3	1.6	0.92	1	1.0	1.03	2	2.2	1.53
10-14	4	2.3	1.14	1	1.1	1.11	3	3.5	2.01
15-19	4	2.4	1.22	4	4.7	2.36	0	0.0	0.00
20-24	3	1.6	0.92	1	1.0	1.05	2	2.2	1.54
25-29	14	6.5	1.73	9	8.3	2.76	5	4.6	2.08
30-34	37	14.4	2.37	22	17.1	3.65	15	11.7	3.02
35-39	44	17.4	2.62	34	27.0	4.63	10	7.9	2.49
40-44	96	43.2	4.41	69	63.2	7.61	27	23.9	4.60
45-49	168	84.6	6.53	125	127.9	11.44	43	42.7	6.51
50-54	198	126.6	9.00	147	192.0	15.83	51	63.9	8.95
55-59	325	256.3	14.22	244	395.5	25.32	81	124.4	13.82
60-64	508	405.2	17.98	348	580.5	31.12	160	244.6	19.34
65-69	846	672.1	23.11	574	1010.7	42.18	272	393.7	23.87
70-74	1,304	1141.5	31.61	759	1546.3	56.13	545	836.5	35.83
75-79	1,832	2008.7	46.93	976	2680.1	85.79	856	1562.5	53.40
80-84	2,033	3490.8	77.42	857	4260.5	145.54	1,176	3084.7	89.95
85+	4,199	8433.9	130.15	1,165	9019.8	264.26	3,034	8228.7	149.39
Total deaths	11,628			5,341			6,287		
Connecticut									
. Crude rate		399.5	-		377.8	-		420.0	-
. AAMR (1940 Std. mil.)		153.5	1.70		206.1	3.05		111.4	1.83
. AAMR (1970 Std. mil.)		241.2	2.34		311.7	4.30		187.8	2.58
. AAMR (2000 Std. mil.)		356.1	3.31		445.0	6.28		292.1	3.72
. HP 2000 (1940 Std. mil.)		84.4	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		182.8	-		238.7	-		137.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diseases of the heart (390-398,402,404-429)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	9	4.5	1.50	4	3.9	1.94	5	5.1	2.30
5-9	2	1.1	0.75	1	1.0	1.03	1	1.1	1.09
10-14	4	2.3	1.14	1	1.1	1.11	3	3.5	2.01
15-19	3	1.8	1.06	3	3.5	2.05	0	0.0	0.00
20-24	2	1.1	0.76	1	1.0	1.05	1	1.1	1.09
25-29	9	4.2	1.39	7	6.4	2.43	2	1.9	1.31
30-34	28	10.9	2.06	20	15.5	3.48	8	6.2	2.21
35-39	34	13.5	2.31	26	20.7	4.05	8	6.3	2.23
40-44	74	33.3	3.87	54	49.4	6.73	20	17.7	3.96
45-49	144	72.5	6.05	111	113.6	10.78	33	32.8	5.70
50-54	166	106.1	8.24	126	164.5	14.66	40	50.1	7.92
55-59	288	227.1	13.38	225	364.7	24.31	63	96.7	12.19
60-64	436	347.8	16.66	306	510.4	29.18	130	198.7	17.43
65-69	708	562.5	21.14	484	852.2	38.74	224	324.3	21.67
70-74	1,064	931.4	28.55	634	1291.7	51.30	430	660.0	31.83
75-79	1,484	1627.2	42.24	807	2216.1	78.01	677	1235.7	47.49
80-84	1,596	2740.4	68.60	683	3395.5	129.92	913	2394.8	79.26
85+	3,222	6471.6	114.01	894	6921.6	231.49	2,328	6313.9	130.86
Total deaths	9,273			4,387			4,886		
Connecticut									
. Crude rate		318.6	-		310.3	-		326.4	-
. AAMR (1940 Std. mil.)		125.2	1.55		172.3	2.81		87.3	1.63
. AAMR (1970 Std. mil.)		194.5	2.11		257.0	3.91		146.7	2.29
. AAMR (2000 Std. mil.)		284.1	2.96		362.0	5.64		227.3	3.28
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		145.3	-		195.5	-		105.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Heart disease-related (390-398,402,404-429)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	34	17.0	2.92	17	16.5	4.01	17	17.5	4.24
5-9	9	4.8	1.59	3	3.1	1.78	6	6.5	2.66
10-14	7	4.0	1.50	2	2.2	1.57	5	5.8	2.60
15-19	13	7.9	2.20	8	9.5	3.34	5	6.3	2.82
20-24	14	7.5	2.00	5	5.2	2.34	9	9.8	3.27
25-29	29	13.4	2.49	19	17.5	4.01	10	9.3	2.94
30-34	82	31.9	3.53	54	42.0	5.71	28	21.8	4.13
35-39	100	39.6	3.96	75	59.6	6.88	25	19.7	3.94
40-44	155	69.8	5.61	108	98.9	9.52	47	41.7	6.08
45-49	218	109.8	7.44	160	163.7	12.94	58	57.6	7.56
50-54	276	176.5	10.62	189	246.8	17.95	87	109.0	11.69
55-59	441	347.7	16.56	319	517.0	28.95	122	187.3	16.96
60-64	698	556.8	21.07	441	735.6	35.03	257	392.9	24.51
65-69	1,195	949.4	27.46	776	1366.3	49.05	419	606.5	29.63
70-74	1,826	1598.4	37.41	1,058	2155.5	66.27	768	1178.8	42.53
75-79	2,401	2632.6	53.73	1,281	3517.7	98.28	1,120	2044.4	61.09
80-84	2,689	4617.2	89.04	1,192	5925.9	171.64	1,497	3926.7	101.49
85+	5,038	10119.1	142.57	1,473	11404.5	297.15	3,565	9668.8	161.94
Total deaths	15,225			7,180			8,045		
Connecticut									
. Crude rate		523.1	-		507.9	-		537.4	-
. AAMR (1940 Std. mil.)		210.3	2.03		279.8	3.57		156.2	2.28
. AAMR (1970 Std. mil.)		321.8	2.73		418.2	4.98		251.8	3.08
. AAMR (2000 Std. mil.)		466.1	3.79		593.3	7.23		379.1	4.28
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									No comparison is available
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	0	0.0	0.00	1	1.0	1.03
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	5	2.3	1.03	5	4.6	2.06	0	0.0	0.00
30-34	10	3.9	1.23	8	6.2	2.20	2	1.6	1.10
35-39	14	5.5	1.48	12	9.5	2.75	2	1.6	1.12
40-44	48	21.6	3.12	34	31.1	5.34	14	12.4	3.32
45-49	106	53.4	5.19	87	89.0	9.54	19	18.9	4.33
50-54	120	76.7	7.00	94	122.8	12.66	26	32.6	6.39
55-59	209	164.8	11.40	169	273.9	21.07	40	61.4	9.71
60-64	344	274.4	14.79	249	415.3	26.32	95	145.2	14.90
65-69	526	417.9	18.22	376	662.0	34.14	150	217.1	17.73
70-74	790	691.5	24.60	469	955.5	44.12	321	492.7	27.50
75-79	1,088	1193.0	36.17	610	1675.1	67.82	478	872.5	39.91
80-84	1,159	1990.1	58.46	496	2465.8	110.72	663	1739.1	67.54
85+	2,336	4692.0	97.08	641	4962.8	196.02	1,695	4597.1	111.66
Total deaths	6,756			3,250			3,506		
Connecticut									
. Crude rate		232.1	-		229.9	-		234.2	-
. AAMR (1940 Std. mil.)		90.9	1.32		128.1	2.42		60.8	1.32
. AAMR (1970 Std. mil.)		141.8	1.80		190.7	3.37		103.9	1.90
. AAMR (2000 Std. mil.)		206.7	2.52		266.8	4.83		162.3	2.77
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		113.8	-		157.4	-		79.2	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.5	0.53	0	0.0	0.00	1	1.1	1.09
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	3	1.5	0.87	2	2.0	1.45	1	1.0	0.99
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	8	6.3	2.23	4	6.5	3.24	4	6.1	3.07
60-64	4	3.2	1.60	2	3.3	2.36	2	3.1	2.16
65-69	16	12.7	3.18	8	14.1	4.98	8	11.6	4.09
70-74	37	32.4	5.32	23	46.9	9.77	14	21.5	5.74
75-79	69	75.7	9.11	30	82.4	15.04	39	71.2	11.40
80-84	76	130.5	14.97	30	149.1	27.23	46	120.7	17.79
85+	233	468.0	30.66	69	534.2	64.31	164	444.8	34.73
Total deaths	447			168			279		
Connecticut									
. Crude rate		15.4	-		11.9	-		18.6	-
. AAMR (1940 Std. mil.)		4.5	0.25		5.1	0.42		4.0	0.32
. AAMR (1970 Std. mil.)		8.3	0.40		9.5	0.74		7.6	0.49
. AAMR (2000 Std. mil.)		13.7	0.65		15.7	1.25		12.6	0.76
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.3	-		7.0	-		5.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	4	1.8	0.90	4	3.7	1.83	0	0.0	0.00
45-49	3	1.5	0.87	2	2.0	1.45	1	1.0	0.99
50-54	3	1.9	1.11	3	3.9	2.26	0	0.0	0.00
55-59	6	4.7	1.93	4	6.5	3.24	2	3.1	2.17
60-64	1	0.8	0.80	0	0.0	0.00	1	1.5	1.53
65-69	10	7.9	2.51	4	7.0	3.52	6	8.7	3.55
70-74	11	9.6	2.90	4	8.1	4.07	7	10.7	4.06
75-79	16	17.5	4.39	8	22.0	7.77	8	14.6	5.16
80-84	24	41.2	8.41	11	54.7	16.49	13	34.1	9.46
85+	41	82.4	12.86	15	116.1	29.99	26	70.5	13.83
Total deaths	120			56			64		
Connecticut									
. Crude rate		4.1	-		4.0	-		4.3	-
. AAMR (1940 Std. mil.)		1.7	0.18		2.2	0.32		1.3	0.20
. AAMR (1970 Std. mil.)		2.5	0.24		3.3	0.44		2.0	0.28
. AAMR (2000 Std. mil.)		3.7	0.34		4.9	0.68		3.0	0.38
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		2.3	-		2.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	1	1.0	0.97	0	0.0	0.00
5-9	1	0.5	0.53	0	0.0	0.00	1	1.1	1.09
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.61	1	1.2	1.18	0	0.0	0.00
20-24	1	0.5	0.53	0	0.0	0.00	1	1.1	1.09
25-29	4	1.8	0.92	1	0.9	0.92	3	2.8	1.61
30-34	6	2.3	0.95	0	0.0	0.00	6	4.7	1.91
35-39	8	3.2	1.12	7	5.6	2.10	1	0.8	0.79
40-44	12	5.4	1.56	8	7.3	2.59	4	3.5	1.77
45-49	14	7.1	1.88	6	6.1	2.51	8	7.9	2.81
50-54	25	16.0	3.20	15	19.6	5.06	10	12.5	3.96
55-59	23	18.1	3.78	8	13.0	4.58	15	23.0	5.95
60-64	55	43.9	5.92	29	48.4	8.98	26	39.7	7.80
65-69	82	65.1	7.19	51	89.8	12.57	31	44.9	8.06
70-74	154	134.8	10.86	79	160.9	18.11	75	115.1	13.29
75-79	255	279.6	17.51	116	318.5	29.58	139	253.7	21.52
80-84	336	576.9	31.47	120	596.6	54.46	216	566.6	38.55
85+	738	1482.3	54.56	195	1509.8	108.12	543	1472.7	63.20
Total deaths	1,716			637			1,079		
Connecticut									
. Crude rate		59.0	-		45.1	-		72.1	-
. AAMR (1940 Std. mil.)		20.0	0.58		22.0	0.94		18.6	0.75
. AAMR (1970 Std. mil.)		33.6	0.84		36.5	1.45		31.6	1.05
. AAMR (2000 Std. mil.)		52.6	1.27		56.3	2.30		50.0	1.54
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		26.6	-		29.0	-		24.4	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Aortic aneurysm (441)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.40	1	0.8	0.79	0	0.0	0.00
40-44	2	0.9	0.64	1	0.9	0.92	1	0.9	0.89
45-49	2	1.0	0.71	2	2.0	1.45	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	3	2.4	1.37	3	4.9	2.81	0	0.0	0.00
60-64	9	7.2	2.39	8	13.3	4.72	1	1.5	1.53
65-69	27	21.5	4.13	24	42.3	8.63	3	4.3	2.51
70-74	41	35.9	5.61	22	44.8	9.56	19	29.2	6.69
75-79	28	30.7	5.80	16	43.9	10.98	12	21.9	6.32
80-84	34	58.4	10.01	18	89.5	21.09	16	42.0	10.49
85+	31	62.3	11.18	17	131.6	31.92	14	38.0	10.15
Total deaths	178			112			66		
Connecticut									
. Crude rate		6.1	-		7.9	-		4.4	-
. AAMR (1940 Std. mil.)		2.7	0.22		4.4	0.44		1.4	0.20
. AAMR (1970 Std. mil.)		4.0	0.30		6.5	0.62		2.2	0.28
. AAMR (2000 Std. mil.)		5.3	0.40		8.7	0.85		3.1	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.4	-		5.6	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	3.0	1.22	1	1.0	0.97	5	5.1	2.30
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.5	0.53	1	1.0	1.05	0	0.0	0.00
25-29	3	1.4	0.80	2	1.8	1.30	1	0.9	0.93
30-34	4	1.6	0.78	4	3.1	1.55	0	0.0	0.00
35-39	4	1.6	0.79	2	1.6	1.12	2	1.6	1.12
40-44	11	5.0	1.49	7	6.4	2.42	4	3.5	1.77
45-49	6	3.0	1.23	3	3.1	1.77	3	3.0	1.72
50-54	9	5.8	1.92	5	6.5	2.92	4	5.0	2.51
55-59	13	10.3	2.84	8	13.0	4.58	5	7.7	3.43
60-64	27	21.5	4.14	17	28.4	6.88	10	15.3	4.83
65-69	42	33.4	5.15	24	42.3	8.63	18	26.1	6.14
70-74	79	69.2	7.78	42	85.6	13.20	37	56.8	9.34
75-79	162	177.6	13.96	94	258.1	26.62	68	124.1	15.05
80-84	200	343.4	24.28	93	462.3	47.94	107	280.7	27.13
85+	575	1154.9	48.16	195	1509.8	108.12	380	1030.6	52.87
Total deaths	1,142			498			644		
Connecticut									
. Crude rate		39.2	-		35.2	-		43.0	-
. AAMR (1940 Std. mil.)		12.3	0.44		16.0	0.78		10.0	0.53
. AAMR (1970 Std. mil.)		21.6	0.66		28.3	1.28		17.8	0.77
. AAMR (2000 Std. mil.)		35.1	1.04		46.3	2.14		29.4	1.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.5	-		17.5	-		10.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	3	1.6	0.92	1	1.0	1.05	2	2.2	1.54
25-29	2	0.9	0.65	1	0.9	0.92	1	0.9	0.93
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.8	0.56	1	0.8	0.79	1	0.8	0.79
40-44	2	0.9	0.64	1	0.9	0.92	1	0.9	0.89
45-49	4	2.0	1.01	3	3.1	1.77	1	1.0	0.99
50-54	9	5.8	1.92	2	2.6	1.85	7	8.8	3.31
55-59	22	17.3	3.70	11	17.8	5.38	11	16.9	5.09
60-64	60	47.9	6.18	30	50.0	9.14	30	45.9	8.37
65-69	98	77.9	7.86	57	100.4	13.29	41	59.4	9.27
70-74	179	156.7	11.71	94	191.5	19.75	85	130.5	14.15
75-79	202	221.5	15.58	96	263.6	26.91	106	193.5	18.79
80-84	233	400.1	26.21	108	536.9	51.66	125	327.9	29.33
85+	229	460.0	30.39	93	720.0	74.66	136	368.9	31.63
Total deaths	1,045			498			547		
Connecticut									
. Crude rate		35.9	-		35.2	-		36.5	-
. AAMR (1940 Std. mil.)		14.7	0.52		18.0	0.86		12.7	0.66
. AAMR (1970 Std. mil.)		22.6	0.72		28.5	1.28		19.3	0.89
. AAMR (2000 Std. mil.)		31.4	0.97		40.8	1.89		26.5	1.15
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		21.4	-		27.8	-		17.2	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	1.1	0.76	0	0.0	0.00	2	2.2	1.54
25-29	2	0.9	0.65	1	0.9	0.92	1	0.9	0.93
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.40	1	0.8	0.79	0	0.0	0.00
40-44	2	0.9	0.64	1	0.9	0.92	1	0.9	0.89
45-49	2	1.0	0.71	1	1.0	1.02	1	1.0	0.99
50-54	3	1.9	1.11	0	0.0	0.00	3	3.8	2.17
55-59	2	1.6	1.12	2	3.2	2.29	0	0.0	0.00
60-64	2	1.6	1.13	2	3.3	2.36	0	0.0	0.00
65-69	4	3.2	1.59	2	3.5	2.49	2	2.9	2.05
70-74	4	3.5	1.75	0	0.0	0.00	4	6.1	3.07
75-79	3	3.3	1.90	0	0.0	0.00	3	5.5	3.16
80-84	5	8.6	3.84	1	5.0	4.97	4	10.5	5.25
85+	10	20.1	6.35	3	23.2	13.41	7	19.0	7.18
Total deaths	42			14			28		
Connecticut									
. Crude rate		1.4	-		1.0	-		1.9	-
. AAMR (1940 Std. mil.)		0.9	0.16		-	-		1.0	0.24
. AAMR (1970 Std. mil.)		1.0	0.17		-	-		1.2	0.25
. AAMR (2000 Std. mil.)		1.3	0.21		-	-		1.5	0.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.5	-		1.3	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	16	8.0	2.00	9	8.8	2.92	7	7.2	2.72
5-9	11	5.8	1.75	9	9.3	3.09	2	2.2	1.53
10-14	12	6.8	1.97	7	7.8	2.94	5	5.8	2.60
15-19	46	28.1	4.14	32	37.8	6.68	14	17.7	4.72
20-24	55	29.4	3.96	44	46.1	6.95	11	12.0	3.61
25-29	68	31.4	3.81	54	49.7	6.76	14	13.0	3.48
30-34	81	31.5	3.50	64	49.8	6.22	17	13.3	3.22
35-39	65	25.7	3.19	52	41.3	5.73	13	10.3	2.84
40-44	43	19.4	2.95	37	33.9	5.57	6	5.3	2.17
45-49	39	19.6	3.15	34	34.8	5.97	5	5.0	2.22
50-54	23	14.7	3.07	12	15.7	4.52	11	13.8	4.16
55-59	27	21.3	4.10	22	35.7	7.60	5	7.7	3.43
60-64	25	19.9	3.99	19	31.7	7.27	6	9.2	3.74
65-69	44	35.0	5.27	21	37.0	8.07	23	33.3	6.94
70-74	42	36.8	5.67	22	44.8	9.56	20	30.7	6.86
75-79	50	54.8	7.75	24	65.9	13.45	26	47.5	9.31
80-84	76	130.5	14.97	38	188.9	30.65	38	99.7	16.17
85+	128	257.1	22.72	46	356.1	52.51	82	222.4	24.56
Total deaths	851			546			305		
Connecticut									
. Crude rate		29.2	-		38.6	-		20.4	-
. AAMR (1940 Std. mil.)		22.7	0.89		33.6	1.54		12.2	0.91
. AAMR (1970 Std. mil.)		23.8	0.88		34.6	1.54		14.0	0.93
. AAMR (2000 Std. mil.)		27.7	0.96		40.3	1.77		16.8	0.99
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		30.2	-		44.2	-		17.0	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Motor vehicle accidents (E810-E825)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	2.5	1.12	3	2.9	1.68	2	2.1	1.46
5-9	7	3.7	1.40	6	6.2	2.52	1	1.1	1.09
10-14	3	1.7	0.98	1	1.1	1.11	2	2.3	1.64
15-19	37	22.6	3.71	27	31.9	6.14	10	12.6	3.99
20-24	40	21.4	3.38	32	33.5	5.92	8	8.7	3.08
25-29	37	17.1	2.81	28	25.7	4.87	9	8.4	2.79
30-34	41	16.0	2.49	27	21.0	4.04	14	10.9	2.92
35-39	20	7.9	1.77	16	12.7	3.18	4	3.2	1.58
40-44	14	6.3	1.69	12	11.0	3.17	2	1.8	1.25
45-49	11	5.5	1.67	9	9.2	3.07	2	2.0	1.40
50-54	12	7.7	2.22	5	6.5	2.92	7	8.8	3.31
55-59	9	7.1	2.37	8	13.0	4.58	1	1.5	1.54
60-64	10	8.0	2.52	8	13.3	4.72	2	3.1	2.16
65-69	21	16.7	3.64	8	14.1	4.98	13	18.8	5.22
70-74	16	14.0	3.50	8	16.3	5.76	8	12.3	4.34
75-79	16	17.5	4.39	8	22.0	7.77	8	14.6	5.16
80-84	19	32.6	7.48	8	39.8	14.06	11	28.9	8.70
85+	5	10.0	4.49	2	15.5	10.95	3	8.1	4.70
Total deaths	323			216			107		
Connecticut									
. Crude rate		11.1	-		15.3	-		7.1	-
. AAMR (1940 Std. mil.)		10.9	0.65		15.5	1.10		6.2	0.69
. AAMR (1970 Std. mil.)		10.6	0.63		14.9	1.06		6.3	0.67
. AAMR (2000 Std. mil.)		10.8	0.61		15.1	1.04		6.6	0.65
. HP 2000 (1940 Std. mil.)		10.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		16.1	-		22.5	-		9.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	0	0.0	0.00	1	1.0	1.03
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.57	1	1.1	1.11	0	0.0	0.00
15-19	2	1.2	0.86	1	1.2	1.18	1	1.3	1.26
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.8	0.55	2	1.6	1.10	0	0.0	0.00
35-39	4	1.6	0.79	3	2.4	1.38	1	0.8	0.79
40-44	1	0.5	0.45	0	0.0	0.00	1	0.9	0.89
45-49	6	3.0	1.23	6	6.1	2.51	0	0.0	0.00
50-54	4	2.6	1.28	2	2.6	1.85	2	2.5	1.77
55-59	4	3.2	1.58	3	4.9	2.81	1	1.5	1.54
60-64	5	4.0	1.78	4	6.7	3.34	1	1.5	1.53
65-69	8	6.4	2.25	5	8.8	3.94	3	4.3	2.51
70-74	9	7.9	2.63	4	8.1	4.07	5	7.7	3.43
75-79	18	19.7	4.65	7	19.2	7.27	11	20.1	6.05
80-84	21	36.1	7.87	10	49.7	15.72	11	28.9	8.70
85+	78	156.7	17.74	26	201.3	39.48	52	141.0	19.56
Total deaths	164			74			90		
Connecticut									
. Crude rate		5.6	-		5.2	-		6.0	-
. AAMR (1940 Std. mil.)		2.2	0.23		3.0	0.39		1.6	0.25
. AAMR (1970 Std. mil.)		3.4	0.28		4.4	0.52		2.7	0.32
. AAMR (2000 Std. mil.)		5.1	0.40		6.7	0.80		4.2	0.45
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.4	-		3.4	-		1.6	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.71	2	1.9	1.38	0	0.0	0.00
5-9	1	0.5	0.53	1	1.0	1.03	0	0.0	0.00
10-14	1	0.6	0.57	1	1.1	1.11	0	0.0	0.00
15-19	2	1.2	0.86	1	1.2	1.18	1	1.3	1.26
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.5	0.46	1	0.9	0.92	0	0.0	0.00
30-34	2	0.8	0.55	2	1.6	1.10	0	0.0	0.00
35-39	5	2.0	0.88	4	3.2	1.59	1	0.8	0.79
40-44	3	1.4	0.78	2	1.8	1.30	1	0.9	0.89
45-49	3	1.5	0.87	3	3.1	1.77	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	1	0.8	0.79	1	1.6	1.62	0	0.0	0.00
60-64	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	1	1.1	1.10	1	2.7	2.75	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	22			19			3		
Connecticut									
. Crude rate		0.8	-		1.3	-		0.2	-
. AAMR (1940 Std. mil.)		0.8	0.17		1.3	0.31		-	-
. AAMR (1970 Std. mil.)		0.7	0.16		1.2	0.30		-	-
. AAMR (2000 Std. mil.)		0.8	0.17		1.3	0.31		-	-
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.6	-		2.8	-		0.6	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.53	0	0.0	0.00	1	1.1	1.09
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.5	0.53	0	0.0	0.00	1	1.1	1.09
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	2	0.9	0.64	2	1.8	1.30	0	0.0	0.00
45-49	2	1.0	0.71	2	2.0	1.45	0	0.0	0.00
50-54	1	0.6	0.64	0	0.0	0.00	1	1.3	1.25
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
65-69	1	0.8	0.79	1	1.8	1.76	0	0.0	0.00
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	4	6.9	3.43	2	9.9	7.03	2	5.2	3.71
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	13			8			5		
Connecticut									
. Crude rate		0.4	-		0.6	-		0.3	-
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. AAMR (1970 Std. mil.)		-	-		-	-		-	-
. AAMR (2000 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.0	-		1.7	-		1.0	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	12	7.3	2.11	9	10.6	3.54	3	3.8	2.18
20-24	22	11.7	2.50	20	20.9	4.68	2	2.2	1.54
25-29	25	11.6	2.31	20	18.4	4.11	5	4.6	2.08
30-34	30	11.7	2.13	23	17.9	3.73	7	5.5	2.06
35-39	33	13.1	2.27	25	19.9	3.97	8	6.3	2.23
40-44	30	13.5	2.47	25	22.9	4.58	5	4.4	1.98
45-49	18	9.1	2.14	15	15.3	3.96	3	3.0	1.72
50-54	22	14.1	3.00	19	24.8	5.69	3	3.8	2.17
55-59	22	17.3	3.70	14	22.7	6.06	8	12.3	4.34
60-64	11	8.8	2.65	9	15.0	5.00	2	3.1	2.16
65-69	13	10.3	2.86	8	14.1	4.98	5	7.2	3.24
70-74	17	14.9	3.61	15	30.6	7.89	2	3.1	2.17
75-79	10	11.0	3.47	6	16.5	6.73	4	7.3	3.65
80-84	7	12.0	4.54	5	24.9	11.12	2	5.2	3.71
85+	4	8.0	4.02	3	23.2	13.41	1	2.7	2.71
Total deaths	276			216			60		
Connecticut									
. Crude rate		9.5	-		15.3	-		4.0	-
. AAMR (1940 Std. mil.)		8.7	0.55		14.0	0.99		3.6	0.49
. AAMR (1970 Std. mil.)		8.3	0.52		13.4	0.95		3.4	0.47
. AAMR (2000 Std. mil.)		9.2	0.56		15.1	1.04		3.8	0.50
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.4	-		18.5	-		4.2	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	1	1.0	0.97	0	0.0	0.00
5-9	2	1.1	0.75	1	1.0	1.03	1	1.1	1.09
10-14	3	1.7	0.98	3	3.3	1.93	0	0.0	0.00
15-19	15	9.1	2.36	13	15.4	4.26	2	2.5	1.78
20-24	22	11.7	2.50	16	16.8	4.19	6	6.5	2.67
25-29	16	7.4	1.85	9	8.3	2.76	7	6.5	2.46
30-34	6	2.3	0.95	6	4.7	1.90	0	0.0	0.00
35-39	9	3.6	1.19	7	5.6	2.10	2	1.6	1.12
40-44	5	2.3	1.01	3	2.7	1.59	2	1.8	1.25
45-49	12	6.0	1.75	9	9.2	3.07	3	3.0	1.72
50-54	4	2.6	1.28	3	3.9	2.26	1	1.3	1.25
55-59	4	3.2	1.58	3	4.9	2.81	1	1.5	1.54
60-64	4	3.2	1.60	3	5.0	2.89	1	1.5	1.53
65-69	1	0.8	0.79	0	0.0	0.00	1	1.4	1.45
70-74	3	2.6	1.52	1	2.0	2.04	2	3.1	2.17
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	2	3.4	2.43	0	0.0	0.00	2	5.2	3.71
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	109			78			31		
Connecticut									
. Crude rate		3.7	-		5.5	-		2.1	-
. AAMR (1940 Std. mil.)		4.3	0.42		6.2	0.73		2.2	0.42
. AAMR (1970 Std. mil.)		4.0	0.40		5.9	0.69		2.1	0.39
. AAMR (2000 Std. mil.)		3.8	0.37		5.6	0.63		2.1	0.37
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.7	-		16.9	-		4.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide (E960-E969)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	1	1.0	0.97	0	0.0	0.00
5-9	2	1.1	0.75	1	1.0	1.03	1	1.1	1.09
10-14	3	1.7	0.98	3	3.3	1.93	0	0.0	0.00
15-19	15	9.1	2.36	13	15.4	4.26	2	2.5	1.78
20-24	22	11.7	2.50	16	16.8	4.19	6	6.5	2.67
25-29	16	7.4	1.85	9	8.3	2.76	7	6.5	2.46
30-34	6	2.3	0.95	6	4.7	1.90	0	0.0	0.00
35-39	8	3.2	1.12	6	4.8	1.95	2	1.6	1.12
40-44	4	1.8	0.90	2	1.8	1.30	2	1.8	1.25
45-49	12	6.0	1.75	9	9.2	3.07	3	3.0	1.72
50-54	4	2.6	1.28	3	3.9	2.26	1	1.3	1.25
55-59	4	3.2	1.58	3	4.9	2.81	1	1.5	1.54
60-64	4	3.2	1.60	3	5.0	2.89	1	1.5	1.53
65-69	1	0.8	0.79	0	0.0	0.00	1	1.4	1.45
70-74	3	2.6	1.52	1	2.0	2.04	2	3.1	2.17
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	2	3.4	2.43	0	0.0	0.00	2	5.2	3.71
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	107			76			31		
Connecticut									
. Crude rate		3.7	-		5.4	-		2.1	-
. AAMR (1940 Std. mil.)		4.2	0.42		6.1	0.72		2.2	0.42
. AAMR (1970 Std. mil.)		4.0	0.40		5.8	0.69		2.1	0.39
. AAMR (2000 Std. mil.)		3.8	0.37		5.4	0.63		2.1	0.37
. HP 2000 (1940 Std. mil.)		5.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.6	-		16.6	-		4.5	-
. HP 2000 (1940 Std. mil.)		7.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	0	0.0	0.00	1	1.0	1.03
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.57	1	1.1	1.11	0	0.0	0.00
15-19	2	1.2	0.86	1	1.2	1.18	1	1.3	1.26
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	3	1.2	0.67	3	2.3	1.35	0	0.0	0.00
35-39	4	1.6	0.79	3	2.4	1.38	1	0.8	0.79
40-44	1	0.5	0.45	0	0.0	0.00	1	0.9	0.89
45-49	6	3.0	1.23	6	6.1	2.51	0	0.0	0.00
50-54	4	2.6	1.28	3	3.9	2.26	1	1.3	1.25
55-59	4	3.2	1.58	3	4.9	2.81	1	1.5	1.54
60-64	5	4.0	1.78	4	6.7	3.34	1	1.5	1.53
65-69	7	5.6	2.10	5	8.8	3.94	2	2.9	2.05
70-74	8	7.0	2.48	5	10.2	4.56	3	4.6	2.66
75-79	13	14.3	3.95	7	19.2	7.27	6	11.0	4.47
80-84	16	27.5	6.87	7	34.8	13.15	9	23.6	7.87
85+	42	84.4	13.02	19	147.1	33.75	23	62.4	13.01
Total deaths	117			67			50		
Connecticut									
. Crude rate		4.0	-		4.7	-		3.3	-
. AAMR (1940 Std. mil.)		1.9	0.22		2.9	0.39		1.1	0.23
. AAMR (1970 Std. mil.)		2.6	0.26		4.0	0.50		1.7	0.27
. AAMR (2000 Std. mil.)		3.7	0.34		5.8	0.73		2.4	0.35
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.3	-		3.4	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	2	1.1	0.75	1	1.0	1.03	1	1.1	1.09
10-14	3	1.7	0.98	3	3.3	1.93	0	0.0	0.00
15-19	19	11.6	2.66	17	20.1	4.87	2	2.5	1.78
20-24	32	17.1	3.02	27	28.3	5.44	5	5.4	2.44
25-29	24	11.1	2.26	15	13.8	3.56	9	8.4	2.79
30-34	18	7.0	1.65	14	10.9	2.91	4	3.1	1.56
35-39	27	10.7	2.06	22	17.5	3.73	5	3.9	1.76
40-44	15	6.8	1.74	11	10.1	3.04	4	3.5	1.77
45-49	16	8.1	2.02	14	14.3	3.83	2	2.0	1.40
50-54	16	10.2	2.56	13	17.0	4.71	3	3.8	2.17
55-59	12	9.5	2.73	8	13.0	4.58	4	6.1	3.07
60-64	7	5.6	2.11	7	11.7	4.41	0	0.0	0.00
65-69	6	4.8	1.95	6	10.6	4.31	0	0.0	0.00
70-74	11	9.6	2.90	8	16.3	5.76	3	4.6	2.66
75-79	3	3.3	1.90	3	8.2	4.76	0	0.0	0.00
80-84	3	5.2	2.97	3	14.9	8.61	0	0.0	0.00
85+	2	4.0	2.84	2	15.5	10.95	0	0.0	0.00
Total deaths	216			174			42		
Connecticut									
. Crude rate		7.4	-		12.3	-		2.8	-
. AAMR (1940 Std. mil.)		7.6	0.54		12.4	0.98		3.0	0.47
. AAMR (1970 Std. mil.)		7.2	0.51		11.8	0.94		2.7	0.43
. AAMR (2000 Std. mil.)		7.4	0.51		12.3	0.95		2.8	0.44
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.5	-		26.9	-		4.6	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	5	3.0	1.36	5	5.9	2.64	0	0.0	0.00
20-24	13	6.9	1.93	12	12.6	3.63	1	1.1	1.09
25-29	10	4.6	1.46	6	5.5	2.25	4	3.7	1.86
30-34	12	4.7	1.35	8	6.2	2.20	4	3.1	1.56
35-39	23	9.1	1.90	19	15.1	3.46	4	3.2	1.58
40-44	11	5.0	1.49	8	7.3	2.59	3	2.7	1.53
45-49	10	5.0	1.59	9	9.2	3.07	1	1.0	0.99
50-54	13	8.3	2.31	11	14.4	4.33	2	2.5	1.77
55-59	12	9.5	2.73	8	13.0	4.58	4	6.1	3.07
60-64	6	4.8	1.95	6	10.0	4.09	0	0.0	0.00
65-69	6	4.8	1.95	6	10.6	4.31	0	0.0	0.00
70-74	9	7.9	2.63	7	14.3	5.39	2	3.1	2.17
75-79	2	2.2	1.55	2	5.5	3.88	0	0.0	0.00
80-84	3	5.2	2.97	3	14.9	8.61	0	0.0	0.00
85+	2	4.0	2.84	2	15.5	10.95	0	0.0	0.00
Total deaths	137			112			25		
Connecticut									
. Crude rate		4.7	-		7.9	-		1.7	-
. AAMR (1940 Std. mil.)		4.4	0.39		7.3	0.71		1.6	0.33
. AAMR (1970 Std. mil.)		4.1	0.37		7.1	0.69		1.4	0.29
. AAMR (2000 Std. mil.)		4.6	0.40		7.9	0.76		1.6	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.8	-		12.1	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	2	1.1	0.75	1	1.0	1.03	1	1.1	1.09
10-14	3	1.7	0.98	3	3.3	1.93	0	0.0	0.00
15-19	14	8.5	2.28	12	14.2	4.09	2	2.5	1.78
20-24	18	9.6	2.27	14	14.7	3.92	4	4.4	2.18
25-29	14	6.5	1.73	9	8.3	2.76	5	4.6	2.08
30-34	6	2.3	0.95	6	4.7	1.90	0	0.0	0.00
35-39	3	1.2	0.69	2	1.6	1.12	1	0.8	0.79
40-44	3	1.4	0.78	2	1.8	1.30	1	0.9	0.89
45-49	6	3.0	1.23	5	5.1	2.29	1	1.0	0.99
50-54	3	1.9	1.11	2	2.6	1.85	1	1.3	1.25
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	1	0.8	0.80	1	1.7	1.67	0	0.0	0.00
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	2	1.8	1.24	1	2.0	2.04	1	1.5	1.53
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	75			58			17		
Connecticut									
. Crude rate		2.6	-		4.1	-		1.1	-
. AAMR (1940 Std. mil.)		3.1	0.37		4.9	0.65		1.4	0.35
. AAMR (1970 Std. mil.)		2.9	0.35		4.5	0.62		1.3	0.32
. AAMR (2000 Std. mil.)		2.7	0.31		4.1	0.55		1.2	0.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.6	-		12.8	-		2.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All poisoning (E850-E869, E950-E952, E962, E972, E980-E982)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	1.7	0.98	1	1.1	1.11	2	2.3	1.64
15-19	7	4.3	1.61	3	3.5	2.05	4	5.0	2.52
20-24	17	9.1	2.20	14	14.7	3.92	3	3.3	1.89
25-29	33	15.3	2.66	29	26.7	4.95	4	3.7	1.86
30-34	42	16.4	2.52	36	28.0	4.66	6	4.7	1.91
35-39	36	14.2	2.37	28	22.2	4.20	8	6.3	2.23
40-44	35	15.8	2.66	31	28.4	5.10	4	3.5	1.77
45-49	17	8.6	2.08	14	14.3	3.83	3	3.0	1.72
50-54	6	3.8	1.57	5	6.5	2.92	1	1.3	1.25
55-59	5	3.9	1.76	2	3.2	2.29	3	4.6	2.66
60-64	6	4.8	1.95	4	6.7	3.34	2	3.1	2.16
65-69	5	4.0	1.78	2	3.5	2.49	3	4.3	2.51
70-74	7	6.1	2.32	6	12.2	4.99	1	1.5	1.53
75-79	6	6.6	2.69	1	2.7	2.75	5	9.1	4.08
80-84	6	10.3	4.21	4	19.9	9.94	2	5.2	3.71
85+	3	6.0	3.48	2	15.5	10.95	1	2.7	2.71
Total deaths	234			182			52		
Connecticut									
. Crude rate		8.0	-		12.9	-		3.5	-
. AAMR (1940 Std. mil.)		7.5	0.51		11.8	0.90		3.2	0.49
. AAMR (1970 Std. mil.)		6.6	0.46		10.4	0.79		3.1	0.47
. AAMR (2000 Std. mil.)		7.6	0.50		12.2	0.91		3.3	0.47
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.7	-		8.1	-		3.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is higher
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.5	0.53	0	0.0	0.00	1	1.1	1.09
25-29	3	1.4	0.80	2	1.8	1.30	1	0.9	0.93
30-34	5	1.9	0.87	3	2.3	1.35	2	1.6	1.10
35-39	10	4.0	1.25	6	4.8	1.95	4	3.2	1.58
40-44	20	9.0	2.01	14	12.8	3.43	6	5.3	2.17
45-49	16	8.1	2.02	13	13.3	3.69	3	3.0	1.72
50-54	21	13.4	2.93	19	24.8	5.69	2	2.5	1.77
55-59	15	11.8	3.05	13	21.1	5.84	2	3.1	2.17
60-64	14	11.2	2.98	11	18.3	5.53	3	4.6	2.65
65-69	21	16.7	3.64	17	29.9	7.26	4	5.8	2.90
70-74	11	9.6	2.90	7	14.3	5.39	4	6.1	3.07
75-79	8	8.8	3.10	6	16.5	6.73	2	3.7	2.58
80-84	4	6.9	3.43	4	19.9	9.94	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	149			115			34		
Connecticut									
. Crude rate		5.1	-		8.1	-		2.3	-
. AAMR (1940 Std. mil.)		4.2	0.35		6.8	0.65		1.8	0.33
. AAMR (1970 Std. mil.)		4.3	0.36		7.2	0.68		1.8	0.32
. AAMR (2000 Std. mil.)		5.0	0.41		8.2	0.77		2.1	0.37
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.8	-		10.9	-		3.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1993

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
 U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
 for
 Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	5	3.0	1.36	2	2.4	1.67	3	3.8	2.18
20-24	13	6.9	1.93	10	10.5	3.31	3	3.3	1.89
25-29	30	13.9	2.53	26	23.9	4.69	4	3.7	1.86
30-34	39	15.2	2.43	35	27.2	4.60	4	3.1	1.56
35-39	38	15.0	2.44	28	22.2	4.20	10	7.9	2.49
40-44	33	14.9	2.59	29	26.6	4.93	4	3.5	1.77
45-49	12	6.0	1.75	11	11.3	3.39	1	1.0	0.99
50-54	3	1.9	1.11	3	3.9	2.26	0	0.0	0.00
55-59	2	1.6	1.12	0	0.0	0.00	2	3.1	2.17
60-64	6	4.8	1.95	4	6.7	3.34	2	3.1	2.16
65-69	4	3.2	1.59	1	1.8	1.76	3	4.3	2.51
70-74	6	5.3	2.14	4	8.1	4.07	2	3.1	2.17
75-79	4	4.4	2.19	0	0.0	0.00	4	7.3	3.65
80-84	1	1.7	1.72	0	0.0	0.00	1	2.6	2.62
85+	2	4.0	2.84	1	7.7	7.74	1	2.7	2.71
Total deaths	198			154			44		
Connecticut									
. Crude rate		6.8	-		10.9	-		2.9	-
. AAMR (1940 Std. mil.)		6.3	0.46		10.1	0.82		2.6	0.43
. AAMR (1970 Std. mil.)		5.4	0.40		8.5	0.70		2.4	0.40
. AAMR (2000 Std. mil.)		6.4	0.46		10.1	0.82		2.8	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.8	-		6.7	-		2.9	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
 . The CT HP 2000 target: No comparison is available
 . The U.S. AAMR (1940 Std. mil.): CT rate is higher
 . THE U.S. HP 2000 target: CT rate is higher