

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Septicemia (038)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.5	0.31	2	0.7	0.50	1	0.4	0.37
5-9	1	0.2	0.17	0	0.0	0.00	1	0.3	0.35
10-14	1	0.2	0.18	0	0.0	0.00	1	0.4	0.37
15-19	1	0.2	0.20	0	0.0	0.00	1	0.4	0.41
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	3	0.5	0.31	3	1.1	0.62	0	0.0	0.00
30-34	2	0.3	0.20	1	0.3	0.29	1	0.3	0.29
35-39	6	0.8	0.32	1	0.3	0.26	5	1.3	0.58
40-44	14	1.9	0.52	9	2.5	0.84	5	1.4	0.61
45-49	14	2.2	0.60	8	2.6	0.92	6	1.9	0.78
50-54	22	4.1	0.88	15	5.7	1.47	7	2.6	0.97
55-59	23	5.7	1.19	12	6.1	1.76	11	5.3	1.61
60-64	35	10.4	1.77	22	13.6	2.90	13	7.5	2.08
65-69	70	20.1	2.40	38	23.6	3.83	32	17.0	3.01
70-74	96	29.2	2.98	57	40.0	5.30	39	20.9	3.34
75-79	154	53.5	4.31	79	67.3	7.57	75	44.0	5.08
80-84	230	117.1	7.72	98	135.6	13.70	132	106.3	9.25
85+	370	216.7	11.26	116	248.8	23.10	254	204.6	12.84
Total deaths	1,045			461			584		
Connecticut									
. Crude rate		12.1	-		10.9	-		13.1	-
. AAMR (1940 Std. mil.)		4.4	0.17		5.3	0.28		3.7	0.21
. AAMR (1970 Std. mil.)		6.8	0.22		8.2	0.39		5.8	0.27
. AAMR (2000 Std. mil.)		10.0	0.31		11.9	0.56		8.7	0.37
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.1	-		4.6	-		3.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Oral cavity or pharynx (140.0-149.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.1	0.14	1	0.3	0.29	0	0.0	0.00
35-39	5	0.6	0.29	5	1.3	0.58	0	0.0	0.00
40-44	4	0.6	0.28	2	0.6	0.40	2	0.5	0.39
45-49	5	0.8	0.36	4	1.3	0.65	1	0.3	0.32
50-54	11	2.1	0.62	10	3.8	1.20	1	0.4	0.37
55-59	25	6.2	1.24	18	9.2	2.16	7	3.4	1.28
60-64	31	9.3	1.66	19	11.8	2.70	12	6.9	2.00
65-69	33	9.5	1.65	24	14.9	3.04	9	4.8	1.59
70-74	37	11.2	1.85	18	12.6	2.98	19	10.2	2.33
75-79	38	13.2	2.14	23	19.6	4.09	15	8.8	2.27
80-84	27	13.7	2.65	14	19.4	5.18	13	10.5	2.90
85+	30	17.6	3.21	11	23.6	7.11	19	15.3	3.51
Total deaths	247			149			98		
Connecticut									
. Crude rate		2.9	-		3.5	-		2.2	-
. AAMR (1940 Std. mil.)		1.6	0.12		2.4	0.21		1.0	0.12
. AAMR (1970 Std. mil.)		2.1	0.14		3.0	0.25		1.4	0.15
. AAMR (2000 Std. mil.)		2.5	0.16		3.6	0.29		1.7	0.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.2	-		1.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.22	1	0.4	0.42	0	0.0	0.00
25-29	2	0.4	0.25	1	0.4	0.36	1	0.4	0.36
30-34	6	0.9	0.35	2	0.6	0.40	4	1.1	0.57
35-39	12	1.5	0.45	8	2.1	0.73	4	1.0	0.52
40-44	32	4.4	0.78	20	5.6	1.25	12	3.3	0.94
45-49	32	5.1	0.91	17	5.5	1.34	15	4.7	1.23
50-54	64	11.9	1.49	40	15.2	2.41	24	8.8	1.79
55-59	89	22.1	2.34	50	25.4	3.60	39	18.9	3.02
60-64	126	37.6	3.35	69	42.7	5.14	57	32.9	4.36
65-69	232	66.5	4.36	130	80.8	7.09	102	54.2	5.37
70-74	312	94.8	5.36	166	116.6	9.05	146	78.1	6.47
75-79	367	127.6	6.66	181	154.2	11.46	186	109.2	8.01
80-84	365	185.8	9.73	167	231.2	17.89	198	159.4	11.33
85+	456	267.0	12.50	147	315.3	26.01	309	248.9	14.16
Total deaths	2,096			999			1,097		
Connecticut									
. Crude rate		24.2	-		23.7	-		24.7	-
. AAMR (1940 Std. mil.)		10.9	0.28		13.1	0.45		9.2	0.35
. AAMR (1970 Std. mil.)		15.4	0.36		18.5	0.60		13.1	0.44
. AAMR (2000 Std. mil.)		20.4	0.45		24.6	0.79		17.5	0.54
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.3	-		14.8	-		10.3	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Rectal cancer (154)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.2	0.18	0	0.0	0.00	1	0.4	0.36
30-34	1	0.1	0.14	0	0.0	0.00	1	0.3	0.29
35-39	2	0.3	0.18	2	0.5	0.37	0	0.0	0.00
40-44	6	0.8	0.34	4	1.1	0.56	2	0.5	0.39
45-49	8	1.3	0.45	7	2.3	0.86	1	0.3	0.32
50-54	14	2.6	0.70	11	4.2	1.26	3	1.1	0.63
55-59	19	4.7	1.08	11	5.6	1.69	8	3.9	1.37
60-64	21	6.3	1.37	10	6.2	1.96	11	6.3	1.91
65-69	32	9.2	1.62	13	8.1	2.24	19	10.1	2.32
70-74	66	20.0	2.47	44	30.9	4.66	22	11.8	2.51
75-79	59	20.5	2.67	29	24.7	4.59	30	17.6	3.22
80-84	52	26.5	3.67	21	29.1	6.34	31	25.0	4.48
85+	64	37.5	4.68	22	47.2	10.06	42	33.8	5.22
Total deaths	345			174			171		
Connecticut									
. Crude rate		4.0	-		4.1	-		3.8	-
. AAMR (1940 Std. mil.)		1.9	0.12		2.4	0.19		1.5	0.15
. AAMR (1970 Std. mil.)		2.6	0.15		3.2	0.25		2.1	0.18
. AAMR (2000 Std. mil.)		3.4	0.18		4.2	0.33		2.8	0.22
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.0	-		2.5	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.2	0.18	0	0.0	0.00	1	0.4	0.36
30-34	2	0.3	0.20	2	0.6	0.40	0	0.0	0.00
35-39	6	0.8	0.32	4	1.0	0.52	2	0.5	0.36
40-44	11	1.5	0.46	8	2.2	0.79	3	0.8	0.47
45-49	34	5.4	0.93	17	5.5	1.34	17	5.4	1.30
50-54	47	8.8	1.28	24	9.1	1.87	23	8.4	1.76
55-59	62	15.4	1.95	36	18.3	3.05	26	12.6	2.47
60-64	77	23.0	2.62	48	29.7	4.29	29	16.7	3.11
65-69	148	42.4	3.49	89	55.3	5.86	59	31.4	4.08
70-74	199	60.4	4.28	91	63.9	6.70	108	57.8	5.56
75-79	211	73.3	5.05	88	75.0	7.99	123	72.2	6.51
80-84	185	94.2	6.92	85	117.7	12.76	100	80.5	8.05
85+	169	99.0	7.61	47	100.8	14.70	122	98.3	8.90
Total deaths	1,152			539			613		
Connecticut									
. Crude rate		13.3	-		12.8	-		13.8	-
. AAMR (1940 Std. mil.)		6.6	0.22		7.7	0.35		5.7	0.28
. AAMR (1970 Std. mil.)		8.9	0.28		10.3	0.45		7.8	0.34
. AAMR (2000 Std. mil.)		11.4	0.34		12.9	0.56		10.1	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.2	-		7.2	-		5.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung and other respiratory cancer (160-165)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.2	0.18	1	0.4	0.36	0	0.0	0.00
30-34	6	0.9	0.35	4	1.1	0.57	2	0.6	0.41
35-39	22	2.8	0.60	13	3.4	0.93	9	2.3	0.77
40-44	59	8.2	1.06	29	8.1	1.51	30	8.2	1.49
45-49	126	20.2	1.80	70	22.7	2.72	56	17.7	2.37
50-54	258	48.2	3.00	134	51.0	4.41	124	45.4	4.08
55-59	373	92.5	4.79	221	112.4	7.56	152	73.6	5.97
60-64	559	166.9	7.06	310	191.8	10.89	249	143.6	9.10
65-69	803	230.0	8.12	465	289.0	13.40	338	179.6	9.77
70-74	982	298.2	9.52	576	404.4	16.85	406	217.3	10.78
75-79	977	339.6	10.86	524	446.4	19.50	453	266.0	12.50
80-84	767	390.4	14.10	406	562.0	27.89	361	290.7	15.30
85+	485	284.0	12.90	245	525.5	33.57	240	193.3	12.48
Total deaths	5,418			2,998			2,420		
Connecticut									
. Crude rate		62.6	-		71.1	-		54.5	-
. AAMR (1940 Std. mil.)		34.2	0.52		43.1	0.84		27.5	0.65
. AAMR (1970 Std. mil.)		45.0	0.64		58.0	1.08		35.6	0.78
. AAMR (2000 Std. mil.)		54.3	0.74		71.7	1.32		42.6	0.88
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		39.2	-		54.1	-		27.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.2	0.18	1	0.4	0.36	0	0.0	0.00
30-34	6	0.9	0.35	4	1.1	0.57	2	0.6	0.41
35-39	21	2.7	0.59	12	3.1	0.89	9	2.3	0.77
40-44	57	7.9	1.04	28	7.9	1.48	29	7.9	1.47
45-49	124	19.9	1.78	69	22.4	2.70	55	17.4	2.35
50-54	245	45.7	2.92	123	46.8	4.22	122	44.7	4.04
55-59	359	89.1	4.70	210	106.8	7.37	149	72.2	5.91
60-64	542	161.8	6.95	299	185.0	10.70	243	140.2	8.99
65-69	780	223.5	8.00	448	278.4	13.15	332	176.5	9.68
70-74	960	291.6	9.41	559	392.5	16.60	401	214.6	10.72
75-79	953	331.2	10.73	510	434.5	19.24	443	260.1	12.36
80-84	752	382.8	13.96	394	545.4	27.48	358	288.2	15.23
85+	468	274.0	12.67	233	499.8	32.74	235	189.3	12.35
Total deaths	5,268			2,890			2,378		
Connecticut									
. Crude rate		60.8	-		68.6	-		53.5	-
. AAMR (1940 Std. mil.)		33.2	0.51		41.4	0.83		27.0	0.64
. AAMR (1970 Std. mil.)		43.7	0.63		55.8	1.06		35.0	0.77
. AAMR (2000 Std. mil.)		52.8	0.73		69.1	1.30		41.8	0.87
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		37.8	-		51.8	-		26.8	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.2	0.22	1	0.4	0.42	0	0.0	0.00
25-29	2	0.4	0.25	1	0.4	0.36	1	0.4	0.36
30-34	4	0.6	0.29	3	0.9	0.49	1	0.3	0.29
35-39	10	1.3	0.41	4	1.0	0.52	6	1.5	0.63
40-44	13	1.8	0.50	6	1.7	0.69	7	1.9	0.72
45-49	14	2.2	0.60	11	3.6	1.08	3	0.9	0.55
50-54	19	3.5	0.81	12	4.6	1.32	7	2.6	0.97
55-59	22	5.5	1.16	10	5.1	1.61	12	5.8	1.68
60-64	19	5.7	1.30	13	8.0	2.23	6	3.5	1.41
65-69	31	8.9	1.60	24	14.9	3.04	7	3.7	1.41
70-74	36	10.9	1.82	22	15.4	3.29	14	7.5	2.00
75-79	41	14.3	2.23	23	19.6	4.09	18	10.6	2.49
80-84	33	16.8	2.92	15	20.8	5.36	18	14.5	3.42
85+	35	20.5	3.46	15	32.2	8.31	20	16.1	3.60
Total deaths	280			160			120		
Connecticut									
. Crude rate		3.2	-		3.8	-		2.7	-
. AAMR (1940 Std. mil.)		1.9	0.13		2.5	0.21		1.4	0.15
. AAMR (1970 Std. mil.)		2.3	0.14		3.1	0.25		1.7	0.17
. AAMR (2000 Std. mil.)		2.9	0.17		3.8	0.31		2.1	0.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.7	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	1	0.2	0.18	-	-	-	1	0.4	0.36
30-34	7	1.0	0.38	-	-	-	7	2.0	0.76
35-39	26	3.4	0.66	-	-	-	26	6.7	1.31
40-44	59	8.2	1.06	-	-	-	59	16.1	2.09
45-49	103	16.5	1.63	-	-	-	103	32.6	3.21
50-54	145	27.1	2.25	-	-	-	145	53.1	4.41
55-59	99	24.6	2.47	-	-	-	99	47.9	4.82
60-64	141	42.1	3.54	-	-	-	141	81.3	6.85
65-69	174	49.8	3.78	-	-	-	174	92.5	7.01
70-74	227	68.9	4.58	-	-	-	227	121.5	8.06
75-79	211	73.3	5.05	-	-	-	211	123.9	8.53
80-84	190	96.7	7.02	-	-	-	190	153.0	11.10
85+	270	158.1	9.62	-	-	-	270	217.5	13.23
Total deaths	1,653			-			1,653		
Connecticut									
. Crude rate		-	-		-	-		37.2	-
. AAMR (1940 Std. mil.)		-	-		-	-		19.7	0.57
. AAMR (1970 Std. mil.)		-	-		-	-		24.2	0.64
. AAMR (2000 Std. mil.)		-	-		-	-		29.7	0.75
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		20.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is lower
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	1	0.2	0.22	-	-	-	1	0.4	0.44
25-29	1	0.2	0.18	-	-	-	1	0.4	0.36
30-34	4	0.6	0.29	-	-	-	4	1.1	0.57
35-39	8	1.0	0.36	-	-	-	8	2.1	0.73
40-44	4	0.6	0.28	-	-	-	4	1.1	0.54
45-49	11	1.8	0.53	-	-	-	11	3.5	1.05
50-54	11	2.1	0.62	-	-	-	11	4.0	1.21
55-59	7	1.7	0.66	-	-	-	7	3.4	1.28
60-64	7	2.1	0.79	-	-	-	7	4.0	1.53
65-69	8	2.3	0.81	-	-	-	8	4.3	1.50
70-74	8	2.4	0.86	-	-	-	8	4.3	1.51
75-79	9	3.1	1.04	-	-	-	9	5.3	1.76
80-84	9	4.6	1.53	-	-	-	9	7.2	2.42
85+	13	7.6	2.11	-	-	-	13	10.5	2.90
Total deaths	101			-			101		
Connecticut									
. Crude rate		-	-		-	-		2.3	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.5	0.16
. AAMR (1970 Std. mil.)		-	-		-	-		1.6	0.17
. AAMR (2000 Std. mil.)		-	-		-	-		1.9	0.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is higher
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	1	0.1	0.14	-	-	-	1	0.3	0.27
45-49	4	0.6	0.32	-	-	-	4	1.3	0.63
50-54	3	0.6	0.32	-	-	-	3	1.1	0.63
55-59	6	1.5	0.61	-	-	-	6	2.9	1.19
60-64	9	2.7	0.90	-	-	-	9	5.2	1.73
65-69	14	4.0	1.07	-	-	-	14	7.4	1.99
70-74	16	4.9	1.21	-	-	-	16	8.6	2.14
75-79	26	9.0	1.77	-	-	-	26	15.3	2.99
80-84	24	12.2	2.49	-	-	-	24	19.3	3.94
85+	23	13.5	2.81	-	-	-	23	18.5	3.86
Total deaths	126			-			126		
Connecticut									
. Crude rate		-	-		-	-		2.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	0.13
. AAMR (1970 Std. mil.)		-	-		-	-		1.6	0.16
. AAMR (2000 Std. mil.)		-	-		-	-		2.1	0.19
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	1	0.2	0.18	-	-	-	1	0.4	0.36
30-34	3	0.4	0.25	-	-	-	3	0.9	0.50
35-39	3	0.4	0.22	-	-	-	3	0.8	0.45
40-44	9	1.2	0.41	-	-	-	9	2.5	0.82
45-49	17	2.7	0.66	-	-	-	17	5.4	1.30
50-54	44	8.2	1.24	-	-	-	44	16.1	2.43
55-59	31	7.7	1.38	-	-	-	31	15.0	2.70
60-64	41	12.2	1.91	-	-	-	41	23.7	3.69
65-69	63	18.0	2.27	-	-	-	63	33.5	4.22
70-74	76	23.1	2.65	-	-	-	76	40.7	4.67
75-79	77	26.8	3.05	-	-	-	77	45.2	5.15
80-84	55	28.0	3.78	-	-	-	55	44.3	5.97
85+	85	49.8	5.40	-	-	-	85	68.5	7.43
Total deaths	505			-			505		
Connecticut									
. Crude rate		-	-		-	-		11.4	-
. AAMR (1940 Std. mil.)		-	-		-	-		5.8	0.30
. AAMR (1970 Std. mil.)		-	-		-	-		7.3	0.35
. AAMR (2000 Std. mil.)		-	-		-	-		8.9	0.41
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Brain & other CNS cancer (191-192)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.5	0.31	1	0.4	0.35	2	0.7	0.53
5-9	6	1.0	0.41	5	1.6	0.74	1	0.3	0.35
10-14	3	0.5	0.31	2	0.7	0.50	1	0.4	0.37
15-19	1	0.2	0.20	0	0.0	0.00	1	0.4	0.41
20-24	3	0.6	0.37	3	1.3	0.73	0	0.0	0.00
25-29	3	0.5	0.31	3	1.1	0.62	0	0.0	0.00
30-34	6	0.9	0.35	4	1.1	0.57	2	0.6	0.41
35-39	14	1.8	0.48	9	2.3	0.77	5	1.3	0.58
40-44	11	1.5	0.46	7	2.0	0.74	4	1.1	0.54
45-49	31	5.0	0.89	18	5.8	1.38	13	4.1	1.14
50-54	38	7.1	1.15	20	7.6	1.70	18	6.6	1.55
55-59	41	10.2	1.59	21	10.7	2.33	20	9.7	2.17
60-64	33	9.9	1.71	24	14.8	3.03	9	5.2	1.73
65-69	53	15.2	2.09	32	19.9	3.52	21	11.2	2.44
70-74	61	18.5	2.37	23	16.1	3.37	38	20.3	3.30
75-79	46	16.0	2.36	25	21.3	4.26	21	12.3	2.69
80-84	46	23.4	3.45	23	31.8	6.64	23	18.5	3.86
85+	35	20.5	3.46	15	32.2	8.31	20	16.1	3.60
Total deaths	434			235			199		
Connecticut									
. Crude rate		5.0	-		5.6	-		4.5	-
. AAMR (1940 Std. mil.)		3.3	0.17		4.0	0.28		2.6	0.21
. AAMR (1970 Std. mil.)		3.9	0.20		4.8	0.32		3.1	0.24
. AAMR (2000 Std. mil.)		4.5	0.22		5.6	0.37		3.7	0.27
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.4	-		4.4	-		2.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	0.7	0.36	1	0.4	0.35	3	1.1	0.65
5-9	6	1.0	0.41	1	0.3	0.33	5	1.7	0.78
10-14	3	0.5	0.31	3	1.1	0.61	0	0.0	0.00
15-19	4	0.8	0.39	2	0.8	0.54	2	0.8	0.57
20-24	8	1.7	0.61	6	2.5	1.03	2	0.9	0.63
25-29	4	0.7	0.36	4	1.4	0.71	0	0.0	0.00
30-34	13	1.9	0.52	7	2.0	0.76	6	1.7	0.70
35-39	14	1.8	0.48	7	1.8	0.68	7	1.8	0.68
40-44	19	2.6	0.60	9	2.5	0.84	10	2.7	0.86
45-49	14	2.2	0.60	8	2.6	0.92	6	1.9	0.78
50-54	33	6.2	1.07	17	6.5	1.57	16	5.9	1.46
55-59	29	7.2	1.34	18	9.2	2.16	11	5.3	1.61
60-64	48	14.3	2.07	23	14.2	2.97	25	14.4	2.88
65-69	70	20.1	2.40	49	30.5	4.35	21	11.2	2.44
70-74	110	33.4	3.19	66	46.3	5.70	44	23.5	3.55
75-79	148	51.4	4.23	84	71.6	7.81	64	37.6	4.70
80-84	108	55.0	5.29	54	74.7	10.17	54	43.5	5.92
85+	116	67.9	6.31	50	107.2	15.17	66	53.2	6.54
Total deaths	751			409			342		
Connecticut									
. Crude rate		8.7	-		9.7	-		7.7	-
. AAMR (1940 Std. mil.)		4.6	0.20		5.8	0.32		3.6	0.25
. AAMR (1970 Std. mil.)		5.9	0.23		7.7	0.39		4.7	0.28
. AAMR (2000 Std. mil.)		7.5	0.28		10.0	0.50		5.9	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.8	-		6.2	-		3.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.18	1	0.4	0.35	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	0.4	0.31	1	0.4	0.42	1	0.4	0.44
25-29	3	0.5	0.31	1	0.4	0.36	2	0.7	0.51
30-34	10	1.4	0.45	3	0.9	0.49	7	2.0	0.76
35-39	28	3.6	0.68	19	4.9	1.13	9	2.3	0.77
40-44	61	8.4	1.08	41	11.5	1.80	20	5.4	1.22
45-49	100	16.0	1.60	60	19.5	2.52	40	12.7	2.00
50-54	168	31.4	2.42	102	38.8	3.85	66	24.2	2.97
55-59	284	70.4	4.18	177	90.0	6.76	107	51.8	5.01
60-64	402	120.0	5.99	240	148.5	9.58	162	93.5	7.34
65-69	648	185.6	7.29	368	228.7	11.92	280	148.8	8.89
70-74	940	285.5	9.31	504	353.9	15.76	436	233.3	11.18
75-79	1,287	447.3	12.47	660	562.2	21.89	627	368.1	14.70
80-84	1,319	671.4	18.49	636	880.3	34.91	683	549.9	21.04
85+	1,606	940.4	23.47	530	1136.8	49.38	1,076	866.7	26.42
Total deaths	6,859			3,343			3,516		
Connecticut									
. Crude rate		79.2	-		79.3	-		79.1	-
. AAMR (1940 Std. mil.)		33.8	0.48		42.2	0.80		27.2	0.58
. AAMR (1970 Std. mil.)		49.1	0.63		61.4	1.08		40.2	0.75
. AAMR (2000 Std. mil.)		66.4	0.81		83.0	1.45		55.1	0.95
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	3	0.4	0.22	2	0.5	0.37	1	0.3	0.26
40-44	3	0.4	0.24	1	0.3	0.28	2	0.5	0.39
45-49	2	0.3	0.23	2	0.6	0.46	0	0.0	0.00
50-54	2	0.4	0.26	2	0.8	0.54	0	0.0	0.00
55-59	3	0.7	0.43	2	1.0	0.72	1	0.5	0.48
60-64	1	0.3	0.30	1	0.6	0.62	0	0.0	0.00
65-69	1	0.3	0.29	1	0.6	0.62	0	0.0	0.00
70-74	6	1.8	0.74	3	2.1	1.22	3	1.6	0.93
75-79	3	1.0	0.60	2	1.7	1.20	1	0.6	0.59
80-84	5	2.5	1.14	3	4.2	2.40	2	1.6	1.14
85+	3	1.8	1.01	0	0.0	0.00	3	2.4	1.40
Total deaths	32			19			13		
Connecticut									
. Crude rate		0.4	-		0.5	-		0.3	-
. AAMR (1940 Std. mil.)		0.2	0.04		0.3	0.07		-	-
. AAMR (1970 Std. mil.)		0.3	0.05		0.4	0.08		-	-
. AAMR (2000 Std. mil.)		0.3	0.06		0.4	0.10		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.4	-		0.5	-		0.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Alzheimer's diseases (331.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.1	0.14	1	0.3	0.28	0	0.0	0.00
45-49	1	0.2	0.16	0	0.0	0.00	1	0.3	0.32
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	4	1.0	0.50	0	0.0	0.00	4	1.9	0.97
60-64	6	1.8	0.73	4	2.5	1.24	2	1.2	0.82
65-69	20	5.7	1.28	14	8.7	2.33	6	3.2	1.30
70-74	43	13.1	1.99	13	9.1	2.53	30	16.1	2.93
75-79	105	36.5	3.56	39	33.2	5.32	66	38.8	4.77
80-84	173	88.1	6.70	57	78.9	10.45	116	93.4	8.67
85+	395	231.3	11.64	92	197.3	20.57	303	244.1	14.02
Total deaths	748			220			528		
Connecticut									
. Crude rate		8.6	-		5.2	-		11.9	-
. AAMR (1940 Std. mil.)		2.1	0.09		1.9	0.14		2.2	0.12
. AAMR (1970 Std. mil.)		4.1	0.16		3.6	0.25		4.3	0.20
. AAMR (2000 Std. mil.)		6.9	0.25		6.1	0.42		7.3	0.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.7	-		2.6	-		2.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Heart disease-related (390-398,402,404-429)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	73	13.3	1.55	46	16.3	2.40	27	10.1	1.94
5-9	21	3.6	0.78	11	3.6	1.09	10	3.5	1.10
10-14	18	3.2	0.76	9	3.2	1.05	9	3.3	1.10
15-19	22	4.3	0.92	16	6.1	1.52	6	2.4	0.99
20-24	41	8.9	1.38	25	10.5	2.10	16	7.1	1.78
25-29	63	11.3	1.42	40	14.2	2.25	23	8.3	1.74
30-34	124	17.7	1.59	79	22.6	2.54	45	12.9	1.92
35-39	256	33.0	2.06	177	45.7	3.44	79	20.3	2.29
40-44	410	56.6	2.80	298	83.6	4.84	112	30.5	2.88
45-49	639	102.4	4.05	434	141.0	6.77	205	64.9	4.53
50-54	945	176.4	5.74	642	244.4	9.65	303	110.9	6.37
55-59	1,334	330.9	9.06	883	449.0	15.11	451	218.4	10.29
60-64	1,889	563.9	12.97	1,209	748.0	21.51	680	392.3	15.04
65-69	3,159	905.0	16.10	1,975	1227.3	27.62	1,184	629.3	18.29
70-74	4,790	1454.7	21.02	2,731	1917.5	36.69	2,059	1102.0	24.29
75-79	7,055	2452.2	29.20	3,719	3168.1	51.95	3,336	1958.8	33.91
80-84	8,515	4334.6	46.97	4,010	5550.6	87.65	4,505	3627.2	54.04
85+	16,630	9737.9	75.51	5,031	10791.0	152.14	11,599	9342.4	86.75
Total deaths	45,984			21,335			24,649		
Connecticut									
. Crude rate		531.1	-		506.2	-		554.8	-
. AAMR (1940 Std. mil.)		197.2	1.14		255.6	1.95		150.5	1.30
. AAMR (1970 Std. mil.)		303.2	1.51		384.6	2.68		242.2	1.74
. AAMR (2000 Std. mil.)		439.8	2.06		547.0	3.80		363.9	2.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									No comparison is available
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.2	0.18	1	0.4	0.35	0	0.0	0.00
15-19	3	0.6	0.34	3	1.1	0.66	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	10	1.8	0.57	7	2.5	0.94	3	1.1	0.63
30-34	24	3.4	0.70	19	5.4	1.24	5	1.4	0.64
35-39	63	8.1	1.02	52	13.4	1.86	11	2.8	0.85
40-44	170	23.5	1.80	136	38.1	3.27	34	9.3	1.59
45-49	283	45.4	2.70	231	75.0	4.94	52	16.5	2.28
50-54	427	79.7	3.86	339	129.1	7.01	88	32.2	3.43
55-59	619	153.5	6.17	452	229.8	10.81	167	80.9	6.26
60-64	808	241.2	8.49	585	361.9	14.96	223	128.6	8.61
65-69	1,366	391.3	10.59	920	571.7	18.85	446	237.0	11.22
70-74	1,995	605.9	13.56	1,240	870.6	24.72	755	404.1	14.71
75-79	2,944	1023.3	18.86	1,646	1402.2	34.56	1,298	762.1	21.15
80-84	3,601	1833.1	30.55	1,727	2390.5	57.52	1,874	1508.9	34.86
85+	7,444	4358.9	50.52	2,176	4667.3	100.05	5,268	4243.1	58.46
Total deaths	19,758			9,534			10,224		
Connecticut									
. Crude rate		228.2	-		226.2	-		230.1	-
. AAMR (1940 Std. mil.)		82.9	0.73		115.7	1.31		56.0	0.75
. AAMR (1970 Std. mil.)		129.1	0.98		172.8	1.80		94.9	1.05
. AAMR (2000 Std. mil.)		188.7	1.35		243.7	2.53		147.5	1.48
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		104.7	-		143.0	-		73.3	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.18	0	0.0	0.00	1	0.4	0.37
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.1	0.13	1	0.3	0.26	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	6	1.0	0.39	2	0.6	0.46	4	1.3	0.63
50-54	9	1.7	0.56	3	1.1	0.66	6	2.2	0.90
55-59	16	4.0	0.99	9	4.6	1.53	7	3.4	1.28
60-64	19	5.7	1.30	9	5.6	1.86	10	5.8	1.82
65-69	50	14.3	2.03	28	17.4	3.29	22	11.7	2.49
70-74	118	35.8	3.30	52	36.5	5.06	66	35.3	4.35
75-79	182	63.3	4.69	90	76.7	8.08	92	54.0	5.63
80-84	323	164.4	9.15	128	177.2	15.66	195	157.0	11.24
85+	901	527.6	17.58	224	480.5	32.10	677	545.3	20.96
Total deaths	1,626			546			1,080		
Connecticut									
. Crude rate		18.8	-		13.0	-		24.3	-
. AAMR (1940 Std. mil.)		4.8	0.15		5.0	0.23		4.7	0.19
. AAMR (1970 Std. mil.)		9.0	0.24		9.2	0.40		8.8	0.30
. AAMR (2000 Std. mil.)		15.1	0.37		15.0	0.65		14.9	0.46
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.0	-		6.8	-		5.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.18	0	0.0	0.00	1	0.4	0.37
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.1	0.13	1	0.3	0.26	0	0.0	0.00
40-44	4	0.6	0.28	4	1.1	0.56	0	0.0	0.00
45-49	3	0.5	0.28	1	0.3	0.32	2	0.6	0.45
50-54	10	1.9	0.59	5	1.9	0.85	5	1.8	0.82
55-59	8	2.0	0.70	5	2.5	1.14	3	1.5	0.84
60-64	13	3.9	1.08	6	3.7	1.52	7	4.0	1.53
65-69	21	6.0	1.31	12	7.5	2.15	9	4.8	1.59
70-74	39	11.8	1.90	15	10.5	2.72	24	12.8	2.62
75-79	62	21.6	2.74	26	22.1	4.34	36	21.1	3.52
80-84	68	34.6	4.20	25	34.6	6.92	43	34.6	5.28
85+	176	103.1	7.77	39	83.7	13.39	137	110.3	9.43
Total deaths	406			139			267		
Connecticut									
. Crude rate		4.7	-		3.3	-		6.0	-
. AAMR (1940 Std. mil.)		1.6	0.10		1.6	0.15		1.5	0.13
. AAMR (1970 Std. mil.)		2.5	0.13		2.5	0.21		2.5	0.17
. AAMR (2000 Std. mil.)		3.8	0.19		3.6	0.31		3.9	0.24
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		2.5	-		2.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	1.1	0.45	3	1.1	0.61	3	1.1	0.65
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	0.4	0.25	1	0.4	0.35	1	0.4	0.37
15-19	1	0.2	0.20	1	0.4	0.38	0	0.0	0.00
20-24	2	0.4	0.31	1	0.4	0.42	1	0.4	0.44
25-29	4	0.7	0.36	1	0.4	0.36	3	1.1	0.63
30-34	14	2.0	0.54	8	2.3	0.81	6	1.7	0.70
35-39	20	2.6	0.58	15	3.9	1.00	5	1.3	0.58
40-44	41	5.7	0.88	18	5.0	1.19	23	6.3	1.31
45-49	50	8.0	1.13	23	7.5	1.56	27	8.5	1.64
50-54	56	10.5	1.40	22	8.4	1.79	34	12.4	2.13
55-59	107	26.5	2.57	71	36.1	4.28	36	17.4	2.91
60-64	114	34.0	3.19	71	43.9	5.21	43	24.8	3.78
65-69	255	73.1	4.57	132	82.0	7.14	123	65.4	5.89
70-74	462	140.3	6.53	236	165.7	10.79	226	121.0	8.05
75-79	809	281.2	9.89	369	314.3	16.36	440	258.3	12.32
80-84	1,161	591.0	17.35	454	628.4	29.49	707	569.2	21.41
85+	2,359	1381.3	28.44	586	1256.9	51.92	1,773	1428.1	33.92
Total deaths	5,463			2,012			3,451		
Connecticut									
. Crude rate		63.1	-		47.7	-		77.7	-
. AAMR (1940 Std. mil.)		19.8	0.33		21.5	0.53		18.3	0.42
. AAMR (1970 Std. mil.)		33.1	0.47		35.0	0.79		31.3	0.59
. AAMR (2000 Std. mil.)		51.4	0.70		52.9	1.20		49.5	0.86
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		26.5	-		28.5	-		24.6	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									
. The U.S. AAMR (1940 Std. mil.):									
. THE U.S. HP 2000 target:									

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Aortic aneurysm (441)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.2	0.18	1	0.4	0.36	0	0.0	0.00
30-34	2	0.3	0.20	2	0.6	0.40	0	0.0	0.00
35-39	5	0.6	0.29	4	1.0	0.52	1	0.3	0.26
40-44	3	0.4	0.24	3	0.8	0.49	0	0.0	0.00
45-49	4	0.6	0.32	2	0.6	0.46	2	0.6	0.45
50-54	7	1.3	0.49	6	2.3	0.93	1	0.4	0.37
55-59	14	3.5	0.93	13	6.6	1.83	1	0.5	0.48
60-64	25	7.5	1.49	21	13.0	2.84	4	2.3	1.15
65-69	69	19.8	2.38	52	32.3	4.48	17	9.0	2.19
70-74	103	31.3	3.08	70	49.1	5.87	33	17.7	3.07
75-79	136	47.3	4.05	78	66.4	7.52	58	34.1	4.47
80-84	131	66.7	5.83	64	88.6	11.07	67	53.9	6.59
85+	162	94.9	7.45	77	165.2	18.82	85	68.5	7.43
Total deaths	662			393			269		
Connecticut									
. Crude rate		7.6	-		9.3	-		6.1	-
. AAMR (1940 Std. mil.)		3.0	0.14		4.8	0.26		1.7	0.13
. AAMR (1970 Std. mil.)		4.5	0.19		7.1	0.36		2.7	0.18
. AAMR (2000 Std. mil.)		6.3	0.25		9.8	0.50		4.0	0.25
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.2	-		4.9	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	8	1.5	0.51	6	2.1	0.87	2	0.7	0.53
5-9	3	0.5	0.29	1	0.3	0.33	2	0.7	0.49
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	3	0.6	0.34	2	0.8	0.54	1	0.4	0.41
20-24	1	0.2	0.22	1	0.4	0.42	0	0.0	0.00
25-29	6	1.1	0.44	2	0.7	0.50	4	1.4	0.72
30-34	5	0.7	0.32	4	1.1	0.57	1	0.3	0.29
35-39	14	1.8	0.48	9	2.3	0.77	5	1.3	0.58
40-44	20	2.8	0.62	15	4.2	1.09	5	1.4	0.61
45-49	28	4.5	0.85	13	4.2	1.17	15	4.7	1.23
50-54	23	4.3	0.90	17	6.5	1.57	6	2.2	0.90
55-59	47	11.7	1.70	28	14.2	2.69	19	9.2	2.11
60-64	63	18.8	2.37	36	22.3	3.71	27	15.6	3.00
65-69	118	33.8	3.11	74	46.0	5.35	44	23.4	3.53
70-74	212	64.4	4.42	131	92.0	8.04	81	43.4	4.82
75-79	407	141.5	7.01	222	189.1	12.69	185	108.6	7.99
80-84	666	339.0	13.14	341	472.0	25.56	325	261.7	14.52
85+	1,792	1049.3	24.79	622	1334.1	53.49	1,170	942.4	27.55
Total deaths	3,416			1,524			1,892		
Connecticut									
. Crude rate		39.5	-		36.2	-		42.6	-
. AAMR (1940 Std. mil.)		11.1	0.24		14.7	0.42		8.8	0.29
. AAMR (1970 Std. mil.)		19.7	0.36		25.9	0.67		15.9	0.41
. AAMR (2000 Std. mil.)		31.9	0.55		41.9	1.09		26.4	0.62
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.0	-		16.3	-		10.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.18	1	0.4	0.35	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.2	0.18	1	0.4	0.35	0	0.0	0.00
15-19	1	0.2	0.20	0	0.0	0.00	1	0.4	0.41
20-24	3	0.6	0.37	3	1.3	0.73	0	0.0	0.00
25-29	2	0.4	0.25	2	0.7	0.50	0	0.0	0.00
30-34	3	0.4	0.25	1	0.3	0.29	2	0.6	0.41
35-39	9	1.2	0.39	4	1.0	0.52	5	1.3	0.58
40-44	17	2.3	0.57	8	2.2	0.79	9	2.5	0.82
45-49	23	3.7	0.77	11	3.6	1.08	12	3.8	1.10
50-54	47	8.8	1.28	22	8.4	1.79	25	9.2	1.83
55-59	78	19.3	2.19	28	14.2	2.69	50	24.2	3.42
60-64	151	45.1	3.67	68	42.1	5.10	83	47.9	5.26
65-69	338	96.8	5.27	177	110.0	8.27	161	85.6	6.74
70-74	542	164.6	7.07	253	177.6	11.17	289	154.7	9.10
75-79	745	259.0	9.49	348	296.5	15.89	397	233.1	11.70
80-84	761	387.4	14.04	338	467.9	25.45	423	340.6	16.56
85+	878	514.1	17.35	380	815.1	41.81	498	401.1	17.97
Total deaths	3,600			1,645			1,955		
Connecticut									
. Crude rate		41.6	-		39.0	-		44.0	-
. AAMR (1940 Std. mil.)		16.2	0.32		18.3	0.50		15.1	0.42
. AAMR (1970 Std. mil.)		24.7	0.43		29.0	0.73		22.5	0.56
. AAMR (2000 Std. mil.)		34.3	0.57		42.0	1.05		30.7	0.71
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		20.9	-		25.8	-		17.6	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.2	0.18	1	0.4	0.35	0	0.0	0.00
15-19	1	0.2	0.20	0	0.0	0.00	1	0.4	0.41
20-24	3	0.6	0.37	3	1.3	0.73	0	0.0	0.00
25-29	2	0.4	0.25	2	0.7	0.50	0	0.0	0.00
30-34	2	0.3	0.20	1	0.3	0.29	1	0.3	0.29
35-39	8	1.0	0.36	4	1.0	0.52	4	1.0	0.52
40-44	8	1.1	0.39	2	0.6	0.40	6	1.6	0.67
45-49	5	0.8	0.36	0	0.0	0.00	5	1.6	0.71
50-54	16	3.0	0.75	8	3.0	1.08	8	2.9	1.04
55-59	9	2.2	0.74	2	1.0	0.72	7	3.4	1.28
60-64	10	3.0	0.94	3	1.9	1.07	7	4.0	1.53
65-69	5	1.4	0.64	1	0.6	0.62	4	2.1	1.06
70-74	12	3.6	1.05	2	1.4	0.99	10	5.4	1.69
75-79	18	6.3	1.47	3	2.6	1.48	15	8.8	2.27
80-84	17	8.7	2.10	4	5.5	2.77	13	10.5	2.90
85+	22	12.9	2.75	3	6.4	3.72	19	15.3	3.51
Total deaths	139			39			100		
Connecticut									
. Crude rate		1.6	-		0.9	-		2.3	-
. AAMR (1940 Std. mil.)		1.0	0.10		0.7	0.13		1.2	0.14
. AAMR (1970 Std. mil.)		1.2	0.10		0.8	0.13		1.4	0.16
. AAMR (2000 Std. mil.)		1.4	0.12		1.0	0.15		1.8	0.18
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.6	-		1.1	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.18	1	0.4	0.35	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.2	0.20	1	0.4	0.38	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	3	0.5	0.31	1	0.4	0.36	2	0.7	0.51
30-34	13	1.9	0.52	9	2.6	0.86	4	1.1	0.57
35-39	27	3.5	0.67	17	4.4	1.07	10	2.6	0.81
40-44	63	8.7	1.10	53	14.9	2.04	10	2.7	0.86
45-49	85	13.6	1.48	61	19.8	2.54	24	7.6	1.55
50-54	79	14.7	1.66	58	22.1	2.90	21	7.7	1.68
55-59	72	17.9	2.10	51	25.9	3.63	21	10.2	2.22
60-64	77	23.0	2.62	54	33.4	4.55	23	13.3	2.77
65-69	97	27.8	2.82	58	36.0	4.73	39	20.7	3.32
70-74	105	31.9	3.11	60	42.1	5.44	45	24.1	3.59
75-79	95	33.0	3.39	36	30.7	5.11	59	34.6	4.51
80-84	71	36.1	4.29	34	47.1	8.07	37	29.8	4.90
85+	31	18.2	3.26	15	32.2	8.31	16	12.9	3.22
Total deaths	820			509			311		
Connecticut									
. Crude rate		9.5	-		12.1	-		7.0	-
. AAMR (1940 Std. mil.)		6.4	0.24		9.0	0.41		4.0	0.26
. AAMR (1970 Std. mil.)		7.3	0.26		10.1	0.46		4.8	0.29
. AAMR (2000 Std. mil.)		8.6	0.30		11.9	0.53		5.7	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.5	-		10.7	-		4.5	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis, nephrotic syndrome/nephrosis (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.5	0.31	1	0.4	0.35	2	0.7	0.53
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.4	0.28	2	0.8	0.54	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	4	0.7	0.36	3	1.1	0.62	1	0.4	0.36
30-34	1	0.1	0.14	0	0.0	0.00	1	0.3	0.29
35-39	6	0.8	0.32	1	0.3	0.26	5	1.3	0.58
40-44	8	1.1	0.39	5	1.4	0.63	3	0.8	0.47
45-49	13	2.1	0.58	9	2.9	0.97	4	1.3	0.63
50-54	18	3.4	0.79	13	4.9	1.37	5	1.8	0.82
55-59	22	5.5	1.16	8	4.1	1.44	14	6.8	1.81
60-64	31	9.3	1.66	17	10.5	2.55	14	8.1	2.16
65-69	53	15.2	2.09	19	11.8	2.71	34	18.1	3.10
70-74	106	32.2	3.13	59	41.4	5.39	47	25.2	3.67
75-79	149	51.8	4.24	77	65.6	7.48	72	42.3	4.98
80-84	195	99.3	7.11	105	145.3	14.18	90	72.5	7.64
85+	340	199.1	10.80	141	302.4	25.47	199	160.3	11.36
Total deaths	951			460			491		
Connecticut									
. Crude rate		11.0	-		10.9	-		11.1	-
. AAMR (1940 Std. mil.)		4.0	0.16		4.9	0.26		3.4	0.20
. AAMR (1970 Std. mil.)		6.2	0.21		8.0	0.38		5.2	0.26
. AAMR (2000 Std. mil.)		9.1	0.29		12.2	0.58		7.5	0.34
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.4	-		5.3	-		3.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----End stage renal disease (ESRD) (581-588,591,593.8,593.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	0.9	0.41	2	0.7	0.50	3	1.1	0.65
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.4	0.28	2	0.8	0.54	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	5	0.9	0.40	4	1.4	0.71	1	0.4	0.36
30-34	2	0.3	0.20	1	0.3	0.29	1	0.3	0.29
35-39	7	0.9	0.34	1	0.3	0.26	6	1.5	0.63
40-44	10	1.4	0.44	7	2.0	0.74	3	0.8	0.47
45-49	18	2.9	0.68	12	3.9	1.13	6	1.9	0.78
50-54	19	3.5	0.81	14	5.3	1.42	5	1.8	0.82
55-59	28	6.9	1.31	11	5.6	1.69	17	8.2	2.00
60-64	40	11.9	1.89	23	14.2	2.97	17	9.8	2.38
65-69	68	19.5	2.36	29	18.0	3.35	39	20.7	3.32
70-74	129	39.2	3.45	68	47.7	5.79	61	32.6	4.18
75-79	181	62.9	4.68	94	80.1	8.26	87	51.1	5.48
80-84	223	113.5	7.60	123	170.3	15.35	100	80.5	8.05
85+	384	224.9	11.47	165	353.9	27.55	219	176.4	11.92
Total deaths	1,121			556			565		
Connecticut									
. Crude rate		12.9	-		13.2	-		12.7	-
. AAMR (1940 Std. mil.)		4.8	0.18		6.1	0.29		4.0	0.22
. AAMR (1970 Std. mil.)		7.4	0.24		9.7	0.42		6.1	0.29
. AAMR (2000 Std. mil.)		10.7	0.32		14.7	0.63		8.6	0.37
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.5	-		6.6	-		4.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	39	7.1	1.14	24	8.5	1.73	15	5.6	1.45
5-9	24	4.1	0.83	19	6.3	1.44	5	1.7	0.78
10-14	31	5.6	1.00	15	5.3	1.36	16	5.9	1.47
15-19	125	24.5	2.19	81	30.8	3.42	44	17.8	2.69
20-24	152	32.8	2.66	117	49.2	4.54	35	15.6	2.63
25-29	153	27.4	2.22	118	42.0	3.86	35	12.7	2.14
30-34	215	30.8	2.10	168	48.0	3.70	47	13.5	1.97
35-39	222	28.6	1.92	174	44.9	3.41	48	12.4	1.78
40-44	200	27.6	1.95	160	44.9	3.55	40	10.9	1.72
45-49	169	27.1	2.08	124	40.3	3.62	45	14.2	2.12
50-54	108	20.2	1.94	77	29.3	3.34	31	11.4	2.04
55-59	78	19.3	2.19	47	23.9	3.49	31	15.0	2.70
60-64	91	27.2	2.85	57	35.3	4.67	34	19.6	3.36
65-69	111	31.8	3.02	76	47.2	5.42	35	18.6	3.14
70-74	171	51.9	3.97	91	63.9	6.70	80	42.8	4.79
75-79	220	76.5	5.16	128	109.0	9.64	92	54.0	5.63
80-84	284	144.6	8.58	132	182.7	15.90	152	122.4	9.93
85+	443	259.4	12.32	181	388.2	28.86	262	211.0	13.04
Total deaths	2,836			1,789			1,047		
Connecticut									
. Crude rate		32.8	-		42.4	-		23.6	-
. AAMR (1940 Std. mil.)		24.2	0.53		34.9	0.91		14.1	0.57
. AAMR (1970 Std. mil.)		25.6	0.53		36.3	0.90		15.9	0.58
. AAMR (2000 Std. mil.)		30.3	0.58		43.5	1.04		19.1	0.61
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		30.5	-		43.2	-		17.8	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Motor vehicle accidents (E810-E825)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	1.1	0.45	1	0.4	0.35	5	1.9	0.84
5-9	14	2.4	0.63	12	4.0	1.14	2	0.7	0.49
10-14	20	3.6	0.80	10	3.5	1.11	10	3.7	1.16
15-19	98	19.2	1.94	57	21.6	2.87	41	16.6	2.60
20-24	94	20.3	2.09	68	28.6	3.46	26	11.6	2.27
25-29	80	14.3	1.60	60	21.3	2.75	20	7.2	1.62
30-34	77	11.0	1.26	56	16.0	2.14	21	6.0	1.31
35-39	67	8.6	1.06	43	11.1	1.69	24	6.2	1.26
40-44	55	7.6	1.02	38	10.7	1.73	17	4.6	1.12
45-49	60	9.6	1.24	44	14.3	2.15	16	5.1	1.27
50-54	45	8.4	1.25	31	11.8	2.12	14	5.1	1.37
55-59	23	5.7	1.19	12	6.1	1.76	11	5.3	1.61
60-64	39	11.6	1.86	22	13.6	2.90	17	9.8	2.38
65-69	33	9.5	1.65	21	13.1	2.85	12	6.4	1.84
70-74	48	14.6	2.10	24	16.9	3.44	24	12.8	2.62
75-79	48	16.7	2.41	27	23.0	4.43	21	12.3	2.69
80-84	47	23.9	3.49	24	33.2	6.78	23	18.5	3.86
85+	32	18.7	3.31	11	23.6	7.11	21	16.9	3.69
Total deaths	886			561			325		
Connecticut									
. Crude rate		10.2	-		13.3	-		7.3	-
. AAMR (1940 Std. mil.)		10.1	0.37		13.3	0.60		6.9	0.44
. AAMR (1970 Std. mil.)		9.9	0.36		12.9	0.57		6.9	0.43
. AAMR (2000 Std. mil.)		10.2	0.35		13.5	0.57		7.1	0.40
. HP 2000 (1940 Std. mil.)		10.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		16.3	-		22.4	-		10.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is lower
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.18	0	0.0	0.00	1	0.4	0.37
5-9	1	0.2	0.17	1	0.3	0.33	0	0.0	0.00
10-14	1	0.2	0.18	1	0.4	0.35	0	0.0	0.00
15-19	2	0.4	0.28	2	0.8	0.54	0	0.0	0.00
20-24	7	1.5	0.57	7	2.9	1.11	0	0.0	0.00
25-29	2	0.4	0.25	2	0.7	0.50	0	0.0	0.00
30-34	12	1.7	0.50	10	2.9	0.90	2	0.6	0.41
35-39	6	0.8	0.32	5	1.3	0.58	1	0.3	0.26
40-44	10	1.4	0.44	7	2.0	0.74	3	0.8	0.47
45-49	12	1.9	0.56	9	2.9	0.97	3	0.9	0.55
50-54	10	1.9	0.59	8	3.0	1.08	2	0.7	0.52
55-59	10	2.5	0.78	8	4.1	1.44	2	1.0	0.68
60-64	13	3.9	1.08	11	6.8	2.05	2	1.2	0.82
65-69	35	10.0	1.69	25	15.5	3.11	10	5.3	1.68
70-74	43	13.1	1.99	23	16.1	3.37	20	10.7	2.39
75-79	79	27.5	3.09	52	44.3	6.14	27	15.9	3.05
80-84	115	58.5	5.46	51	70.6	9.89	64	51.5	6.44
85+	242	141.7	9.11	92	197.3	20.57	150	120.8	9.86
Total deaths	601			314			287		
Connecticut									
. Crude rate		6.9	-		7.4	-		6.5	-
. AAMR (1940 Std. mil.)		2.6	0.14		3.9	0.26		1.5	0.12
. AAMR (1970 Std. mil.)		3.9	0.17		5.7	0.33		2.6	0.17
. AAMR (2000 Std. mil.)		5.8	0.24		8.2	0.47		4.1	0.25
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.6	-		3.6	-		1.7	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	10	1.8	0.57	8	2.8	1.00	2	0.7	0.53
5-9	2	0.3	0.24	2	0.7	0.47	0	0.0	0.00
10-14	1	0.2	0.18	0	0.0	0.00	1	0.4	0.37
15-19	5	1.0	0.44	5	1.9	0.85	0	0.0	0.00
20-24	7	1.5	0.57	7	2.9	1.11	0	0.0	0.00
25-29	7	1.3	0.47	6	2.1	0.87	1	0.4	0.36
30-34	5	0.7	0.32	5	1.4	0.64	0	0.0	0.00
35-39	8	1.0	0.36	7	1.8	0.68	1	0.3	0.26
40-44	2	0.3	0.20	2	0.6	0.40	0	0.0	0.00
45-49	3	0.5	0.28	3	1.0	0.56	0	0.0	0.00
50-54	4	0.7	0.37	4	1.5	0.76	0	0.0	0.00
55-59	3	0.7	0.43	3	1.5	0.88	0	0.0	0.00
60-64	2	0.6	0.42	0	0.0	0.00	2	1.2	0.82
65-69	2	0.6	0.41	1	0.6	0.62	1	0.5	0.53
70-74	2	0.6	0.43	1	0.7	0.70	1	0.5	0.54
75-79	5	1.7	0.78	3	2.6	1.48	2	1.2	0.83
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	2	1.2	0.83	1	2.1	2.14	1	0.8	0.81
Total deaths	70			58			12		
Connecticut									
. Crude rate		0.8	-		1.4	-		0.3	-
. AAMR (1940 Std. mil.)		0.8	0.11		1.5	0.20		-	-
. AAMR (1970 Std. mil.)		0.8	0.10		1.4	0.19		-	-
. AAMR (2000 Std. mil.)		0.8	0.10		1.4	0.18		-	-
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.5	-		2.3	-		0.4	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	0.9	0.41	2	0.7	0.50	3	1.1	0.65
5-9	4	0.7	0.34	1	0.3	0.33	3	1.0	0.60
10-14	2	0.4	0.25	2	0.7	0.50	0	0.0	0.00
15-19	1	0.2	0.20	0	0.0	0.00	1	0.4	0.41
20-24	4	0.9	0.43	4	1.7	0.84	0	0.0	0.00
25-29	1	0.2	0.18	1	0.4	0.36	0	0.0	0.00
30-34	3	0.4	0.25	2	0.6	0.40	1	0.3	0.29
35-39	5	0.6	0.29	5	1.3	0.58	0	0.0	0.00
40-44	6	0.8	0.34	4	1.1	0.56	2	0.5	0.39
45-49	6	1.0	0.39	3	1.0	0.56	3	0.9	0.55
50-54	6	1.1	0.46	3	1.1	0.66	3	1.1	0.63
55-59	5	1.2	0.55	2	1.0	0.72	3	1.5	0.84
60-64	4	1.2	0.60	2	1.2	0.87	2	1.2	0.82
65-69	4	1.1	0.57	3	1.9	1.08	1	0.5	0.53
70-74	8	2.4	0.86	4	2.8	1.40	4	2.1	1.07
75-79	5	1.7	0.78	1	0.9	0.85	4	2.3	1.17
80-84	9	4.6	1.53	6	8.3	3.39	3	2.4	1.39
85+	2	1.2	0.83	0	0.0	0.00	2	1.6	1.14
Total deaths	80			45			35		
Connecticut									
. Crude rate		0.9	-		1.1	-		0.8	-
. AAMR (1940 Std. mil.)		0.8	0.09		0.9	0.15		0.6	0.12
. AAMR (1970 Std. mil.)		0.8	0.10		1.0	0.15		0.7	0.13
. AAMR (2000 Std. mil.)		0.9	0.10		1.1	0.16		0.7	0.12
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.0	-		1.4	-		0.9	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	10	1.8	0.57	8	2.8	0.99	2	0.7	0.52
15-19	27	5.3	1.02	19	7.2	1.66	8	3.2	1.15
20-24	45	9.7	1.45	37	15.5	2.56	8	3.6	1.26
25-29	51	9.1	1.28	44	15.6	2.36	7	2.5	0.96
30-34	96	13.7	1.40	75	21.4	2.47	21	6.0	1.31
35-39	92	11.9	1.24	67	17.3	2.11	25	6.4	1.29
40-44	100	13.8	1.38	77	21.6	2.46	23	6.3	1.31
45-49	60	9.6	1.24	48	15.6	2.25	12	3.8	1.10
50-54	64	11.9	1.49	42	16.0	2.47	22	8.1	1.72
55-59	41	10.2	1.59	34	17.3	2.96	7	3.4	1.28
60-64	31	9.3	1.66	27	16.7	3.21	4	2.3	1.15
65-69	35	10.0	1.69	25	15.5	3.11	10	5.3	1.68
70-74	29	8.8	1.64	22	15.4	3.29	7	3.7	1.42
75-79	28	9.7	1.84	24	20.4	4.17	4	2.3	1.17
80-84	30	15.3	2.79	21	29.1	6.34	9	7.2	2.42
85+	20	11.7	2.62	16	34.3	8.58	4	3.2	1.61
Total deaths	759			586			173		
Connecticut									
. Crude rate		8.8	-		13.9	-		3.9	-
. AAMR (1940 Std. mil.)		7.9	0.31		12.5	0.54		3.5	0.29
. AAMR (1970 Std. mil.)		7.4	0.29		12.0	0.52		3.3	0.27
. AAMR (2000 Std. mil.)		8.4	0.31		13.7	0.57		3.7	0.29
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.8	-		17.9	-		4.0	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	10	1.8	0.57	5	1.8	0.79	5	1.9	0.84
5-9	3	0.5	0.29	2	0.7	0.47	1	0.3	0.35
10-14	6	1.1	0.44	3	1.1	0.61	3	1.1	0.63
15-19	27	5.3	1.02	22	8.4	1.78	5	2.0	0.91
20-24	31	6.7	1.20	30	12.6	2.30	1	0.4	0.44
25-29	25	4.5	0.90	18	6.4	1.51	7	2.5	0.96
30-34	18	2.6	0.61	14	4.0	1.07	4	1.1	0.57
35-39	19	2.5	0.56	13	3.4	0.93	6	1.5	0.63
40-44	24	3.3	0.68	18	5.0	1.19	6	1.6	0.67
45-49	7	1.1	0.42	4	1.3	0.65	3	0.9	0.55
50-54	20	3.7	0.83	12	4.6	1.32	8	2.9	1.04
55-59	4	1.0	0.50	2	1.0	0.72	2	1.0	0.68
60-64	7	2.1	0.79	4	2.5	1.24	3	1.7	1.00
65-69	6	1.7	0.70	5	3.1	1.39	1	0.5	0.53
70-74	5	1.5	0.68	4	2.8	1.40	1	0.5	0.54
75-79	3	1.0	0.60	1	0.9	0.85	2	1.2	0.83
80-84	3	1.5	0.88	1	1.4	1.38	2	1.6	1.14
85+	1	0.6	0.59	0	0.0	0.00	1	0.8	0.81
Total deaths	219			158			61		
Connecticut									
. Crude rate		2.5	-		3.7	-		1.4	-
. AAMR (1940 Std. mil.)		2.8	0.20		4.2	0.35		1.4	0.19
. AAMR (1970 Std. mil.)		2.7	0.19		3.9	0.33		1.4	0.19
. AAMR (2000 Std. mil.)		2.6	0.18		3.8	0.31		1.4	0.18
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.5	-		13.3	-		3.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide (E960-E969)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	10	1.8	0.57	5	1.8	0.79	5	1.9	0.84
5-9	3	0.5	0.29	2	0.7	0.47	1	0.3	0.35
10-14	6	1.1	0.44	3	1.1	0.61	3	1.1	0.63
15-19	27	5.3	1.02	22	8.4	1.78	5	2.0	0.91
20-24	31	6.7	1.20	30	12.6	2.30	1	0.4	0.44
25-29	25	4.5	0.90	18	6.4	1.51	7	2.5	0.96
30-34	17	2.4	0.59	13	3.7	1.03	4	1.1	0.57
35-39	19	2.5	0.56	13	3.4	0.93	6	1.5	0.63
40-44	22	3.0	0.65	16	4.5	1.12	6	1.6	0.67
45-49	7	1.1	0.42	4	1.3	0.65	3	0.9	0.55
50-54	20	3.7	0.83	12	4.6	1.32	8	2.9	1.04
55-59	4	1.0	0.50	2	1.0	0.72	2	1.0	0.68
60-64	7	2.1	0.79	4	2.5	1.24	3	1.7	1.00
65-69	6	1.7	0.70	5	3.1	1.39	1	0.5	0.53
70-74	5	1.5	0.68	4	2.8	1.40	1	0.5	0.54
75-79	3	1.0	0.60	1	0.9	0.85	2	1.2	0.83
80-84	3	1.5	0.88	1	1.4	1.38	2	1.6	1.14
85+	1	0.6	0.59	0	0.0	0.00	1	0.8	0.81
Total deaths	216			155			61		
Connecticut									
. Crude rate		2.5	-		3.7	-		1.4	-
. AAMR (1940 Std. mil.)		2.8	0.20		4.2	0.35		1.4	0.19
. AAMR (1970 Std. mil.)		2.6	0.19		3.9	0.33		1.4	0.19
. AAMR (2000 Std. mil.)		2.6	0.18		3.8	0.30		1.4	0.18
. HP 2000 (1940 Std. mil.)		5.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.3	-		13.1	-		3.6	-
. HP 2000 (1940 Std. mil.)		7.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	1.1	0.45	1	0.4	0.35	5	1.9	0.84
5-9	13	2.2	0.61	11	3.6	1.09	2	0.7	0.49
10-14	20	3.6	0.80	10	3.5	1.11	10	3.7	1.16
15-19	97	19.0	1.93	56	21.3	2.84	41	16.6	2.60
20-24	91	19.7	2.06	65	27.3	3.39	26	11.6	2.27
25-29	76	13.6	1.56	57	20.3	2.68	19	6.9	1.58
30-34	76	10.9	1.25	54	15.4	2.10	22	6.3	1.34
35-39	63	8.1	1.02	40	10.3	1.63	23	5.9	1.24
40-44	54	7.5	1.02	37	10.4	1.71	17	4.6	1.12
45-49	60	9.6	1.24	44	14.3	2.15	16	5.1	1.27
50-54	44	8.2	1.24	30	11.4	2.09	14	5.1	1.37
55-59	23	5.7	1.19	12	6.1	1.76	11	5.3	1.61
60-64	37	11.0	1.82	21	13.0	2.84	16	9.2	2.31
65-69	34	9.7	1.67	22	13.7	2.91	12	6.4	1.84
70-74	47	14.3	2.08	25	17.6	3.51	22	11.8	2.51
75-79	47	16.3	2.38	27	23.0	4.43	20	11.7	2.63
80-84	46	23.4	3.45	23	31.8	6.64	23	18.5	3.86
85+	31	18.2	3.26	11	23.6	7.11	20	16.1	3.60
Total deaths	865			546			319		
Connecticut									
. Crude rate		10.0	-		13.0	-		7.2	-
. AAMR (1940 Std. mil.)		9.8	0.37		12.9	0.59		6.8	0.43
. AAMR (1970 Std. mil.)		9.6	0.35		12.6	0.57		6.8	0.42
. AAMR (2000 Std. mil.)		9.9	0.34		13.1	0.56		7.0	0.40
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.8	-		21.9	-		10.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.18	0	0.0	0.00	1	0.4	0.37
5-9	1	0.2	0.17	1	0.3	0.33	0	0.0	0.00
10-14	1	0.2	0.18	1	0.4	0.35	0	0.0	0.00
15-19	2	0.4	0.28	2	0.8	0.54	0	0.0	0.00
20-24	7	1.5	0.57	7	2.9	1.11	0	0.0	0.00
25-29	4	0.7	0.36	3	1.1	0.62	1	0.4	0.36
30-34	13	1.9	0.52	11	3.1	0.95	2	0.6	0.41
35-39	6	0.8	0.32	5	1.3	0.58	1	0.3	0.26
40-44	11	1.5	0.46	8	2.2	0.79	3	0.8	0.47
45-49	15	2.4	0.62	12	3.9	1.13	3	0.9	0.55
50-54	12	2.2	0.65	9	3.4	1.14	3	1.1	0.63
55-59	8	2.0	0.70	6	3.1	1.25	2	1.0	0.68
60-64	15	4.5	1.16	13	8.0	2.23	2	1.2	0.82
65-69	30	8.6	1.57	23	14.3	2.98	7	3.7	1.41
70-74	34	10.3	1.77	22	15.4	3.29	12	6.4	1.85
75-79	64	22.2	2.78	43	36.6	5.59	21	12.3	2.69
80-84	76	38.7	4.44	37	51.2	8.42	39	31.4	5.03
85+	141	82.6	6.95	52	111.5	15.47	89	71.7	7.60
Total deaths	441			255			186		
Connecticut									
. Crude rate		5.1	-		6.1	-		4.2	-
. AAMR (1940 Std. mil.)		2.3	0.14		3.6	0.25		1.1	0.12
. AAMR (1970 Std. mil.)		3.1	0.16		4.7	0.31		1.8	0.15
. AAMR (2000 Std. mil.)		4.3	0.21		6.4	0.41		2.7	0.21
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.5	-		3.6	-		1.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.18	1	0.4	0.35	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	5	0.9	0.40	4	1.4	0.70	1	0.4	0.37
15-19	23	4.5	0.94	19	7.2	1.66	4	1.6	0.81
20-24	48	10.4	1.50	45	18.9	2.82	3	1.3	0.77
25-29	42	7.5	1.16	36	12.8	2.13	6	2.2	0.89
30-34	53	7.6	1.04	51	14.6	2.04	2	0.6	0.41
35-39	46	5.9	0.87	37	9.6	1.57	9	2.3	0.77
40-44	47	6.5	0.95	40	11.2	1.77	7	1.9	0.72
45-49	21	3.4	0.73	16	5.2	1.30	5	1.6	0.71
50-54	35	6.5	1.10	26	9.9	1.94	9	3.3	1.10
55-59	19	4.7	1.08	16	8.1	2.03	3	1.5	0.84
60-64	20	6.0	1.34	19	11.8	2.70	1	0.6	0.58
65-69	15	4.3	1.11	14	8.7	2.33	1	0.5	0.53
70-74	15	4.6	1.18	15	10.5	2.72	0	0.0	0.00
75-79	9	3.1	1.04	9	7.7	2.56	0	0.0	0.00
80-84	14	7.1	1.90	14	19.4	5.18	0	0.0	0.00
85+	9	5.3	1.76	8	17.2	6.07	1	0.8	0.81
Total deaths	422			370			52		
Connecticut									
. Crude rate		4.9	-		8.8	-		1.2	-
. AAMR (1940 Std. mil.)		4.8	0.25		8.5	0.47		1.2	0.18
. AAMR (1970 Std. mil.)		4.5	0.23		8.0	0.44		1.1	0.16
. AAMR (2000 Std. mil.)		4.8	0.24		8.8	0.46		1.2	0.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.8	-		22.3	-		3.3	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	0.5	0.31	3	1.1	0.61	0	0.0	0.00
15-19	6	1.2	0.48	5	1.9	0.85	1	0.4	0.41
20-24	19	4.1	0.94	17	7.1	1.73	2	0.9	0.63
25-29	20	3.6	0.80	19	6.8	1.55	1	0.4	0.36
30-34	41	5.9	0.92	40	11.4	1.81	1	0.3	0.29
35-39	33	4.3	0.74	27	7.0	1.34	6	1.5	0.63
40-44	34	4.7	0.81	28	7.9	1.48	6	1.6	0.67
45-49	17	2.7	0.66	14	4.5	1.22	3	0.9	0.55
50-54	22	4.1	0.88	17	6.5	1.57	5	1.8	0.82
55-59	15	3.7	0.96	13	6.6	1.83	2	1.0	0.68
60-64	17	5.1	1.23	17	10.5	2.55	0	0.0	0.00
65-69	14	4.0	1.07	14	8.7	2.33	0	0.0	0.00
70-74	15	4.6	1.18	15	10.5	2.72	0	0.0	0.00
75-79	9	3.1	1.04	9	7.7	2.56	0	0.0	0.00
80-84	14	7.1	1.90	14	19.4	5.18	0	0.0	0.00
85+	9	5.3	1.76	8	17.2	6.07	1	0.8	0.81
Total deaths	288			260			28		
Connecticut									
. Crude rate		3.3	-		6.2	-		0.6	-
. AAMR (1940 Std. mil.)		2.9	0.19		5.4	0.36		0.6	0.12
. AAMR (1970 Std. mil.)		2.8	0.17		5.3	0.34		0.5	0.11
. AAMR (2000 Std. mil.)		3.2	0.19		6.1	0.38		0.6	0.12
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.2	-		11.3	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.18	1	0.4	0.35	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	0.4	0.25	1	0.4	0.35	1	0.4	0.37
15-19	16	3.1	0.78	13	4.9	1.37	3	1.2	0.70
20-24	28	6.0	1.14	27	11.3	2.18	1	0.4	0.44
25-29	21	3.8	0.82	16	5.7	1.42	5	1.8	0.81
30-34	10	1.4	0.45	9	2.6	0.86	1	0.3	0.29
35-39	12	1.5	0.45	10	2.6	0.82	2	0.5	0.36
40-44	9	1.2	0.41	8	2.2	0.79	1	0.3	0.27
45-49	4	0.6	0.32	2	0.6	0.46	2	0.6	0.45
50-54	13	2.4	0.67	9	3.4	1.14	4	1.5	0.73
55-59	2	0.5	0.35	1	0.5	0.51	1	0.5	0.48
60-64	2	0.6	0.42	1	0.6	0.62	1	0.6	0.58
65-69	1	0.3	0.29	0	0.0	0.00	1	0.5	0.53
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	121			98			23		
Connecticut									
. Crude rate		1.4	-		2.3	-		0.5	-
. AAMR (1940 Std. mil.)		1.7	0.16		2.8	0.29		0.6	0.13
. AAMR (1970 Std. mil.)		1.6	0.15		2.5	0.27		0.6	0.12
. AAMR (2000 Std. mil.)		1.5	0.14		2.4	0.25		0.5	0.11
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.9	-		9.8	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.4	0.25	2	0.7	0.50	0	0.0	0.00
30-34	20	2.9	0.64	16	4.6	1.14	4	1.1	0.57
35-39	27	3.5	0.67	20	5.2	1.16	7	1.8	0.68
40-44	47	6.5	0.95	35	9.8	1.66	12	3.3	0.94
45-49	71	11.4	1.35	58	18.8	2.47	13	4.1	1.14
50-54	56	10.5	1.40	41	15.6	2.44	15	5.5	1.42
55-59	35	8.7	1.47	23	11.7	2.44	12	5.8	1.68
60-64	50	14.9	2.11	39	24.1	3.86	11	6.3	1.91
65-69	46	13.2	1.94	25	15.5	3.11	21	11.2	2.44
70-74	30	9.1	1.66	21	14.7	3.22	9	4.8	1.61
75-79	25	8.7	1.74	16	13.6	3.41	9	5.3	1.76
80-84	13	6.6	1.84	11	15.2	4.59	2	1.6	1.14
85+	6	3.5	1.43	4	8.6	4.29	2	1.6	1.14
Total deaths	428			311			117		
Connecticut									
. Crude rate		4.9	-		7.4	-		2.6	-
. AAMR (1940 Std. mil.)		3.9	0.19		5.9	0.34		2.0	0.19
. AAMR (1970 Std. mil.)		4.0	0.20		6.2	0.36		2.1	0.20
. AAMR (2000 Std. mil.)		4.7	0.23		7.2	0.41		2.4	0.22
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.4	-		10.1	-		3.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.4	0.26	1	0.4	0.35	1	0.4	0.37
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	12	2.4	0.68	8	3.0	1.07	4	1.6	0.81
20-24	39	8.4	1.35	30	12.6	2.30	9	4.0	1.33
25-29	58	10.4	1.37	46	16.4	2.41	12	4.3	1.25
30-34	124	17.7	1.59	90	25.7	2.71	34	9.7	1.67
35-39	147	19.0	1.56	113	29.2	2.75	34	8.8	1.50
40-44	131	18.1	1.58	100	28.0	2.80	31	8.4	1.52
45-49	83	13.3	1.46	54	17.5	2.39	29	9.2	1.70
50-54	23	4.3	0.90	14	5.3	1.42	9	3.3	1.10
55-59	18	4.5	1.05	8	4.1	1.44	10	4.8	1.53
60-64	6	1.8	0.73	2	1.2	0.87	4	2.3	1.15
65-69	4	1.1	0.57	0	0.0	0.00	4	2.1	1.06
70-74	9	2.7	0.91	5	3.5	1.57	4	2.1	1.07
75-79	7	2.4	0.92	4	3.4	1.70	3	1.8	1.02
80-84	8	4.1	1.44	3	4.2	2.40	5	4.0	1.80
85+	6	3.5	1.43	1	2.1	2.14	5	4.0	1.80
Total deaths	677			479			198		
Connecticut									
. Crude rate		7.8	-		11.4	-		4.5	-
. AAMR (1940 Std. mil.)		7.3	0.29		10.5	0.50		4.1	0.31
. AAMR (1970 Std. mil.)		6.3	0.25		8.9	0.42		3.6	0.27
. AAMR (2000 Std. mil.)		7.5	0.29		10.8	0.49		4.3	0.31
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.3	-		7.3	-		3.1	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is higher
. THE U.S. HP 2000 target: CT rate is higher