

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

## Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Infectious diseases and parasitic disease (1-139)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	13	3.6	0.99	9	4.8	1.61	4	2.3	1.13
5-9	1	0.3	0.25	0	0.0	0.00	1	0.5	0.52
10-14	3	0.8	0.46	0	0.0	0.00	3	1.6	0.94
15-19	1	0.3	0.29	0	0.0	0.00	1	0.6	0.60
20-24	6	2.0	0.80	2	1.3	0.90	4	2.7	1.35
25-29	16	4.4	1.10	13	7.1	1.97	3	1.7	0.96
30-34	43	9.4	1.43	29	12.7	2.35	14	6.1	1.64
35-39	65	12.6	1.56	45	17.5	2.61	20	7.7	1.73
40-44	72	14.8	1.74	56	23.3	3.11	16	6.5	1.61
45-49	63	15.3	1.92	45	22.1	3.29	18	8.6	2.03
50-54	44	12.0	1.81	32	17.8	3.15	12	6.4	1.85
55-59	45	16.4	2.44	29	21.6	4.02	16	11.4	2.84
60-64	51	23.0	3.22	32	29.8	5.28	19	16.6	3.80
65-69	73	31.9	3.73	39	36.9	5.90	34	27.6	4.73
70-74	100	45.9	4.59	50	53.0	7.49	50	40.5	5.73
75-79	134	69.6	6.01	75	95.2	10.99	59	51.9	6.75
80-84	189	142.3	10.35	81	164.9	18.32	108	129.1	12.42
85+	278	240.2	14.40	87	272.5	29.22	191	227.9	16.49
Total deaths	1,197			624			573		
Connecticut									
. Crude rate		20.8	-		22.2	-		19.4	-
. AAMR (1940 Std. mil.)		10.9	0.38		14.2	0.62		7.8	0.44
. AAMR (1970 Std. mil.)		13.3	0.41		17.0	0.70		10.2	0.49
. AAMR (2000 Std. mil.)		17.9	0.52		22.7	0.92		13.9	0.60
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.5	-		18.1	-		9.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----HIV infection (042-044)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.3	0.28	0	0.0	0.00	1	0.6	0.57
5-9	1	0.3	0.25	0	0.0	0.00	1	0.5	0.52
10-14	2	0.5	0.38	0	0.0	0.00	2	1.1	0.77
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	3	1.0	0.57	1	0.6	0.64	2	1.3	0.95
25-29	14	3.8	1.03	11	6.0	1.81	3	1.7	0.96
30-34	35	7.6	1.29	25	10.9	2.18	10	4.4	1.38
35-39	47	9.1	1.33	35	13.6	2.30	12	4.6	1.34
40-44	45	9.2	1.37	38	15.8	2.56	7	2.8	1.07
45-49	34	8.2	1.41	26	12.8	2.50	8	3.8	1.35
50-54	12	3.3	0.95	12	6.7	1.93	0	0.0	0.00
55-59	15	5.5	1.41	12	9.0	2.59	3	2.1	1.23
60-64	5	2.3	1.01	5	4.7	2.09	0	0.0	0.00
65-69	4	1.7	0.87	4	3.8	1.89	0	0.0	0.00
70-74	1	0.5	0.46	1	1.1	1.06	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	219			170			49		
Connecticut									
. Crude rate		3.8	-		6.1	-		1.7	-
. AAMR (1940 Std. mil.)		3.5	0.24		5.3	0.41		1.6	0.24
. AAMR (1970 Std. mil.)		3.0	0.21		4.7	0.36		1.4	0.22
. AAMR (2000 Std. mil.)		3.6	0.25		5.7	0.44		1.6	0.24
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.7	-		9.1	-		2.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	7	1.9	0.73	3	1.6	0.93	4	2.3	1.13
5-9	9	2.3	0.76	3	1.5	0.86	6	3.1	1.28
10-14	7	1.9	0.71	5	2.6	1.17	2	1.1	0.77
15-19	6	1.8	0.72	3	1.7	0.98	3	1.8	1.05
20-24	22	7.2	1.54	15	9.6	2.47	7	4.7	1.78
25-29	19	5.2	1.20	11	6.0	1.81	8	4.4	1.57
30-34	60	13.1	1.69	24	10.5	2.14	36	15.7	2.62
35-39	121	23.5	2.13	53	20.6	2.83	68	26.3	3.19
40-44	224	45.9	3.07	105	43.7	4.26	119	48.0	4.40
45-49	349	84.5	4.52	148	72.7	5.97	201	96.1	6.78
50-54	613	167.3	6.76	270	150.3	9.15	343	183.7	9.92
55-59	742	270.2	9.92	391	291.8	14.76	351	249.6	13.32
60-64	1,029	463.7	14.46	543	506.4	21.73	486	423.8	19.22
65-69	1,540	672.3	17.13	862	814.8	27.75	678	550.0	21.12
70-74	2,018	926.7	20.63	1,040	1102.3	34.18	978	792.5	25.34
75-79	2,314	1201.7	24.98	1,179	1496.5	43.58	1,135	997.5	29.61
80-84	2,024	1524.0	33.87	1,011	2058.0	64.73	1,013	1210.5	38.03
85+	2,090	1805.6	39.50	834	2612.4	90.46	1,256	1498.3	42.28
Total deaths	13,194			6,500			6,694		
Connecticut									
. Crude rate		228.8	-		231.5	-		226.1	-
. AAMR (1940 Std. mil.)		115.8	1.16		132.5	1.79		104.2	1.54
. AAMR (1970 Std. mil.)		154.7	1.42		182.9	2.31		136.3	1.82
. AAMR (2000 Std. mil.)		195.7	1.71		236.6	2.96		171.1	2.14
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		125.6	-		150.4	-		107.4	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Oral cavity or pharynx (140.0-149.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.2	0.22	1	0.4	0.44	0	0.0	0.00
35-39	5	1.0	0.43	5	1.9	0.87	0	0.0	0.00
40-44	3	0.6	0.35	2	0.8	0.59	1	0.4	0.40
45-49	3	0.7	0.42	2	1.0	0.69	1	0.5	0.48
50-54	5	1.4	0.61	4	2.2	1.11	1	0.5	0.54
55-59	19	6.9	1.59	13	9.7	2.69	6	4.3	1.74
60-64	15	6.8	1.75	10	9.3	2.95	5	4.4	1.95
65-69	21	9.2	2.00	15	14.2	3.66	6	4.9	1.99
70-74	15	6.9	1.78	6	6.4	2.60	9	7.3	2.43
75-79	30	15.6	2.84	18	22.8	5.39	12	10.5	3.04
80-84	20	15.1	3.37	12	24.4	7.05	8	9.6	3.38
85+	23	19.9	4.14	8	25.1	8.86	15	17.9	4.62
Total deaths	160			96			64		
Connecticut									
. Crude rate		2.8	-		3.4	-		2.2	-
. AAMR (1940 Std. mil.)		1.5	0.14		2.2	0.24		0.9	0.14
. AAMR (1970 Std. mil.)		2.0	0.16		2.8	0.29		1.3	0.18
. AAMR (2000 Std. mil.)		2.4	0.19		3.4	0.35		1.6	0.21
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.0	-		3.1	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available







## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Rectal cancer (154)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.3	0.27	0	0.0	0.00	1	0.6	0.55
30-34	1	0.2	0.22	0	0.0	0.00	1	0.4	0.44
35-39	2	0.4	0.27	2	0.8	0.55	0	0.0	0.00
40-44	5	1.0	0.46	3	1.2	0.72	2	0.8	0.57
45-49	3	0.7	0.42	2	1.0	0.69	1	0.5	0.48
50-54	6	1.6	0.67	5	2.8	1.24	1	0.5	0.54
55-59	14	5.1	1.36	8	6.0	2.11	6	4.3	1.74
60-64	14	6.3	1.69	7	6.5	2.47	7	6.1	2.31
65-69	19	8.3	1.90	8	7.6	2.67	11	8.9	2.69
70-74	43	19.7	3.01	29	30.7	5.71	14	11.3	3.03
75-79	40	20.8	3.28	17	21.6	5.23	23	20.2	4.21
80-84	36	27.1	4.52	17	34.6	8.39	19	22.7	5.21
85+	44	38.0	5.73	15	47.0	12.13	29	34.6	6.42
Total deaths	228			113			115		
Connecticut									
. Crude rate		4.0	-		4.0	-		3.9	-
. AAMR (1940 Std. mil.)		1.9	0.14		2.3	0.23		1.5	0.18
. AAMR (1970 Std. mil.)		2.6	0.18		3.1	0.30		2.1	0.22
. AAMR (2000 Std. mil.)		3.3	0.22		4.1	0.39		2.8	0.27
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.3	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.3	0.27	0	0.0	0.00	1	0.6	0.55
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	3	0.6	0.34	1	0.4	0.39	2	0.8	0.55
40-44	8	1.6	0.58	6	2.5	1.02	2	0.8	0.57
45-49	20	4.8	1.08	8	3.9	1.39	12	5.7	1.66
50-54	34	9.3	1.59	15	8.3	2.16	19	10.2	2.33
55-59	43	15.7	2.39	27	20.1	3.88	16	11.4	2.84
60-64	50	22.5	3.19	36	33.6	5.60	14	12.2	3.26
65-69	101	44.1	4.39	61	57.7	7.38	40	32.4	5.13
70-74	122	56.0	5.07	55	58.3	7.86	67	54.3	6.63
75-79	137	71.1	6.08	57	72.3	9.58	80	70.3	7.86
80-84	130	97.9	8.58	62	126.2	16.03	68	81.3	9.85
85+	107	92.4	8.94	29	90.8	16.87	78	93.0	10.54
Total deaths	756			357			399		
Connecticut									
. Crude rate		13.1	-		12.7	-		13.5	-
. AAMR (1940 Std. mil.)		6.5	0.27		7.6	0.43		5.5	0.34
. AAMR (1970 Std. mil.)		8.8	0.34		10.3	0.55		7.6	0.41
. AAMR (2000 Std. mil.)		11.2	0.41		12.8	0.68		9.9	0.50
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.2	-		7.3	-		5.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Lung and other respiratory cancer (160-165)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	4	0.9	0.44	2	0.9	0.62	2	0.9	0.62
35-39	14	2.7	0.73	10	3.9	1.23	4	1.5	0.77
40-44	39	8.0	1.28	20	8.3	1.86	19	7.7	1.76
45-49	89	21.6	2.28	50	24.5	3.47	39	18.6	2.99
50-54	165	45.0	3.51	84	46.8	5.10	81	43.4	4.82
55-59	241	87.8	5.65	140	104.5	8.83	101	71.8	7.15
60-64	371	167.2	8.68	204	190.2	13.32	167	145.6	11.27
65-69	534	233.1	10.09	322	304.4	16.96	212	172.0	11.81
70-74	640	293.9	11.62	364	385.8	20.22	276	223.7	13.46
75-79	647	336.0	13.21	344	436.6	23.54	303	266.3	15.30
80-84	503	378.7	16.89	260	529.3	32.82	243	290.4	18.63
85+	337	291.1	15.86	174	545.0	41.32	163	194.4	15.23
Total deaths	3,584			1,974			1,610		
Connecticut									
. Crude rate		62.1	-		70.3	-		54.4	-
. AAMR (1940 Std. mil.)		33.8	0.63		42.4	1.03		27.3	0.79
. AAMR (1970 Std. mil.)		44.5	0.78		57.0	1.31		35.4	0.95
. AAMR (2000 Std. mil.)		53.7	0.90		70.6	1.60		42.4	1.08
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		38.7	-		52.7	-		27.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	4	0.9	0.44	2	0.9	0.62	2	0.9	0.62
35-39	13	2.5	0.70	9	3.5	1.17	4	1.5	0.77
40-44	39	8.0	1.28	20	8.3	1.86	19	7.7	1.76
45-49	88	21.3	2.27	49	24.1	3.44	39	18.6	2.99
50-54	156	42.6	3.41	76	42.3	4.85	80	42.8	4.79
55-59	231	84.1	5.53	133	99.3	8.61	98	69.7	7.04
60-64	359	161.8	8.54	196	182.8	13.06	163	142.1	11.13
65-69	523	228.3	9.98	315	297.7	16.78	208	168.7	11.70
70-74	627	287.9	11.50	354	375.2	19.94	273	221.2	13.39
75-79	631	327.7	13.04	334	423.9	23.20	297	261.0	15.15
80-84	489	368.2	16.65	249	506.9	32.12	240	286.8	18.51
85+	322	278.2	15.50	164	513.7	40.11	158	188.5	14.99
Total deaths	3,482			1,901			1,581		
Connecticut									
. Crude rate		60.4	-		67.7	-		53.4	-
. AAMR (1940 Std. mil.)		32.9	0.62		40.8	1.01		26.8	0.78
. AAMR (1970 Std. mil.)		43.2	0.77		54.9	1.28		34.8	0.94
. AAMR (2000 Std. mil.)		52.2	0.89		67.9	1.57		41.6	1.07
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		37.4	-		50.4	-		27.0	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.4	0.31	1	0.4	0.44	1	0.4	0.44
35-39	6	1.2	0.48	2	0.8	0.55	4	1.5	0.77
40-44	7	1.4	0.54	3	1.2	0.72	4	1.6	0.81
45-49	8	1.9	0.69	8	3.9	1.39	0	0.0	0.00
50-54	11	3.0	0.91	7	3.9	1.47	4	2.1	1.07
55-59	14	5.1	1.36	5	3.7	1.67	9	6.4	2.13
60-64	12	5.4	1.56	8	7.5	2.64	4	3.5	1.74
65-69	17	7.4	1.80	12	11.3	3.27	5	4.1	1.81
70-74	23	10.6	2.20	13	13.8	3.82	10	8.1	2.56
75-79	28	14.5	2.75	14	17.8	4.75	14	12.3	3.29
80-84	21	15.8	3.45	7	14.2	5.39	14	16.7	4.47
85+	27	23.3	4.49	9	28.2	9.40	18	21.5	5.06
Total deaths	176			89			87		
Connecticut									
. Crude rate		3.1	-		3.2	-		2.9	-
. AAMR (1940 Std. mil.)		1.7	0.14		2.0	0.23		1.4	0.18
. AAMR (1970 Std. mil.)		2.1	0.17		2.6	0.27		1.7	0.20
. AAMR (2000 Std. mil.)		2.7	0.20		3.2	0.34		2.2	0.24
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.7	-		2.6	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	5	1.1	0.49	-	-	-	5	2.2	0.98
35-39	17	3.3	0.80	-	-	-	17	6.6	1.60
40-44	42	8.6	1.33	-	-	-	42	17.0	2.62
45-49	65	15.7	1.95	-	-	-	65	31.1	3.85
50-54	95	25.9	2.66	-	-	-	95	50.9	5.22
55-59	62	22.6	2.87	-	-	-	62	44.1	5.60
60-64	86	38.8	4.18	-	-	-	86	75.0	8.09
65-69	112	48.9	4.62	-	-	-	112	90.9	8.58
70-74	162	74.4	5.85	-	-	-	162	131.3	10.31
75-79	124	64.4	5.78	-	-	-	124	109.0	9.79
80-84	127	95.6	8.49	-	-	-	127	151.8	13.47
85+	192	165.9	11.97	-	-	-	192	229.0	16.53
Total deaths	1,089			-			1,089		
Connecticut									
. Crude rate		-	-		-	-		36.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		19.1	0.68
. AAMR (1970 Std. mil.)		-	-		-	-		23.6	0.77
. AAMR (2000 Std. mil.)		-	-		-	-		29.0	0.90
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		19.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is lower
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									CT rate is lower



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	1	0.3	0.27	-	-	-	1	0.6	0.55
30-34	3	0.7	0.38	-	-	-	3	1.3	0.76
35-39	6	1.2	0.48	-	-	-	6	2.3	0.95
40-44	1	0.2	0.20	-	-	-	1	0.4	0.40
45-49	7	1.7	0.64	-	-	-	7	3.3	1.26
50-54	7	1.9	0.72	-	-	-	7	3.7	1.42
55-59	5	1.8	0.81	-	-	-	5	3.6	1.59
60-64	3	1.4	0.78	-	-	-	3	2.6	1.51
65-69	6	2.6	1.07	-	-	-	6	4.9	1.99
70-74	2	0.9	0.65	-	-	-	2	1.6	1.15
75-79	5	2.6	1.16	-	-	-	5	4.4	1.97
80-84	4	3.0	1.51	-	-	-	4	4.8	2.39
85+	9	7.8	2.59	-	-	-	9	10.7	3.58
Total deaths	59			-			59		
Connecticut									
. Crude rate		-	-		-	-		2.0	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.3	0.19
. AAMR (1970 Std. mil.)		-	-		-	-		1.4	0.19
. AAMR (2000 Std. mil.)		-	-		-	-		1.7	0.22
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	0	0.0	0.00	-	-	-	0	0.0	0.00
45-49	3	0.7	0.42	-	-	-	3	1.4	0.83
50-54	1	0.3	0.27	-	-	-	1	0.5	0.54
55-59	5	1.8	0.81	-	-	-	5	3.6	1.59
60-64	6	2.7	1.10	-	-	-	6	5.2	2.14
65-69	5	2.2	0.98	-	-	-	5	4.1	1.81
70-74	9	4.1	1.38	-	-	-	9	7.3	2.43
75-79	16	8.3	2.08	-	-	-	16	14.1	3.52
80-84	14	10.5	2.82	-	-	-	14	16.7	4.47
85+	17	14.7	3.56	-	-	-	17	20.3	4.92
Total deaths	76			-			76		
Connecticut									
. Crude rate		-	-		-	-		2.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.0	0.15
. AAMR (1970 Std. mil.)		-	-		-	-		1.5	0.18
. AAMR (2000 Std. mil.)		-	-		-	-		1.9	0.22
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	1	0.3	0.27	-	-	-	1	0.6	0.55
30-34	2	0.4	0.31	-	-	-	2	0.9	0.62
35-39	2	0.4	0.27	-	-	-	2	0.8	0.55
40-44	7	1.4	0.54	-	-	-	7	2.8	1.07
45-49	11	2.7	0.80	-	-	-	11	5.3	1.59
50-54	34	9.3	1.59	-	-	-	34	18.2	3.12
55-59	22	8.0	1.71	-	-	-	22	15.6	3.34
60-64	29	13.1	2.43	-	-	-	29	25.3	4.70
65-69	49	21.4	3.06	-	-	-	49	39.7	5.68
70-74	54	24.8	3.37	-	-	-	54	43.8	5.95
75-79	50	26.0	3.67	-	-	-	50	43.9	6.21
80-84	43	32.4	4.94	-	-	-	43	51.4	7.84
85+	58	50.1	6.58	-	-	-	58	69.2	9.09
Total deaths	362			-			362		
Connecticut									
. Crude rate		-	-		-	-		12.2	-
. AAMR (1940 Std. mil.)		-	-		-	-		6.3	0.39
. AAMR (1970 Std. mil.)		-	-		-	-		7.9	0.45
. AAMR (2000 Std. mil.)		-	-		-	-		9.6	0.51
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		5.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Brain & other CNS cancer (191-192)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.8	0.48	1	0.5	0.54	2	1.1	0.80
5-9	1	0.3	0.25	1	0.5	0.50	0	0.0	0.00
10-14	3	0.8	0.46	2	1.0	0.74	1	0.5	0.54
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.3	0.33	1	0.6	0.64	0	0.0	0.00
25-29	3	0.8	0.48	3	1.6	0.94	0	0.0	0.00
30-34	5	1.1	0.49	4	1.7	0.87	1	0.4	0.44
35-39	12	2.3	0.67	8	3.1	1.10	4	1.5	0.77
40-44	6	1.2	0.50	5	2.1	0.93	1	0.4	0.40
45-49	20	4.8	1.08	11	5.4	1.63	9	4.3	1.43
50-54	21	5.7	1.25	13	7.2	2.01	8	4.3	1.51
55-59	28	10.2	1.93	13	9.7	2.69	15	10.7	2.75
60-64	20	9.0	2.02	15	14.0	3.61	5	4.4	1.95
65-69	34	14.8	2.55	22	20.8	4.43	12	9.7	2.81
70-74	37	17.0	2.79	12	12.7	3.67	25	20.3	4.05
75-79	36	18.7	3.12	19	24.1	5.53	17	14.9	3.62
80-84	31	23.3	4.19	16	32.6	8.14	15	17.9	4.63
85+	25	21.6	4.32	10	31.3	9.91	15	17.9	4.62
Total deaths	286			156			130		
Connecticut									
. Crude rate		5.0	-		5.6	-		4.4	-
. AAMR (1940 Std. mil.)		3.2	0.21		3.9	0.34		2.4	0.25
. AAMR (1970 Std. mil.)		3.7	0.23		4.6	0.38		3.0	0.28
. AAMR (2000 Std. mil.)		4.4	0.26		5.5	0.45		3.6	0.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.5	-		4.1	-		2.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.6	0.39	0	0.0	0.00	2	1.1	0.80
5-9	5	1.3	0.57	0	0.0	0.00	5	2.6	1.17
10-14	2	0.5	0.38	2	1.0	0.74	0	0.0	0.00
15-19	3	0.9	0.51	1	0.6	0.57	2	1.2	0.85
20-24	7	2.3	0.87	6	3.8	1.56	1	0.7	0.67
25-29	3	0.8	0.48	3	1.6	0.94	0	0.0	0.00
30-34	9	2.0	0.66	4	1.7	0.87	5	2.2	0.98
35-39	9	1.7	0.58	5	1.9	0.87	4	1.5	0.77
40-44	13	2.7	0.74	5	2.1	0.93	8	3.2	1.14
45-49	6	1.5	0.59	1	0.5	0.49	5	2.4	1.07
50-54	23	6.3	1.31	9	5.0	1.67	14	7.5	2.00
55-59	21	7.6	1.67	14	10.4	2.79	7	5.0	1.88
60-64	33	14.9	2.59	18	16.8	3.96	15	13.1	3.38
65-69	51	22.3	3.12	39	36.9	5.90	12	9.7	2.81
70-74	72	33.1	3.90	41	43.5	6.79	31	25.1	4.51
75-79	99	51.4	5.17	62	78.7	9.99	37	32.5	5.35
80-84	71	53.5	6.34	36	73.3	12.21	35	41.8	7.07
85+	81	70.0	7.78	31	97.1	17.44	50	59.6	8.44
Total deaths	510			277			233		
Connecticut									
. Crude rate		8.8	-		9.9	-		7.9	-
. AAMR (1940 Std. mil.)		4.7	0.25		5.9	0.40		3.8	0.31
. AAMR (1970 Std. mil.)		6.1	0.29		7.9	0.49		4.8	0.35
. AAMR (2000 Std. mil.)		7.6	0.34		10.0	0.61		6.0	0.41
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.8	-		6.1	-		3.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.3	0.28	1	0.5	0.54	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	0.7	0.46	1	0.6	0.64	1	0.7	0.67
25-29	3	0.8	0.48	1	0.5	0.55	2	1.1	0.78
30-34	6	1.3	0.53	2	0.9	0.62	4	1.7	0.87
35-39	20	3.9	0.87	14	5.4	1.45	6	2.3	0.95
40-44	38	7.8	1.26	29	12.1	2.24	9	3.6	1.21
45-49	63	15.3	1.92	39	19.1	3.07	24	11.5	2.34
50-54	120	32.8	2.99	72	40.1	4.72	48	25.7	3.71
55-59	194	70.6	5.07	124	92.5	8.31	70	49.8	5.95
60-64	269	121.2	7.39	163	152.0	11.91	106	92.4	8.98
65-69	414	180.7	8.88	229	216.5	14.30	185	150.1	11.03
70-74	616	282.9	11.40	338	358.3	19.49	278	225.3	13.51
75-79	858	445.6	15.21	449	569.9	26.90	409	359.5	17.77
80-84	873	657.3	22.25	436	887.5	42.51	437	522.2	24.98
85+	1,064	919.2	28.18	359	1124.5	59.35	705	841.0	31.67
Total deaths	4,541			2,257			2,284		
Connecticut									
. Crude rate		78.7	-		80.4	-		77.2	-
. AAMR (1940 Std. mil.)		33.5	0.59		42.4	0.98		26.6	0.71
. AAMR (1970 Std. mil.)		48.6	0.77		61.6	1.32		39.1	0.91
. AAMR (2000 Std. mil.)		65.6	0.98		83.3	1.77		53.5	1.14
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									
. The U.S. AAMR (1940 Std. mil.):									
. THE U.S. HP 2000 target:									

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): No comparison is available  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.2	0.19	1	0.4	0.39	0	0.0	0.00
40-44	2	0.4	0.29	1	0.4	0.42	1	0.4	0.40
45-49	1	0.2	0.24	1	0.5	0.49	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	2	0.7	0.51	1	0.7	0.75	1	0.7	0.71
60-64	1	0.5	0.45	1	0.9	0.93	0	0.0	0.00
65-69	1	0.4	0.44	1	0.9	0.95	0	0.0	0.00
70-74	1	0.5	0.46	0	0.0	0.00	1	0.8	0.81
75-79	2	1.0	0.73	1	1.3	1.27	1	0.9	0.88
80-84	4	3.0	1.51	3	6.1	3.53	1	1.2	1.19
85+	2	1.7	1.22	0	0.0	0.00	2	2.4	1.69
Total deaths	17			10			7		
Connecticut									
. Crude rate		0.3	-		0.4	-		0.2	-
. AAMR (1940 Std. mil.)		0.2	0.04		-	-		-	-
. AAMR (1970 Std. mil.)		0.2	0.05		-	-		-	-
. AAMR (2000 Std. mil.)		0.3	0.06		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.5	-		0.5	-		0.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Alzheimer's diseases (331.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.2	0.20	1	0.4	0.42	0	0.0	0.00
45-49	1	0.2	0.24	0	0.0	0.00	1	0.5	0.48
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	3	1.1	0.63	0	0.0	0.00	3	2.1	1.23
60-64	2	0.9	0.64	2	1.9	1.32	0	0.0	0.00
65-69	14	6.1	1.63	12	11.3	3.27	2	1.6	1.15
70-74	31	14.2	2.56	9	9.5	3.18	22	17.8	3.80
75-79	66	34.3	4.22	23	29.2	6.09	43	37.8	5.76
80-84	121	91.1	8.28	42	85.5	13.19	79	94.4	10.62
85+	273	235.8	14.27	62	194.2	24.66	211	251.7	17.33
Total deaths	512			151			361		
Connecticut									
. Crude rate		8.9	-		5.4	-		12.2	-
. AAMR (1940 Std. mil.)		2.1	0.11		2.0	0.18		2.2	0.14
. AAMR (1970 Std. mil.)		4.1	0.19		3.7	0.30		4.3	0.24
. AAMR (2000 Std. mil.)		7.0	0.31		6.1	0.51		7.4	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.7	-		2.5	-		2.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Cardiovascular disease (390-459)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	19	5.2	1.20	8	4.3	1.52	11	6.2	1.88
5-9	2	0.5	0.36	1	0.5	0.50	1	0.5	0.52
10-14	6	1.6	0.65	2	1.0	0.74	4	2.2	1.09
15-19	7	2.0	0.77	6	3.4	1.39	1	0.6	0.60
20-24	11	3.6	1.09	5	3.2	1.43	6	4.0	1.65
25-29	29	8.0	1.48	18	9.8	2.31	11	6.1	1.84
30-34	42	9.2	1.42	24	10.5	2.14	18	7.9	1.85
35-39	102	19.8	1.96	75	29.1	3.36	27	10.5	2.01
40-44	193	39.5	2.85	143	59.5	4.97	50	20.2	2.85
45-49	292	70.7	4.14	214	105.1	7.18	78	37.3	4.22
50-54	432	117.9	5.67	314	174.8	9.86	118	63.2	5.82
55-59	660	240.3	9.35	473	353.0	16.23	187	133.0	9.72
60-64	833	375.4	13.01	578	539.0	22.42	255	222.4	13.92
65-69	1,452	633.9	16.63	921	870.5	28.69	531	430.7	18.69
70-74	2,184	1003.0	21.46	1,308	1386.4	38.33	876	709.9	23.98
75-79	3,407	1769.3	30.31	1,816	2305.0	54.09	1,591	1398.3	35.06
80-84	4,348	3273.8	49.65	1,984	4038.7	90.67	2,364	2824.8	58.10
85+	9,262	8001.6	83.14	2,579	8078.3	159.07	6,683	7972.4	97.52
Total deaths	23,281			10,469			12,812		
Connecticut									
. Crude rate		403.6	-		372.9	-		432.8	-
. AAMR (1940 Std. mil.)		141.0	1.15		184.5	2.01		104.9	1.28
. AAMR (1970 Std. mil.)		222.3	1.56		279.5	2.78		176.7	1.76
. AAMR (2000 Std. mil.)		329.5	2.17		399.5	3.95		275.2	2.48
. HP 2000 (1940 Std. mil.)		84.4	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		167.1	-		214.2	-		128.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									
. The U.S. AAMR (1940 Std. mil.):									
. THE U.S. HP 2000 target:									

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is higher  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.6	0.41	2	1.1	0.80	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	8	2.2	0.78	6	3.3	1.34	2	1.1	0.78
30-34	13	2.8	0.79	9	3.9	1.31	4	1.7	0.87
35-39	42	8.1	1.26	34	13.2	2.27	8	3.1	1.10
40-44	107	21.9	2.12	90	37.4	3.95	17	6.9	1.66
45-49	179	43.4	3.24	148	72.7	5.97	31	14.8	2.66
50-54	286	78.1	4.62	229	127.5	8.42	57	30.5	4.04
55-59	426	155.1	7.52	310	231.4	13.14	116	82.5	7.66
60-64	528	237.9	10.35	380	354.4	18.18	148	129.1	10.61
65-69	898	392.0	13.08	598	565.2	23.11	300	243.4	14.05
70-74	1,249	573.6	16.23	796	843.7	29.90	453	367.1	17.25
75-79	1,924	999.1	22.78	1,082	1373.4	41.75	842	740.0	25.50
80-84	2,390	1799.5	36.81	1,165	2371.5	69.48	1,225	1463.8	41.82
85+	4,981	4303.2	60.97	1,445	4526.2	119.07	3,536	4218.2	70.94
Total deaths	13,033			6,294			6,739		
Connecticut									
. Crude rate		226.0	-		224.2	-		227.7	-
. AAMR (1940 Std. mil.)		81.3	0.88		113.6	1.59		54.7	0.91
. AAMR (1970 Std. mil.)		126.6	1.19		169.4	2.17		92.8	1.27
. AAMR (2000 Std. mil.)		185.1	1.63		238.7	3.04		144.5	1.79
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		100.4	-		137.4	-		70.0	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.3	0.28	0	0.0	0.00	1	0.6	0.57
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.2	0.19	1	0.4	0.39	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	4	1.0	0.48	2	1.0	0.69	2	1.0	0.68
50-54	6	1.6	0.67	3	1.7	0.96	3	1.6	0.93
55-59	13	4.7	1.31	6	4.5	1.83	7	5.0	1.88
60-64	14	6.3	1.69	6	5.6	2.28	8	7.0	2.47
65-69	33	14.4	2.51	18	17.0	4.01	15	12.2	3.14
70-74	73	33.5	3.92	35	37.1	6.27	38	30.8	5.00
75-79	124	64.4	5.78	64	81.2	10.15	60	52.7	6.81
80-84	224	168.7	11.27	91	185.2	19.42	133	158.9	13.78
85+	630	544.3	21.68	160	501.2	39.62	470	560.7	25.86
Total deaths	1,123			386			737		
Connecticut									
. Crude rate		19.5	-		13.7	-		24.9	-
. AAMR (1940 Std. mil.)		4.9	0.18		5.2	0.29		4.7	0.24
. AAMR (1970 Std. mil.)		9.2	0.29		9.5	0.49		8.9	0.36
. AAMR (2000 Std. mil.)		15.4	0.46		15.7	0.81		15.1	0.56
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.1	-		6.6	-		5.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available







## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.8	0.48	1	0.5	0.54	2	1.1	0.80
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.3	0.27	0	0.0	0.00	1	0.5	0.54
15-19	1	0.3	0.29	1	0.6	0.57	0	0.0	0.00
20-24	1	0.3	0.33	0	0.0	0.00	1	0.7	0.67
25-29	3	0.8	0.48	1	0.5	0.55	2	1.1	0.78
30-34	9	2.0	0.66	4	1.7	0.87	5	2.2	0.98
35-39	15	2.9	0.75	11	4.3	1.29	4	1.5	0.77
40-44	31	6.4	1.14	14	5.8	1.56	17	6.9	1.66
45-49	32	7.8	1.37	15	7.4	1.90	17	8.1	1.97
50-54	39	10.6	1.70	15	8.3	2.16	24	12.9	2.62
55-59	74	26.9	3.13	49	36.6	5.22	25	17.8	3.56
60-64	66	29.7	3.66	41	38.2	5.97	25	21.8	4.36
65-69	160	69.8	5.52	90	85.1	8.97	70	56.8	6.79
70-74	309	141.9	8.07	156	165.3	13.24	153	124.0	10.02
75-79	542	281.5	12.09	251	318.6	20.11	291	255.8	14.99
80-84	763	574.5	20.80	295	600.5	34.96	468	559.2	25.85
85+	1,584	1368.4	34.38	394	1234.1	62.18	1,190	1419.6	41.15
Total deaths	3,633			1,338			2,295		
Connecticut									
. Crude rate		63.0	-		47.7	-		77.5	-
. AAMR (1940 Std. mil.)		19.5	0.40		21.1	0.65		18.0	0.51
. AAMR (1970 Std. mil.)		32.6	0.57		34.4	0.95		30.8	0.72
. AAMR (2000 Std. mil.)		50.7	0.84		52.0	1.44		49.0	1.04
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		25.9	-		27.9	-		24.1	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is higher  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Atherosclerosis (440)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.2	0.19	0	0.0	0.00	1	0.4	0.39
40-44	1	0.2	0.20	1	0.4	0.42	0	0.0	0.00
45-49	5	1.2	0.54	4	2.0	0.98	1	0.5	0.48
50-54	2	0.5	0.39	0	0.0	0.00	2	1.1	0.76
55-59	2	0.7	0.51	2	1.5	1.06	0	0.0	0.00
60-64	4	1.8	0.90	1	0.9	0.93	3	2.6	1.51
65-69	15	6.5	1.69	8	7.6	2.67	7	5.7	2.15
70-74	20	9.2	2.05	13	13.8	3.82	7	5.7	2.14
75-79	29	15.1	2.80	10	12.7	4.01	19	16.7	3.83
80-84	55	41.4	5.58	23	46.8	9.76	32	38.2	6.76
85+	194	167.6	12.03	56	175.4	23.44	138	164.6	14.01
Total deaths	328			118			210		
Connecticut									
. Crude rate		5.7	-		4.2	-		7.1	-
. AAMR (1940 Std. mil.)		1.5	0.10		1.6	0.17		1.4	0.13
. AAMR (1970 Std. mil.)		2.7	0.16		3.0	0.27		2.5	0.19
. AAMR (2000 Std. mil.)		4.5	0.25		4.9	0.46		4.3	0.30
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.1	-		2.4	-		1.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available







## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.3	0.28	1	0.5	0.54	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.3	0.27	1	0.5	0.52	0	0.0	0.00
15-19	1	0.3	0.29	0	0.0	0.00	1	0.6	0.60
20-24	2	0.7	0.46	2	1.3	0.90	0	0.0	0.00
25-29	1	0.3	0.27	1	0.5	0.55	0	0.0	0.00
30-34	1	0.2	0.22	1	0.4	0.44	0	0.0	0.00
35-39	5	1.0	0.43	2	0.8	0.55	3	1.2	0.67
40-44	12	2.5	0.71	4	1.7	0.83	8	3.2	1.14
45-49	13	3.1	0.87	6	2.9	1.20	7	3.3	1.26
50-54	36	9.8	1.64	15	8.3	2.16	21	11.2	2.45
55-59	55	20.0	2.70	21	15.7	3.42	34	24.2	4.15
60-64	98	44.2	4.46	45	42.0	6.26	53	46.2	6.35
65-69	220	96.0	6.48	104	98.3	9.64	116	94.1	8.74
70-74	356	163.5	8.66	176	186.5	14.06	180	145.9	10.87
75-79	505	262.2	11.67	236	299.6	19.50	269	236.4	14.41
80-84	523	393.8	17.22	226	460.1	30.60	297	354.9	20.59
85+	601	519.2	21.18	252	789.4	49.72	349	416.3	22.29
Total deaths	2,431			1,093			1,338		
Connecticut									
. Crude rate		42.1	-		38.9	-		45.2	-
. AAMR (1940 Std. mil.)		16.3	0.39		18.1	0.60		15.4	0.52
. AAMR (1970 Std. mil.)		24.9	0.53		28.7	0.88		23.0	0.69
. AAMR (2000 Std. mil.)		34.6	0.70		41.4	1.27		31.3	0.87
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		21.1	-		26.1	-		17.8	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.3	0.27	1	0.5	0.52	0	0.0	0.00
15-19	1	0.3	0.29	0	0.0	0.00	1	0.6	0.60
20-24	2	0.7	0.46	2	1.3	0.90	0	0.0	0.00
25-29	1	0.3	0.27	1	0.5	0.55	0	0.0	0.00
30-34	1	0.2	0.22	1	0.4	0.44	0	0.0	0.00
35-39	4	0.8	0.39	2	0.8	0.55	2	0.8	0.55
40-44	6	1.2	0.50	1	0.4	0.42	5	2.0	0.90
45-49	2	0.5	0.34	0	0.0	0.00	2	1.0	0.68
50-54	14	3.8	1.02	7	3.9	1.47	7	3.7	1.42
55-59	6	2.2	0.89	2	1.5	1.06	4	2.8	1.42
60-64	8	3.6	1.27	2	1.9	1.32	6	5.2	2.14
65-69	1	0.4	0.44	0	0.0	0.00	1	0.8	0.81
70-74	6	2.8	1.12	1	1.1	1.06	5	4.1	1.81
75-79	14	7.3	1.94	2	2.5	1.80	12	10.5	3.04
80-84	14	10.5	2.82	3	6.1	3.53	11	13.1	3.96
85+	17	14.7	3.56	1	3.1	3.13	16	19.1	4.77
Total deaths	98			26			72		
Connecticut									
. Crude rate		1.7	-		0.9	-		2.4	-
. AAMR (1940 Std. mil.)		1.0	0.12		0.8	0.16		1.2	0.17
. AAMR (1970 Std. mil.)		1.2	0.13		0.8	0.17		1.5	0.19
. AAMR (2000 Std. mil.)		1.5	0.15		0.9	0.18		1.9	0.23
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.6	-		1.1	-		1.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.3	0.28	1	0.5	0.54	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	3	0.8	0.48	1	0.5	0.55	2	1.1	0.78
30-34	6	1.3	0.53	5	2.2	0.98	1	0.4	0.44
35-39	17	3.3	0.80	9	3.5	1.17	8	3.1	1.10
40-44	41	8.4	1.31	35	14.6	2.46	6	2.4	0.99
45-49	54	13.1	1.78	37	18.2	2.99	17	8.1	1.97
50-54	53	14.5	1.99	39	21.7	3.48	14	7.5	2.00
55-59	47	17.1	2.50	33	24.6	4.29	14	10.0	2.66
60-64	55	24.8	3.34	40	37.3	5.90	15	13.1	3.38
65-69	63	27.5	3.46	38	35.9	5.83	25	20.3	4.06
70-74	66	30.3	3.73	38	40.3	6.53	28	22.7	4.29
75-79	65	33.8	4.19	21	26.7	5.82	44	38.7	5.83
80-84	50	37.6	5.32	23	46.8	9.76	27	32.3	6.21
85+	20	17.3	3.86	8	25.1	8.86	12	14.3	4.13
Total deaths	541			328			213		
Connecticut									
. Crude rate		9.4	-		11.7	-		7.2	-
. AAMR (1940 Std. mil.)		6.3	0.29		8.7	0.50		4.0	0.31
. AAMR (1970 Std. mil.)		7.2	0.32		9.8	0.55		4.9	0.35
. AAMR (2000 Std. mil.)		8.5	0.37		11.5	0.63		5.9	0.41
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.3	-		10.5	-		4.5	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Nephritis, nephrotic syndrome/nephrosis (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.6	0.39	0	0.0	0.00	2	1.1	0.80
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.29	1	0.6	0.57	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.5	0.39	1	0.5	0.55	1	0.6	0.55
30-34	1	0.2	0.22	0	0.0	0.00	1	0.4	0.44
35-39	2	0.4	0.27	0	0.0	0.00	2	0.8	0.55
40-44	6	1.2	0.50	3	1.2	0.72	3	1.2	0.70
45-49	9	2.2	0.73	6	2.9	1.20	3	1.4	0.83
50-54	14	3.8	1.02	10	5.6	1.76	4	2.1	1.07
55-59	15	5.5	1.41	7	5.2	1.97	8	5.7	2.01
60-64	20	9.0	2.02	9	8.4	2.80	11	9.6	2.89
65-69	28	12.2	2.31	11	10.4	3.13	17	13.8	3.34
70-74	71	32.6	3.87	41	43.5	6.79	30	24.3	4.44
75-79	101	52.4	5.22	52	66.0	9.15	49	43.1	6.15
80-84	136	102.4	8.78	73	148.6	17.39	63	75.3	9.48
85+	236	203.9	13.27	99	310.1	31.17	137	163.4	13.96
Total deaths	644			313			331		
Connecticut									
. Crude rate		11.2	-		11.1	-		11.2	-
. AAMR (1940 Std. mil.)		3.9	0.19		4.8	0.31		3.4	0.25
. AAMR (1970 Std. mil.)		6.2	0.26		8.0	0.46		5.2	0.32
. AAMR (2000 Std. mil.)		9.1	0.36		12.3	0.70		7.5	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.4	-		5.5	-		3.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----End stage renal disease (ESRD) (581-588,591,593.8,593.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	1.1	0.55	1	0.5	0.54	3	1.7	0.98
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.3	0.29	1	0.6	0.57	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	3	0.8	0.48	2	1.1	0.77	1	0.6	0.55
30-34	1	0.2	0.22	0	0.0	0.00	1	0.4	0.44
35-39	2	0.4	0.27	0	0.0	0.00	2	0.8	0.55
40-44	8	1.6	0.58	5	2.1	0.93	3	1.2	0.70
45-49	12	2.9	0.84	7	3.4	1.30	5	2.4	1.07
50-54	15	4.1	1.06	11	6.1	1.85	4	2.1	1.07
55-59	17	6.2	1.50	9	6.7	2.24	8	5.7	2.01
60-64	25	11.3	2.25	13	12.1	3.36	12	10.5	3.02
65-69	37	16.2	2.66	18	17.0	4.01	19	15.4	3.54
70-74	87	40.0	4.28	46	48.8	7.19	41	33.2	5.19
75-79	124	64.4	5.78	63	80.0	10.07	61	53.6	6.86
80-84	152	114.4	9.28	84	171.0	18.66	68	81.3	9.85
85+	264	228.1	14.04	115	360.2	33.59	149	177.7	14.56
Total deaths	752			375			377		
Connecticut									
. Crude rate		13.0	-		13.4	-		12.7	-
. AAMR (1940 Std. mil.)		4.7	0.21		6.0	0.35		3.9	0.27
. AAMR (1970 Std. mil.)		7.3	0.28		9.7	0.51		6.0	0.35
. AAMR (2000 Std. mil.)		10.7	0.39		14.7	0.77		8.6	0.45
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.5	-		6.8	-		4.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	25	6.9	1.38	17	9.1	2.21	8	4.5	1.60
5-9	11	2.8	0.84	10	5.0	1.57	1	0.5	0.52
10-14	22	5.9	1.25	10	5.2	1.65	12	6.5	1.89
15-19	91	26.6	2.79	61	34.5	4.42	30	18.1	3.31
20-24	98	32.1	3.25	73	46.6	5.45	25	16.9	3.37
25-29	114	31.3	2.93	91	49.6	5.20	23	12.7	2.66
30-34	160	34.9	2.76	121	52.8	4.80	39	17.0	2.73
35-39	157	30.4	2.43	120	46.6	4.26	37	14.3	2.36
40-44	130	26.6	2.34	100	41.6	4.16	30	12.1	2.21
45-49	121	29.3	2.66	88	43.2	4.61	33	15.8	2.75
50-54	64	17.5	2.18	47	26.2	3.82	17	9.1	2.21
55-59	54	19.7	2.68	30	22.4	4.09	24	17.1	3.48
60-64	58	26.1	3.43	36	33.6	5.60	22	19.2	4.09
65-69	87	38.0	4.07	60	56.7	7.32	27	21.9	4.22
70-74	108	49.6	4.77	58	61.5	8.07	50	40.5	5.73
75-79	151	78.4	6.38	81	102.8	11.42	70	61.5	7.35
80-84	198	149.1	10.59	94	191.3	19.74	104	124.3	12.19
85+	300	259.2	14.96	129	404.1	35.58	171	204.0	15.60
Total deaths	1,949			1,226			723		
Connecticut									
. Crude rate		33.8	-		43.7	-		24.4	-
. AAMR (1940 Std. mil.)		25.1	0.67		36.0	1.13		14.8	0.71
. AAMR (1970 Std. mil.)		26.3	0.65		37.3	1.12		16.5	0.72
. AAMR (2000 Std. mil.)		31.2	0.71		44.6	1.29		19.8	0.77
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		30.1	-		42.8	-		17.8	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Motor vehicle accidents (E810-E825)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.6	0.39	0	0.0	0.00	2	1.1	0.80
5-9	7	1.8	0.67	7	3.5	1.31	0	0.0	0.00
10-14	14	3.7	1.00	5	2.6	1.17	9	4.9	1.63
15-19	71	20.7	2.46	44	24.9	3.75	27	16.3	3.14
20-24	58	19.0	2.50	41	26.2	4.09	17	11.5	2.78
25-29	58	15.9	2.09	47	25.6	3.74	11	6.1	1.84
30-34	60	13.1	1.69	40	17.5	2.76	20	8.7	1.95
35-39	45	8.7	1.30	28	10.9	2.06	17	6.6	1.60
40-44	37	7.6	1.25	24	10.0	2.04	13	5.2	1.46
45-49	41	9.9	1.55	29	14.2	2.64	12	5.7	1.66
50-54	28	7.6	1.44	20	11.1	2.49	8	4.3	1.51
55-59	15	5.5	1.41	6	4.5	1.83	9	6.4	2.13
60-64	21	9.5	2.07	13	12.1	3.36	8	7.0	2.47
65-69	26	11.4	2.23	17	16.1	3.90	9	7.3	2.43
70-74	33	15.2	2.64	16	17.0	4.24	17	13.8	3.34
75-79	33	17.1	2.98	16	20.3	5.08	17	14.9	3.62
80-84	34	25.6	4.39	15	30.5	7.88	19	22.7	5.21
85+	23	19.9	4.14	9	28.2	9.40	14	16.7	4.46
Total deaths	606			377			229		
Connecticut									
. Crude rate		10.5	-		13.4	-		7.7	-
. AAMR (1940 Std. mil.)		10.3	0.46		13.5	0.74		7.0	0.54
. AAMR (1970 Std. mil.)		10.0	0.44		13.1	0.71		7.1	0.52
. AAMR (2000 Std. mil.)		10.4	0.43		13.6	0.70		7.4	0.50
. HP 2000 (1940 Std. mil.)		10.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		16.0	-		21.7	-		10.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.3	0.25	1	0.5	0.50	0	0.0	0.00
10-14	1	0.3	0.27	1	0.5	0.52	0	0.0	0.00
15-19	1	0.3	0.29	1	0.6	0.57	0	0.0	0.00
20-24	4	1.3	0.66	4	2.6	1.28	0	0.0	0.00
25-29	2	0.5	0.39	2	1.1	0.77	0	0.0	0.00
30-34	7	1.5	0.58	6	2.6	1.07	1	0.4	0.44
35-39	3	0.6	0.34	3	1.2	0.67	0	0.0	0.00
40-44	4	0.8	0.41	4	1.7	0.83	0	0.0	0.00
45-49	7	1.7	0.64	4	2.0	0.98	3	1.4	0.83
50-54	7	1.9	0.72	6	3.3	1.36	1	0.5	0.54
55-59	6	2.2	0.89	5	3.7	1.67	1	0.7	0.71
60-64	8	3.6	1.27	6	5.6	2.28	2	1.7	1.23
65-69	26	11.4	2.23	18	17.0	4.01	8	6.5	2.29
70-74	21	9.6	2.10	11	11.7	3.52	10	8.1	2.56
75-79	50	26.0	3.67	32	40.6	7.18	18	15.8	3.73
80-84	81	61.0	6.78	40	81.4	12.87	41	49.0	7.65
85+	160	138.2	10.93	64	200.5	25.06	96	114.5	11.69
Total deaths	389			208			181		
Connecticut									
. Crude rate		6.7	-		7.4	-		6.1	-
. AAMR (1940 Std. mil.)		2.4	0.16		3.7	0.30		1.4	0.14
. AAMR (1970 Std. mil.)		3.7	0.20		5.5	0.39		2.4	0.20
. AAMR (2000 Std. mil.)		5.5	0.28		8.1	0.57		3.8	0.29
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.6	-		3.6	-		1.7	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	1.7	0.67	5	2.7	1.20	1	0.6	0.57
5-9	1	0.3	0.25	1	0.5	0.50	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	3	0.9	0.51	3	1.7	0.98	0	0.0	0.00
20-24	4	1.3	0.66	4	2.6	1.28	0	0.0	0.00
25-29	6	1.6	0.67	5	2.7	1.22	1	0.6	0.55
30-34	4	0.9	0.44	4	1.7	0.87	0	0.0	0.00
35-39	5	1.0	0.43	5	1.9	0.87	0	0.0	0.00
40-44	2	0.4	0.29	2	0.8	0.59	0	0.0	0.00
45-49	2	0.5	0.34	2	1.0	0.69	0	0.0	0.00
50-54	2	0.5	0.39	2	1.1	0.79	0	0.0	0.00
55-59	1	0.4	0.36	1	0.7	0.75	0	0.0	0.00
60-64	2	0.9	0.64	0	0.0	0.00	2	1.7	1.23
65-69	1	0.4	0.44	1	0.9	0.95	0	0.0	0.00
70-74	1	0.5	0.46	1	1.1	1.06	0	0.0	0.00
75-79	5	2.6	1.16	3	3.8	2.20	2	1.8	1.24
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	2	1.7	1.22	1	3.1	3.13	1	1.2	1.19
Total deaths	47			40			7		
Connecticut									
. Crude rate		0.8	-		1.4	-		0.2	-
. AAMR (1940 Std. mil.)		0.8	0.13		1.5	0.24		-	-
. AAMR (1970 Std. mil.)		0.8	0.12		1.4	0.23		-	-
. AAMR (2000 Std. mil.)		0.8	0.12		1.4	0.23		-	-
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.6	-		2.5	-		0.4	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.6	0.39	1	0.5	0.54	1	0.6	0.57
5-9	1	0.3	0.25	0	0.0	0.00	1	0.5	0.52
10-14	2	0.5	0.38	2	1.0	0.74	0	0.0	0.00
15-19	1	0.3	0.29	0	0.0	0.00	1	0.6	0.60
20-24	3	1.0	0.57	3	1.9	1.11	0	0.0	0.00
25-29	1	0.3	0.27	1	0.5	0.55	0	0.0	0.00
30-34	1	0.2	0.22	0	0.0	0.00	1	0.4	0.44
35-39	3	0.6	0.34	3	1.2	0.67	0	0.0	0.00
40-44	4	0.8	0.41	2	0.8	0.59	2	0.8	0.57
45-49	4	1.0	0.48	1	0.5	0.49	3	1.4	0.83
50-54	3	0.8	0.47	1	0.6	0.56	2	1.1	0.76
55-59	2	0.7	0.51	0	0.0	0.00	2	1.4	1.01
60-64	3	1.4	0.78	2	1.9	1.32	1	0.9	0.87
65-69	4	1.7	0.87	3	2.8	1.64	1	0.8	0.81
70-74	3	1.4	0.80	0	0.0	0.00	3	2.4	1.40
75-79	5	2.6	1.16	1	1.3	1.27	4	3.5	1.76
80-84	5	3.8	1.68	4	8.1	4.07	1	1.2	1.19
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	47			24			23		
Connecticut									
. Crude rate		0.8	-		0.9	-		0.8	-
. AAMR (1940 Std. mil.)		0.7	0.11		0.8	0.17		0.6	0.14
. AAMR (1970 Std. mil.)		0.7	0.11		0.8	0.17		0.7	0.15
. AAMR (2000 Std. mil.)		0.8	0.11		0.9	0.18		0.7	0.15
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.9	-		1.4	-		0.9	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	1.7	0.67	3	1.6	0.93	3	1.7	0.98
5-9	3	0.8	0.44	2	1.0	0.70	1	0.5	0.52
10-14	4	1.1	0.53	2	1.0	0.74	2	1.1	0.77
15-19	17	5.0	1.20	14	7.9	2.12	3	1.8	1.05
20-24	22	7.2	1.54	21	13.4	2.92	1	0.7	0.67
25-29	12	3.3	0.95	9	4.9	1.64	3	1.7	0.96
30-34	8	1.7	0.62	6	2.6	1.07	2	0.9	0.62
35-39	12	2.3	0.67	7	2.7	1.03	5	1.9	0.87
40-44	15	3.1	0.79	11	4.6	1.38	4	1.6	0.81
45-49	5	1.2	0.54	4	2.0	0.98	1	0.5	0.48
50-54	9	2.5	0.82	8	4.5	1.57	1	0.5	0.54
55-59	3	1.1	0.63	1	0.7	0.75	2	1.4	1.01
60-64	6	2.7	1.10	4	3.7	1.87	2	1.7	1.23
65-69	3	1.3	0.76	2	1.9	1.34	1	0.8	0.81
70-74	2	0.9	0.65	2	2.1	1.50	0	0.0	0.00
75-79	2	1.0	0.73	1	1.3	1.27	1	0.9	0.88
80-84	2	1.5	1.06	1	2.0	2.04	1	1.2	1.19
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	131			98			33		
Connecticut									
. Crude rate		2.3	-		3.5	-		1.1	-
. AAMR (1940 Std. mil.)		2.6	0.24		4.0	0.42		1.2	0.22
. AAMR (1970 Std. mil.)		2.5	0.23		3.8	0.40		1.2	0.21
. AAMR (2000 Std. mil.)		2.4	0.21		3.6	0.37		1.1	0.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.1	-		12.5	-		3.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.6	0.39	0	0.0	0.00	2	1.1	0.80
5-9	6	1.5	0.62	6	3.0	1.21	0	0.0	0.00
10-14	14	3.7	1.00	5	2.6	1.17	9	4.9	1.63
15-19	70	20.4	2.44	43	24.3	3.71	27	16.3	3.14
20-24	55	18.0	2.43	38	24.3	3.93	17	11.5	2.78
25-29	55	15.1	2.04	45	24.5	3.66	10	5.5	1.75
30-34	60	13.1	1.69	39	17.0	2.73	21	9.2	2.00
35-39	43	8.3	1.27	27	10.5	2.02	16	6.2	1.55
40-44	36	7.4	1.23	23	9.6	2.00	13	5.2	1.46
45-49	41	9.9	1.55	29	14.2	2.64	12	5.7	1.66
50-54	28	7.6	1.44	20	11.1	2.49	8	4.3	1.51
55-59	15	5.5	1.41	6	4.5	1.83	9	6.4	2.13
60-64	20	9.0	2.02	13	12.1	3.36	7	6.1	2.31
65-69	27	11.8	2.27	18	17.0	4.01	9	7.3	2.43
70-74	32	14.7	2.60	17	18.0	4.37	15	12.2	3.14
75-79	33	17.1	2.98	16	20.3	5.08	17	14.9	3.62
80-84	33	24.8	4.33	14	28.5	7.62	19	22.7	5.21
85+	23	19.9	4.14	9	28.2	9.40	14	16.7	4.46
Total deaths	593			368			225		
Connecticut									
. Crude rate		10.3	-		13.1	-		7.6	-
. AAMR (1940 Std. mil.)		10.0	0.45		13.1	0.73		6.9	0.53
. AAMR (1970 Std. mil.)		9.8	0.43		12.7	0.70		7.0	0.52
. AAMR (2000 Std. mil.)		10.2	0.42		13.3	0.70		7.2	0.50
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.6	-		21.3	-		10.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.3	0.25	1	0.5	0.50	0	0.0	0.00
10-14	1	0.3	0.27	1	0.5	0.52	0	0.0	0.00
15-19	2	0.6	0.41	2	1.1	0.80	0	0.0	0.00
20-24	4	1.3	0.66	4	2.6	1.28	0	0.0	0.00
25-29	3	0.8	0.48	2	1.1	0.77	1	0.6	0.55
30-34	8	1.7	0.62	7	3.1	1.15	1	0.4	0.44
35-39	3	0.6	0.34	3	1.2	0.67	0	0.0	0.00
40-44	5	1.0	0.46	5	2.1	0.93	0	0.0	0.00
45-49	9	2.2	0.73	6	2.9	1.20	3	1.4	0.83
50-54	7	1.9	0.72	6	3.3	1.36	1	0.5	0.54
55-59	5	1.8	0.81	4	3.0	1.49	1	0.7	0.71
60-64	10	4.5	1.43	8	7.5	2.64	2	1.7	1.23
65-69	23	10.0	2.09	17	16.1	3.90	6	4.9	1.99
70-74	15	6.9	1.78	10	10.6	3.35	5	4.1	1.81
75-79	42	21.8	3.37	29	36.8	6.84	13	11.4	3.17
80-84	53	39.9	5.48	30	61.1	11.15	23	27.5	5.73
85+	88	76.0	8.10	36	112.8	18.79	52	62.0	8.60
Total deaths	279			171			108		
Connecticut									
. Crude rate		4.8	-		6.1	-		3.6	-
. AAMR (1940 Std. mil.)		2.1	0.16		3.5	0.31		1.0	0.13
. AAMR (1970 Std. mil.)		2.9	0.19		4.7	0.37		1.5	0.17
. AAMR (2000 Std. mil.)		4.1	0.24		6.5	0.50		2.4	0.23
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.5	-		3.6	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	15	4.4	1.13	14	7.9	2.12	1	0.6	0.60
20-24	27	8.9	1.70	25	16.0	3.19	2	1.3	0.95
25-29	23	6.3	1.32	22	12.0	2.56	1	0.6	0.55
30-34	29	6.3	1.18	28	12.2	2.31	1	0.4	0.44
35-39	33	6.4	1.11	26	10.1	1.98	7	2.7	1.02
40-44	29	5.9	1.10	23	9.6	2.00	6	2.4	0.99
45-49	11	2.7	0.80	8	3.9	1.39	3	1.4	0.83
50-54	21	5.7	1.25	19	10.6	2.43	2	1.1	0.76
55-59	13	4.7	1.31	11	8.2	2.48	2	1.4	1.01
60-64	14	6.3	1.69	13	12.1	3.36	1	0.9	0.87
65-69	9	3.9	1.31	8	7.6	2.67	1	0.8	0.81
70-74	7	3.2	1.22	7	7.4	2.80	0	0.0	0.00
75-79	5	2.6	1.16	5	6.3	2.84	0	0.0	0.00
80-84	10	7.5	2.38	10	20.4	6.44	0	0.0	0.00
85+	7	6.0	2.29	6	18.8	7.67	1	1.2	1.19
Total deaths	253			225			28		
Connecticut									
. Crude rate		4.4	-		8.0	-		0.9	-
. AAMR (1940 Std. mil.)		4.3	0.29		7.7	0.55		0.9	0.18
. AAMR (1970 Std. mil.)		4.0	0.27		7.3	0.51		0.8	0.16
. AAMR (2000 Std. mil.)		4.3	0.27		8.0	0.54		0.9	0.18
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.2	-		21.1	-		3.3	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	5	1.5	0.65	5	2.8	1.27	0	0.0	0.00
20-24	8	2.6	0.93	7	4.5	1.69	1	0.7	0.67
25-29	13	3.6	0.99	13	7.1	1.97	0	0.0	0.00
30-34	26	5.7	1.11	25	10.9	2.18	1	0.4	0.44
35-39	25	4.8	0.97	21	8.2	1.78	4	1.5	0.77
40-44	22	4.5	0.96	17	7.1	1.72	5	2.0	0.90
45-49	9	2.2	0.73	6	2.9	1.20	3	1.4	0.83
50-54	15	4.1	1.06	13	7.2	2.01	2	1.1	0.76
55-59	10	3.6	1.15	9	6.7	2.24	1	0.7	0.71
60-64	11	5.0	1.49	11	10.3	3.09	0	0.0	0.00
65-69	8	3.5	1.23	8	7.6	2.67	0	0.0	0.00
70-74	7	3.2	1.22	7	7.4	2.80	0	0.0	0.00
75-79	5	2.6	1.16	5	6.3	2.84	0	0.0	0.00
80-84	10	7.5	2.38	10	20.4	6.44	0	0.0	0.00
85+	7	6.0	2.29	6	18.8	7.67	1	1.2	1.19
Total deaths	181			163			18		
Connecticut									
. Crude rate		3.1	-		5.8	-		0.6	-
. AAMR (1940 Std. mil.)		2.7	0.22		5.0	0.42		0.5	0.13
. AAMR (1970 Std. mil.)		2.6	0.20		4.9	0.40		0.5	0.12
. AAMR (2000 Std. mil.)		3.0	0.22		5.7	0.45		0.6	0.14
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.9	-		10.7	-		1.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	9	2.6	0.88	8	4.5	1.60	1	0.6	0.60
20-24	19	6.2	1.43	18	11.5	2.71	1	0.7	0.67
25-29	9	2.5	0.82	8	4.4	1.54	1	0.6	0.55
30-34	2	0.4	0.31	2	0.9	0.62	0	0.0	0.00
35-39	7	1.4	0.51	5	1.9	0.87	2	0.8	0.55
40-44	6	1.2	0.50	5	2.1	0.93	1	0.4	0.40
45-49	2	0.5	0.34	2	1.0	0.69	0	0.0	0.00
50-54	6	1.6	0.67	6	3.3	1.36	0	0.0	0.00
55-59	1	0.4	0.36	0	0.0	0.00	1	0.7	0.71
60-64	2	0.9	0.64	1	0.9	0.93	1	0.9	0.87
65-69	1	0.4	0.44	0	0.0	0.00	1	0.8	0.81
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	64			55			9		
Connecticut									
. Crude rate		1.1	-		2.0	-		0.3	-
. AAMR (1940 Std. mil.)		1.4	0.18		2.4	0.34		-	-
. AAMR (1970 Std. mil.)		1.3	0.17		2.2	0.31		-	-
. AAMR (2000 Std. mil.)		1.2	0.15		2.1	0.28		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.5	-		9.3	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	15	3.3	0.85	14	6.1	1.63	1	0.4	0.44
35-39	18	3.5	0.82	13	5.0	1.40	5	1.9	0.87
40-44	31	6.4	1.14	22	9.2	1.95	9	3.6	1.21
45-49	41	9.9	1.55	33	16.2	2.82	8	3.8	1.35
50-54	39	10.6	1.70	28	15.6	2.95	11	5.9	1.78
55-59	25	9.1	1.82	17	12.7	3.08	8	5.7	2.01
60-64	31	14.0	2.51	26	24.2	4.76	5	4.4	1.95
65-69	28	12.2	2.31	17	16.1	3.90	11	8.9	2.69
70-74	17	7.8	1.89	11	11.7	3.52	6	4.9	1.98
75-79	12	6.2	1.80	8	10.2	3.59	4	3.5	1.76
80-84	10	7.5	2.38	8	16.3	5.76	2	2.4	1.69
85+	2	1.7	1.22	2	6.3	4.43	0	0.0	0.00
Total deaths	269			199			70		
Connecticut									
. Crude rate		4.7	-		7.1	-		2.4	-
. AAMR (1940 Std. mil.)		3.7	0.23		5.7	0.41		1.8	0.22
. AAMR (1970 Std. mil.)		3.8	0.24		5.9	0.43		1.9	0.23
. AAMR (2000 Std. mil.)		4.4	0.27		6.9	0.49		2.2	0.26
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.3	-		9.9	-		3.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.3	0.28	0	0.0	0.00	1	0.6	0.57
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	9	2.6	0.88	6	3.4	1.39	3	1.8	1.05
20-24	29	9.5	1.77	22	14.0	2.99	7	4.7	1.78
25-29	43	11.8	1.80	33	18.0	3.13	10	5.5	1.75
30-34	89	19.4	2.06	66	28.8	3.55	23	10.1	2.10
35-39	107	20.8	2.01	83	32.2	3.54	24	9.3	1.90
40-44	95	19.5	2.00	70	29.1	3.48	25	10.1	2.02
45-49	62	15.0	1.91	41	20.1	3.14	21	10.0	2.19
50-54	18	4.9	1.16	11	6.1	1.85	7	3.7	1.42
55-59	13	4.7	1.31	6	4.5	1.83	7	5.0	1.88
60-64	4	1.8	0.90	0	0.0	0.00	4	3.5	1.74
65-69	3	1.3	0.76	0	0.0	0.00	3	2.4	1.41
70-74	8	3.7	1.30	5	5.3	2.37	3	2.4	1.40
75-79	7	3.6	1.37	4	5.1	2.54	3	2.6	1.52
80-84	7	5.3	1.99	2	4.1	2.88	5	6.0	2.67
85+	4	3.5	1.73	1	3.1	3.13	3	3.6	2.07
Total deaths	499			350			149		
Connecticut									
. Crude rate		8.7	-		12.5	-		5.0	-
. AAMR (1940 Std. mil.)		8.1	0.38		11.6	0.64		4.6	0.40
. AAMR (1970 Std. mil.)		7.0	0.33		9.8	0.55		4.1	0.36
. AAMR (2000 Std. mil.)		8.4	0.38		11.9	0.64		4.9	0.40
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.6	-		7.8	-		3.3	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is higher  
. THE U.S. HP 2000 target: CT rate is higher