

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Infectious diseases and parasitic disease (1-139)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	3.3	1.33	5	5.3	2.36	1	1.1	1.12
5-9	1	0.5	0.51	0	0.0	0.00	1	1.0	1.04
10-14	1	0.5	0.54	0	0.0	0.00	1	1.1	1.10
15-19	1	0.6	0.59	0	0.0	0.00	1	1.2	1.22
20-24	3	2.0	1.13	1	1.3	1.26	2	2.7	1.90
25-29	10	5.4	1.70	7	7.4	2.81	3	3.3	1.89
30-34	24	10.3	2.10	17	14.5	3.53	7	6.0	2.28
35-39	41	15.8	2.47	31	24.0	4.30	10	7.7	2.44
40-44	38	15.8	2.56	30	25.2	4.60	8	6.5	2.31
45-49	28	13.6	2.57	20	19.7	4.40	8	7.7	2.71
50-54	21	11.5	2.52	14	15.7	4.19	7	7.5	2.85
55-59	21	15.8	3.45	12	18.5	5.34	9	13.2	4.41
60-64	26	23.4	4.59	18	33.6	7.91	8	13.9	4.93
65-69	42	35.9	5.54	18	33.4	7.86	24	38.1	7.78
70-74	41	37.5	5.85	22	46.5	9.92	19	30.6	7.01
75-79	60	62.2	8.03	35	89.2	15.07	25	43.7	8.74
80-84	94	144.0	14.85	43	179.0	27.30	51	123.6	17.31
85+	135	237.9	20.47	38	248.1	40.25	97	234.1	23.77
Total deaths	593			311			282		
Connecticut									
. Crude rate		20.5	-		22.1	-		19.1	-
. AAMR (1940 Std. mil.)		11.0	0.54		14.5	0.89		7.9	0.63
. AAMR (1970 Std. mil.)		13.3	0.58		17.0	0.98		10.2	0.69
. AAMR (2000 Std. mil.)		17.9	0.74		22.6	1.30		13.8	0.85
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.5	-		18.1	-		9.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Septicemia (038)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.59	0	0.0	0.00	1	1.2	1.22
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
30-34	1	0.4	0.43	1	0.9	0.86	0	0.0	0.00
35-39	3	1.2	0.67	1	0.8	0.77	2	1.5	1.09
40-44	6	2.5	1.02	4	3.4	1.68	2	1.6	1.16
45-49	4	1.9	0.97	4	3.9	1.97	0	0.0	0.00
50-54	6	3.3	1.34	3	3.4	1.94	3	3.2	1.87
55-59	9	6.8	2.26	3	4.6	2.67	6	8.8	3.60
60-64	12	10.8	3.12	7	13.1	4.93	5	8.7	3.89
65-69	25	21.4	4.28	11	20.4	6.15	14	22.2	5.94
70-74	24	21.9	4.48	15	31.7	8.19	9	14.5	4.83
75-79	42	43.5	6.72	21	53.5	11.67	21	36.7	8.01
80-84	77	118.0	13.44	30	124.9	22.80	47	114.0	16.62
85+	124	218.5	19.62	35	228.5	38.63	89	214.8	22.77
Total deaths	336			137			199		
Connecticut									
. Crude rate		11.6	-		9.7	-		13.4	-
. AAMR (1940 Std. mil.)		4.3	0.29		4.8	0.46		3.8	0.38
. AAMR (1970 Std. mil.)		6.6	0.38		7.3	0.63		6.0	0.48
. AAMR (2000 Std. mil.)		9.6	0.53		10.7	0.93		8.9	0.64
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.1	-		4.7	-		3.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----HIV infection (042-044)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.51	0	0.0	0.00	1	1.0	1.04
10-14	1	0.5	0.54	0	0.0	0.00	1	1.1	1.10
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.7	0.65	0	0.0	0.00	1	1.3	1.34
25-29	9	4.8	1.61	6	6.4	2.60	3	3.3	1.89
30-34	21	9.0	1.97	15	12.8	3.31	6	5.2	2.11
35-39	29	11.2	2.08	23	17.8	3.71	6	4.6	1.89
40-44	23	9.5	1.99	20	16.8	3.76	3	2.5	1.42
45-49	18	8.7	2.06	13	12.8	3.55	5	4.8	2.15
50-54	5	2.7	1.23	5	5.6	2.50	0	0.0	0.00
55-59	5	3.8	1.68	4	6.2	3.09	1	1.5	1.47
60-64	4	3.6	1.80	4	7.5	3.73	0	0.0	0.00
65-69	3	2.6	1.48	3	5.6	3.21	0	0.0	0.00
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	120			93			27		
Connecticut									
. Crude rate		4.2	-		6.6	-		1.8	-
. AAMR (1940 Std. mil.)		3.8	0.35		5.8	0.60		1.8	0.36
. AAMR (1970 Std. mil.)		3.3	0.31		5.0	0.53		1.6	0.32
. AAMR (2000 Std. mil.)		4.0	0.36		6.2	0.64		1.8	0.35
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.7	-		9.1	-		2.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.1	0.77	0	0.0	0.00	2	2.2	1.59
5-9	2	1.0	0.72	1	1.0	0.99	1	1.0	1.04
10-14	4	2.2	1.08	3	3.2	1.83	1	1.1	1.10
15-19	4	2.4	1.18	2	2.3	1.61	2	2.4	1.73
20-24	13	8.5	2.35	9	11.4	3.79	4	5.4	2.69
25-29	11	5.9	1.78	7	7.4	2.81	4	4.4	2.18
30-34	30	12.9	2.35	9	7.7	2.57	21	18.1	3.95
35-39	65	25.1	3.11	25	19.3	3.86	40	30.9	4.88
40-44	118	48.9	4.50	53	44.6	6.12	65	53.2	6.60
45-49	169	82.1	6.32	69	67.9	8.18	100	96.0	9.60
50-54	302	165.8	9.54	129	144.3	12.70	173	186.5	14.18
55-59	361	271.7	14.30	190	293.1	21.26	171	251.2	19.21
60-64	533	480.0	20.79	281	524.0	31.26	252	438.9	27.65
65-69	780	667.0	23.88	430	796.9	38.43	350	555.7	29.70
70-74	999	913.1	28.89	519	1098.1	48.20	480	772.4	35.26
75-79	1,139	1180.8	34.99	579	1474.9	61.29	560	979.0	41.37
80-84	1,005	1539.8	48.57	487	2027.1	91.86	518	1255.9	55.18
85+	1,077	1897.7	57.83	431	2814.4	135.57	646	1559.0	61.34
Total deaths	6,614			3,224			3,390		
Connecticut									
. Crude rate		229.2	-		229.3	-		229.1	-
. AAMR (1940 Std. mil.)		116.3	1.65		132.1	2.54		105.8	2.19
. AAMR (1970 Std. mil.)		155.4	2.02		183.1	3.29		138.0	2.60
. AAMR (2000 Std. mil.)		197.1	2.44		237.8	4.24		173.7	3.05
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		125.6	-		150.4	-		107.4	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	1.1	0.76	1	1.1	1.06	1	1.1	1.09
30-34	2	0.9	0.61	0	0.0	0.00	2	1.7	1.22
35-39	8	3.1	1.09	5	3.9	1.73	3	2.3	1.34
40-44	13	5.4	1.50	9	7.6	2.52	4	3.3	1.64
45-49	12	5.8	1.68	6	5.9	2.41	6	5.8	2.35
50-54	24	13.2	2.69	12	13.4	3.87	12	12.9	3.73
55-59	30	22.6	4.12	17	26.2	6.36	13	19.1	5.30
60-64	45	40.5	6.04	26	48.5	9.51	19	33.1	7.59
65-69	74	63.3	7.36	39	72.3	11.57	35	55.6	9.39
70-74	101	92.3	9.19	52	110.0	15.26	49	78.9	11.26
75-79	117	121.3	11.21	63	160.5	20.22	54	94.4	12.85
80-84	117	179.3	16.57	55	228.9	30.87	62	150.3	19.09
85+	151	266.1	21.65	46	300.4	44.29	105	253.4	24.73
Total deaths	696			331			365		
Connecticut									
. Crude rate		24.1	-		23.5	-		24.7	-
. AAMR (1940 Std. mil.)		11.1	0.49		13.1	0.79		9.5	0.63
. AAMR (1970 Std. mil.)		15.4	0.62		18.4	1.03		13.3	0.77
. AAMR (2000 Std. mil.)		20.4	0.78		24.5	1.36		17.7	0.95
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.1	-		14.6	-		10.1	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colon cancer (153)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
30-34	1	0.4	0.43	0	0.0	0.00	1	0.9	0.86
35-39	6	2.3	0.95	3	2.3	1.34	3	2.3	1.34
40-44	10	4.1	1.31	7	5.9	2.22	3	2.5	1.42
45-49	11	5.3	1.61	5	4.9	2.20	6	5.8	2.35
50-54	21	11.5	2.52	10	11.2	3.54	11	11.9	3.58
55-59	21	15.8	3.45	12	18.5	5.34	9	13.2	4.41
60-64	42	37.8	5.84	24	44.8	9.14	18	31.4	7.39
65-69	67	57.3	7.00	35	64.9	10.96	32	50.8	8.98
70-74	75	68.6	7.92	35	74.1	12.52	40	64.4	10.18
75-79	97	100.6	10.21	52	132.5	18.37	45	78.7	11.73
80-84	98	150.1	15.17	45	187.3	27.92	53	128.5	17.65
85+	125	220.3	19.70	36	235.1	39.18	89	214.8	22.77
Total deaths	575			265			310		
Connecticut									
. Crude rate		19.9	-		18.8	-		20.9	-
. AAMR (1940 Std. mil.)		9.2	0.45		10.6	0.71		8.1	0.58
. AAMR (1970 Std. mil.)		12.8	0.56		14.8	0.93		11.3	0.71
. AAMR (2000 Std. mil.)		16.9	0.71		19.6	1.22		15.0	0.87
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.1	-		12.1	-		8.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Rectal cancer (154)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.5	0.54	0	0.0	0.00	1	1.1	1.09
30-34	1	0.4	0.43	0	0.0	0.00	1	0.9	0.86
35-39	2	0.8	0.55	2	1.5	1.09	0	0.0	0.00
40-44	3	1.2	0.72	2	1.7	1.19	1	0.8	0.82
45-49	1	0.5	0.49	1	1.0	0.98	0	0.0	0.00
50-54	3	1.6	0.95	2	2.2	1.58	1	1.1	1.08
55-59	9	6.8	2.26	5	7.7	3.45	4	5.9	2.94
60-64	3	2.7	1.56	2	3.7	2.64	1	1.7	1.74
65-69	7	6.0	2.26	4	7.4	3.71	3	4.8	2.75
70-74	21	19.2	4.19	16	33.9	8.46	5	8.0	3.60
75-79	18	18.7	4.40	9	22.9	7.64	9	15.7	5.24
80-84	17	26.0	6.32	8	33.3	11.77	9	21.8	7.27
85+	22	38.8	8.26	7	45.7	17.28	15	36.2	9.35
Total deaths	108			58			50		
Connecticut									
. Crude rate		3.7	-		4.1	-		3.4	-
. AAMR (1940 Std. mil.)		1.8	0.20		2.3	0.33		1.3	0.24
. AAMR (1970 Std. mil.)		2.4	0.24		3.2	0.43		1.8	0.28
. AAMR (2000 Std. mil.)		3.2	0.31		4.2	0.56		2.4	0.35
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.3	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.8	0.55	0	0.0	0.00	2	1.5	1.09
40-44	4	1.7	0.83	3	2.5	1.46	1	0.8	0.82
45-49	8	3.9	1.37	2	2.0	1.39	6	5.8	2.35
50-54	17	9.3	2.26	8	8.9	3.16	9	9.7	3.23
55-59	23	17.3	3.61	15	23.1	5.97	8	11.8	4.16
60-64	27	24.3	4.68	19	35.4	8.13	8	13.9	4.93
65-69	52	44.5	6.17	30	55.6	10.15	22	34.9	7.45
70-74	59	53.9	7.02	26	55.0	10.79	33	53.1	9.24
75-79	71	73.6	8.74	29	73.9	13.72	42	73.4	11.33
80-84	62	95.0	12.06	26	108.2	21.22	36	87.3	14.55
85+	42	74.0	11.42	10	65.3	20.65	32	77.2	13.65
Total deaths	367			168			199		
Connecticut									
. Crude rate		12.7	-		11.9	-		13.4	-
. AAMR (1940 Std. mil.)		6.5	0.38		7.4	0.61		5.7	0.48
. AAMR (1970 Std. mil.)		8.7	0.48		9.9	0.78		7.7	0.59
. AAMR (2000 Std. mil.)		10.9	0.57		12.0	0.93		9.9	0.72
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.2	-		7.3	-		5.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	4	1.5	0.77	2	1.5	1.09	2	1.5	1.09
40-44	4	1.7	0.83	0	0.0	0.00	4	3.3	1.64
45-49	5	2.4	1.09	5	4.9	2.20	0	0.0	0.00
50-54	4	2.2	1.10	2	2.2	1.58	2	2.2	1.52
55-59	7	5.3	1.99	2	3.1	2.18	5	7.3	3.29
60-64	7	6.3	2.38	5	9.3	4.17	2	3.5	2.46
65-69	7	6.0	2.26	4	7.4	3.71	3	4.8	2.75
70-74	10	9.1	2.89	6	12.7	5.18	4	6.4	3.22
75-79	17	17.6	4.27	8	20.4	7.20	9	15.7	5.24
80-84	10	15.3	4.84	4	16.7	8.33	6	14.5	5.94
85+	15	26.4	6.82	4	26.1	13.06	11	26.5	8.00
Total deaths	90			42			48		
Connecticut									
. Crude rate		3.1	-		3.0	-		3.2	-
. AAMR (1940 Std. mil.)		1.7	0.20		1.9	0.31		1.5	0.26
. AAMR (1970 Std. mil.)		2.1	0.24		2.4	0.38		1.9	0.30
. AAMR (2000 Std. mil.)		2.7	0.29		3.0	0.47		2.5	0.37
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.7	-		2.6	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	3	1.3	0.74	-	-	-	3	2.6	1.49
35-39	7	2.7	1.02	-	-	-	7	5.4	2.04
40-44	27	11.2	2.15	-	-	-	27	22.1	4.25
45-49	34	16.5	2.83	-	-	-	34	32.6	5.60
50-54	48	26.4	3.80	-	-	-	48	51.7	7.47
55-59	31	23.3	4.19	-	-	-	31	45.5	8.18
60-64	43	38.7	5.91	-	-	-	43	74.9	11.42
65-69	57	48.7	6.46	-	-	-	57	90.5	11.99
70-74	73	66.7	7.81	-	-	-	73	117.5	13.75
75-79	67	69.5	8.49	-	-	-	67	117.1	14.31
80-84	62	95.0	12.06	-	-	-	62	150.3	19.09
85+	95	167.4	17.17	-	-	-	95	229.3	23.52
Total deaths	547			-			547		
Connecticut									
. Crude rate		-	-		-	-		37.0	-
. AAMR (1940 Std. mil.)		-	-		-	-		19.4	0.97
. AAMR (1970 Std. mil.)		-	-		-	-		23.8	1.10
. AAMR (2000 Std. mil.)		-	-		-	-		29.4	1.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		19.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is lower
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	1	0.5	0.54	-	-	-	1	1.1	1.09
30-34	2	0.9	0.61	-	-	-	2	1.7	1.22
35-39	3	1.2	0.67	-	-	-	3	2.3	1.34
40-44	1	0.4	0.41	-	-	-	1	0.8	0.82
45-49	5	2.4	1.09	-	-	-	5	4.8	2.15
50-54	2	1.1	0.78	-	-	-	2	2.2	1.52
55-59	4	3.0	1.50	-	-	-	4	5.9	2.94
60-64	1	0.9	0.90	-	-	-	1	1.7	1.74
65-69	2	1.7	1.21	-	-	-	2	3.2	2.25
70-74	1	0.9	0.91	-	-	-	1	1.6	1.61
75-79	3	3.1	1.80	-	-	-	3	5.2	3.03
80-84	0	0.0	0.00	-	-	-	0	0.0	0.00
85+	4	7.0	3.52	-	-	-	4	9.7	4.83
Total deaths	29			-			29		
Connecticut									
. Crude rate		-	-		-	-		2.0	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.4	0.29
. AAMR (1970 Std. mil.)		-	-		-	-		1.4	0.28
. AAMR (2000 Std. mil.)		-	-		-	-		1.7	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	0	0.0	0.00	-	-	-	0	0.0	0.00
45-49	1	0.5	0.49	-	-	-	1	1.0	0.96
50-54	0	0.0	0.00	-	-	-	0	0.0	0.00
55-59	1	0.8	0.75	-	-	-	1	1.5	1.47
60-64	2	1.8	1.27	-	-	-	2	3.5	2.46
65-69	3	2.6	1.48	-	-	-	3	4.8	2.75
70-74	6	5.5	2.24	-	-	-	6	9.7	3.94
75-79	9	9.3	3.11	-	-	-	9	15.7	5.24
80-84	4	6.1	3.06	-	-	-	4	9.7	4.85
85+	9	15.9	5.29	-	-	-	9	21.7	7.24
Total deaths	35			-			35		
Connecticut									
. Crude rate		-	-		-	-		2.4	-
. AAMR (1940 Std. mil.)		-	-		-	-		0.9	0.18
. AAMR (1970 Std. mil.)		-	-		-	-		1.3	0.23
. AAMR (2000 Std. mil.)		-	-		-	-		1.7	0.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	1	0.5	0.54	-	-	-	1	1.1	1.09
30-34	2	0.9	0.61	-	-	-	2	1.7	1.22
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	3	1.2	0.72	-	-	-	3	2.5	1.42
45-49	5	2.4	1.09	-	-	-	5	4.8	2.15
50-54	17	9.3	2.26	-	-	-	17	18.3	4.44
55-59	9	6.8	2.26	-	-	-	9	13.2	4.41
60-64	14	12.6	3.37	-	-	-	14	24.4	6.52
65-69	21	18.0	3.92	-	-	-	21	33.3	7.28
70-74	25	22.9	4.57	-	-	-	25	40.2	8.05
75-79	24	24.9	5.08	-	-	-	24	42.0	8.56
80-84	22	33.7	7.19	-	-	-	22	53.3	11.37
85+	27	47.6	9.16	-	-	-	27	65.2	12.54
Total deaths	170			-			170		
Connecticut									
. Crude rate		-	-		-	-		11.5	-
. AAMR (1940 Std. mil.)		-	-		-	-		5.9	0.53
. AAMR (1970 Std. mil.)		-	-		-	-		7.4	0.61
. AAMR (2000 Std. mil.)		-	-		-	-		9.0	0.71
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		5.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Bladder cancer (188)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.4	0.41	1	0.8	0.84	0	0.0	0.00
45-49	2	1.0	0.69	1	1.0	0.98	1	1.0	0.96
50-54	3	1.6	0.95	2	2.2	1.58	1	1.1	1.08
55-59	6	4.5	1.84	5	7.7	3.45	1	1.5	1.47
60-64	10	9.0	2.85	8	14.9	5.27	2	3.5	2.46
65-69	14	12.0	3.20	11	20.4	6.15	3	4.8	2.75
70-74	17	15.5	3.77	13	27.5	7.63	4	6.4	3.22
75-79	39	40.4	6.47	27	68.8	13.24	12	21.0	6.06
80-84	41	62.8	9.81	21	87.4	19.07	20	48.5	10.84
85+	47	82.8	12.08	28	182.8	34.55	19	45.9	10.52
Total deaths	180			117			63		
Connecticut									
. Crude rate		6.2	-		8.3	-		4.3	-
. AAMR (1940 Std. mil.)		2.4	0.22		4.0	0.41		1.2	0.20
. AAMR (1970 Std. mil.)		3.7	0.29		6.3	0.59		2.0	0.27
. AAMR (2000 Std. mil.)		5.2	0.39		9.1	0.85		2.9	0.37
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.5	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Brain & other CNS cancer (191-192)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.1	0.77	0	0.0	0.00	2	2.2	1.59
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	1.1	0.76	2	2.1	1.50	0	0.0	0.00
30-34	3	1.3	0.74	3	2.6	1.48	0	0.0	0.00
35-39	6	2.3	0.95	4	3.1	1.55	2	1.5	1.09
40-44	1	0.4	0.41	0	0.0	0.00	1	0.8	0.82
45-49	10	4.9	1.54	5	4.9	2.20	5	4.8	2.15
50-54	8	4.4	1.55	6	6.7	2.74	2	2.2	1.52
55-59	14	10.5	2.82	5	7.7	3.45	9	13.2	4.41
60-64	9	8.1	2.70	7	13.1	4.93	2	3.5	2.46
65-69	16	13.7	3.42	9	16.7	5.56	7	11.1	4.20
70-74	19	17.4	3.98	10	21.2	6.69	9	14.5	4.83
75-79	20	20.7	4.64	8	20.4	7.20	12	21.0	6.06
80-84	19	29.1	6.68	10	41.6	13.16	9	21.8	7.27
85+	13	22.9	6.35	4	26.1	13.06	9	21.7	7.24
Total deaths	143			74			69		
Connecticut									
. Crude rate		5.0	-		5.3	-		4.7	-
. AAMR (1940 Std. mil.)		3.0	0.28		3.6	0.45		2.5	0.36
. AAMR (1970 Std. mil.)		3.7	0.32		4.3	0.52		3.1	0.41
. AAMR (2000 Std. mil.)		4.4	0.37		5.2	0.61		3.7	0.46
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.5	-		4.1	-		2.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.51	0	0.0	0.00	1	1.0	1.04
10-14	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
15-19	3	1.8	1.02	1	1.1	1.14	2	2.4	1.73
20-24	5	3.3	1.46	4	5.1	2.53	1	1.3	1.34
25-29	3	1.6	0.93	3	3.2	1.84	0	0.0	0.00
30-34	3	1.3	0.74	1	0.9	0.86	2	1.7	1.22
35-39	3	1.2	0.67	1	0.8	0.77	2	1.5	1.09
40-44	4	1.7	0.83	2	1.7	1.19	2	1.6	1.16
45-49	3	1.5	0.84	1	1.0	0.98	2	1.9	1.36
50-54	11	6.0	1.82	3	3.4	1.94	8	8.6	3.05
55-59	10	7.5	2.38	8	12.3	4.36	2	2.9	2.08
60-64	12	10.8	3.12	7	13.1	4.93	5	8.7	3.89
65-69	36	30.8	5.13	28	51.9	9.81	8	12.7	4.49
70-74	32	29.2	5.17	14	29.6	7.92	18	29.0	6.83
75-79	44	45.6	6.88	24	61.1	12.48	20	35.0	7.82
80-84	37	56.7	9.32	21	87.4	19.07	16	38.8	9.70
85+	50	88.1	12.46	19	124.1	28.46	31	74.8	13.44
Total deaths	258			138			120		
Connecticut									
. Crude rate		8.9	-		9.8	-		8.1	-
. AAMR (1940 Std. mil.)		4.7	0.35		6.0	0.58		3.6	0.42
. AAMR (1970 Std. mil.)		6.1	0.41		8.1	0.71		4.7	0.48
. AAMR (2000 Std. mil.)		7.7	0.48		10.2	0.88		6.0	0.57
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.8	-		6.1	-		3.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	1.1	0.76	0	0.0	0.00	2	2.2	1.54
30-34	5	2.1	0.96	2	1.7	1.21	3	2.6	1.49
35-39	14	5.4	1.45	10	7.7	2.44	4	3.1	1.54
40-44	19	7.9	1.81	16	13.5	3.36	3	2.5	1.42
45-49	35	17.0	2.88	21	20.7	4.51	14	13.4	3.59
50-54	69	37.9	4.56	39	43.6	6.99	30	32.3	5.90
55-59	97	73.0	7.41	59	91.0	11.85	38	55.8	9.06
60-64	126	113.5	10.11	79	147.3	16.58	47	81.9	11.94
65-69	206	176.2	12.27	116	215.0	19.96	90	142.9	15.06
70-74	293	267.8	15.65	155	328.0	26.34	138	222.1	18.90
75-79	421	436.5	21.27	228	580.8	38.46	193	337.4	24.29
80-84	430	658.8	31.77	220	915.8	61.74	210	509.1	35.13
85+	498	877.5	39.32	181	1181.9	87.85	317	765.0	42.97
Total deaths	2,215			1,126			1,089		
Connecticut									
. Crude rate		76.8	-		80.1	-		73.6	-
. AAMR (1940 Std. mil.)		33.2	0.83		42.5	1.39		26.2	1.01
. AAMR (1970 Std. mil.)		47.8	1.08		61.9	1.88		38.0	1.28
. AAMR (2000 Std. mil.)		64.5	1.38		84.3	2.55		51.7	1.60
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.39	1	0.8	0.77	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	1	0.8	0.75	0	0.0	0.00	1	1.5	1.47
60-64	1	0.9	0.90	1	1.9	1.86	0	0.0	0.00
65-69	1	0.9	0.86	1	1.9	1.85	0	0.0	0.00
70-74	1	0.9	0.91	0	0.0	0.00	1	1.6	1.61
75-79	1	1.0	1.04	1	2.5	2.55	0	0.0	0.00
80-84	2	3.1	2.17	2	8.3	5.89	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	8			6			2		
Connecticut									
. Crude rate		0.3	-		0.4	-		0.1	-
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. AAMR (1970 Std. mil.)		-	-		-	-		-	-
. AAMR (2000 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.5	-		0.5	-		0.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.59	1	1.1	1.14	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	5	2.7	1.20	3	3.2	1.84	2	2.2	1.54
30-34	5	2.1	0.96	3	2.6	1.48	2	1.7	1.22
35-39	22	8.5	1.81	20	15.5	3.46	2	1.5	1.09
40-44	61	25.3	3.24	52	43.7	6.06	9	7.4	2.45
45-49	97	47.1	4.79	81	79.8	8.86	16	15.4	3.84
50-54	155	85.1	6.83	125	139.8	12.51	30	32.3	5.90
55-59	215	161.8	11.03	149	229.8	18.83	66	97.0	11.94
60-64	266	239.6	14.69	187	348.7	25.50	79	137.6	15.48
65-69	467	399.3	18.48	315	583.8	32.89	152	241.3	19.57
70-74	630	575.8	22.94	405	856.9	42.58	225	362.1	24.14
75-79	1,019	1056.4	33.09	593	1510.5	62.03	426	744.8	36.08
80-84	1,213	1858.4	53.36	617	2568.3	103.39	596	1445.0	59.19
85+	2,482	4373.4	87.78	705	4603.6	173.38	1,777	4288.3	101.73
Total deaths	6,638			3,256			3,382		
Connecticut									
. Crude rate		230.0	-		231.6	-		228.5	-
. AAMR (1940 Std. mil.)		84.0	1.27		118.6	2.30		55.8	1.31
. AAMR (1970 Std. mil.)		130.4	1.71		177.0	3.15		94.2	1.82
. AAMR (2000 Std. mil.)		190.5	2.35		249.7	4.44		146.3	2.56
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		100.4	-		137.4	-		70.0	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	0	0.0	0.00	1	1.1	1.12
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	3	1.5	0.84	2	2.0	1.39	1	1.0	0.96
50-54	2	1.1	0.78	1	1.1	1.12	1	1.1	1.08
55-59	6	4.5	1.84	3	4.6	2.67	3	4.4	2.54
60-64	3	2.7	1.56	1	1.9	1.86	2	3.5	2.46
65-69	19	16.2	3.73	8	14.8	5.24	11	17.5	5.27
70-74	35	32.0	5.41	17	36.0	8.72	18	29.0	6.83
75-79	63	65.3	8.23	28	71.3	13.48	35	61.2	10.34
80-84	109	167.0	16.00	44	183.2	27.61	65	157.6	19.55
85+	312	549.8	31.12	88	574.6	61.26	224	540.6	36.12
Total deaths	553			192			361		
Connecticut									
. Crude rate		19.2	-		13.7	-		24.4	-
. AAMR (1940 Std. mil.)		4.8	0.25		5.1	0.40		4.7	0.34
. AAMR (1970 Std. mil.)		9.1	0.41		9.6	0.70		8.9	0.52
. AAMR (2000 Std. mil.)		15.4	0.65		16.3	1.19		14.9	0.80
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.1	-		6.6	-		5.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	0	0.0	0.00	1	1.1	1.12
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.39	1	0.8	0.77	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	2	1.0	0.69	0	0.0	0.00	2	1.9	1.36
50-54	1	0.5	0.55	0	0.0	0.00	1	1.1	1.08
55-59	3	2.3	1.30	1	1.5	1.54	2	2.9	2.08
60-64	1	0.9	0.90	0	0.0	0.00	1	1.7	1.74
65-69	10	8.6	2.70	7	13.0	4.90	3	4.8	2.75
70-74	13	11.9	3.30	6	12.7	5.18	7	11.3	4.26
75-79	20	20.7	4.64	7	17.8	6.74	13	22.7	6.30
80-84	24	36.8	7.51	10	41.6	13.16	14	33.9	9.07
85+	71	125.1	14.85	16	104.5	26.12	55	132.7	17.90
Total deaths	147			48			99		
Connecticut									
. Crude rate		5.1	-		3.4	-		6.7	-
. AAMR (1940 Std. mil.)		1.6	0.16		1.5	0.23		1.6	0.23
. AAMR (1970 Std. mil.)		2.7	0.23		2.5	0.36		2.7	0.31
. AAMR (2000 Std. mil.)		4.1	0.34		3.8	0.56		4.3	0.44
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.3	-		2.6	-		2.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.1	0.77	1	1.1	1.06	1	1.1	1.12
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.5	0.54	0	0.0	0.00	1	1.1	1.10
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
30-34	5	2.1	0.96	3	2.6	1.48	2	1.7	1.22
35-39	4	1.5	0.77	4	3.1	1.55	0	0.0	0.00
40-44	21	8.7	1.90	10	8.4	2.66	11	9.0	2.71
45-49	15	7.3	1.88	7	6.9	2.61	8	7.7	2.71
50-54	20	11.0	2.46	8	8.9	3.16	12	12.9	3.73
55-59	30	22.6	4.12	20	30.9	6.90	10	14.7	4.65
60-64	29	26.1	4.85	19	35.4	8.13	10	17.4	5.51
65-69	88	75.3	8.02	43	79.7	12.15	45	71.4	10.65
70-74	158	144.4	11.49	86	182.0	19.62	72	115.9	13.65
75-79	272	282.0	17.10	126	321.0	28.59	146	255.2	21.12
80-84	360	551.6	29.07	134	557.8	48.18	226	547.9	36.45
85+	788	1388.5	49.46	199	1299.5	92.12	589	1421.4	58.57
Total deaths	1,794			661			1,133		
Connecticut									
. Crude rate		62.2	-		47.0	-		76.6	-
. AAMR (1940 Std. mil.)		19.3	0.56		21.1	0.91		17.8	0.71
. AAMR (1970 Std. mil.)		32.5	0.81		34.5	1.36		30.6	1.01
. AAMR (2000 Std. mil.)		50.6	1.20		52.5	2.08		48.6	1.47
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		25.9	-		27.9	-		24.1	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.6	0.94	2	2.1	1.49	1	1.1	1.12
5-9	1	0.5	0.51	1	1.0	0.99	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.7	0.65	1	1.3	1.26	0	0.0	0.00
25-29	4	2.2	1.08	0	0.0	0.00	4	4.4	2.18
30-34	3	1.3	0.74	3	2.6	1.48	0	0.0	0.00
35-39	6	2.3	0.95	5	3.9	1.73	1	0.8	0.77
40-44	11	4.6	1.38	8	6.7	2.38	3	2.5	1.42
45-49	9	4.4	1.46	7	6.9	2.61	2	1.9	1.36
50-54	8	4.4	1.55	5	5.6	2.50	3	3.2	1.87
55-59	22	16.6	3.53	15	23.1	5.97	7	10.3	3.89
60-64	19	17.1	3.93	12	22.4	6.46	7	12.2	4.61
65-69	42	35.9	5.54	27	50.0	9.63	15	23.8	6.15
70-74	66	60.3	7.43	42	88.9	13.71	24	38.6	7.88
75-79	145	150.3	12.48	87	221.6	23.76	58	101.4	13.31
80-84	227	347.8	23.08	118	491.2	45.22	109	264.3	25.31
85+	605	1066.0	43.34	212	1384.4	95.08	393	948.4	47.84
Total deaths	1,172			545			627		
Connecticut									
. Crude rate		40.6	-		38.8	-		42.4	-
. AAMR (1940 Std. mil.)		11.8	0.44		16.3	0.79		8.6	0.50
. AAMR (1970 Std. mil.)		20.4	0.63		28.0	1.22		15.7	0.70
. AAMR (2000 Std. mil.)		33.0	0.97		44.9	1.96		26.1	1.06
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.9	-		16.3	-		10.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
30-34	1	0.4	0.43	1	0.9	0.86	0	0.0	0.00
35-39	2	0.8	0.55	1	0.8	0.77	1	0.8	0.77
40-44	3	1.2	0.72	0	0.0	0.00	3	2.5	1.42
45-49	8	3.9	1.37	4	3.9	1.97	4	3.8	1.92
50-54	22	12.1	2.57	8	8.9	3.16	14	15.1	4.03
55-59	29	21.8	4.05	11	17.0	5.12	18	26.4	6.23
60-64	51	45.9	6.43	25	46.6	9.32	26	45.3	8.88
65-69	121	103.5	9.41	58	107.5	14.11	63	100.0	12.60
70-74	190	173.7	12.60	98	207.4	20.95	92	148.0	15.43
75-79	255	264.4	16.56	119	303.1	27.79	136	237.8	20.39
80-84	248	380.0	24.13	91	378.8	39.71	157	380.6	30.38
85+	297	523.3	30.37	121	790.1	71.83	176	424.7	32.02
Total deaths	1,229			539			690		
Connecticut									
. Crude rate		42.6	-		38.3	-		46.6	-
. AAMR (1940 Std. mil.)		16.8	0.56		18.5	0.87		16.0	0.75
. AAMR (1970 Std. mil.)		25.5	0.77		28.9	1.26		23.8	0.99
. AAMR (2000 Std. mil.)		35.2	1.01		41.2	1.80		32.5	1.26
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		21.1	-		26.1	-		17.8	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
30-34	1	0.4	0.43	1	0.9	0.86	0	0.0	0.00
35-39	2	0.8	0.55	1	0.8	0.77	1	0.8	0.77
40-44	1	0.4	0.41	0	0.0	0.00	1	0.8	0.82
45-49	2	1.0	0.69	0	0.0	0.00	2	1.9	1.36
50-54	9	4.9	1.65	4	4.5	2.24	5	5.4	2.41
55-59	4	3.0	1.50	1	1.5	1.54	3	4.4	2.54
60-64	4	3.6	1.80	1	1.9	1.86	3	5.2	3.02
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	3	2.7	1.58	1	2.1	2.12	2	3.2	2.28
75-79	7	7.3	2.74	0	0.0	0.00	7	12.2	4.63
80-84	8	12.3	4.33	2	8.3	5.89	6	14.5	5.94
85+	11	19.4	5.84	0	0.0	0.00	11	26.5	8.00
Total deaths	53			12			41		
Connecticut									
. Crude rate		1.8	-		0.9	-		2.8	-
. AAMR (1940 Std. mil.)		1.0	0.16		-	-		1.3	0.25
. AAMR (1970 Std. mil.)		1.2	0.18		-	-		1.6	0.28
. AAMR (2000 Std. mil.)		1.6	0.23		-	-		2.1	0.34
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.6	-		1.1	-		1.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	1.1	0.76	1	1.1	1.06	1	1.1	1.09
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	8	3.1	1.09	5	3.9	1.73	3	2.3	1.34
40-44	21	8.7	1.90	18	15.1	3.57	3	2.5	1.42
45-49	28	13.6	2.57	21	20.7	4.51	7	6.7	2.54
50-54	31	17.0	3.06	25	28.0	5.59	6	6.5	2.64
55-59	22	16.6	3.53	15	23.1	5.97	7	10.3	3.89
60-64	30	27.0	4.93	21	39.2	8.55	9	15.7	5.23
65-69	35	29.9	5.06	24	44.5	9.08	11	17.5	5.27
70-74	29	26.5	4.92	14	29.6	7.92	15	24.1	6.23
75-79	34	35.2	6.05	13	33.1	9.18	21	36.7	8.01
80-84	23	35.2	7.35	10	41.6	13.16	13	31.5	8.74
85+	11	19.4	5.84	4	26.1	13.06	7	16.9	6.38
Total deaths	274			171			103		
Connecticut									
. Crude rate		9.5	-		12.2	-		7.0	-
. AAMR (1940 Std. mil.)		6.4	0.42		9.2	0.73		3.8	0.43
. AAMR (1970 Std. mil.)		7.4	0.46		10.3	0.80		4.7	0.49
. AAMR (2000 Std. mil.)		8.7	0.53		12.0	0.92		5.6	0.57
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.3	-		10.5	-		4.5	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis, nephrotic syndrome/nephrosis (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	0	0.0	0.00	1	1.1	1.12
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.59	1	1.1	1.14	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
30-34	1	0.4	0.43	0	0.0	0.00	1	0.9	0.86
35-39	1	0.4	0.39	0	0.0	0.00	1	0.8	0.77
40-44	4	1.7	0.83	2	1.7	1.19	2	1.6	1.16
45-49	4	1.9	0.97	3	3.0	1.71	1	1.0	0.96
50-54	6	3.3	1.34	5	5.6	2.50	1	1.1	1.08
55-59	7	5.3	1.99	3	4.6	2.67	4	5.9	2.94
60-64	7	6.3	2.38	2	3.7	2.64	5	8.7	3.89
65-69	11	9.4	2.84	5	9.3	4.14	6	9.5	3.89
70-74	34	31.1	5.33	18	38.1	8.98	16	25.7	6.44
75-79	47	48.7	7.11	28	71.3	13.48	19	33.2	7.62
80-84	67	102.7	12.54	39	162.3	25.99	28	67.9	12.83
85+	121	213.2	19.38	55	359.1	48.43	66	159.3	19.61
Total deaths	313			162			151		
Connecticut									
. Crude rate		10.8	-		11.5	-		10.2	-
. AAMR (1940 Std. mil.)		3.7	0.26		4.9	0.43		3.0	0.33
. AAMR (1970 Std. mil.)		5.9	0.36		8.3	0.66		4.6	0.43
. AAMR (2000 Std. mil.)		8.9	0.51		13.1	1.05		6.8	0.57
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.4	-		5.5	-		3.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----End stage renal disease (ESRD) (581-588,591,593.8,593.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.6	0.94	1	1.1	1.06	2	2.2	1.59
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.59	1	1.1	1.14	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
30-34	1	0.4	0.43	0	0.0	0.00	1	0.9	0.86
35-39	1	0.4	0.39	0	0.0	0.00	1	0.8	0.77
40-44	4	1.7	0.83	2	1.7	1.19	2	1.6	1.16
45-49	5	2.4	1.09	4	3.9	1.97	1	1.0	0.96
50-54	7	3.8	1.45	6	6.7	2.74	1	1.1	1.08
55-59	8	6.0	2.13	4	6.2	3.09	4	5.9	2.94
60-64	10	9.0	2.85	4	7.5	3.73	6	10.5	4.27
65-69	16	13.7	3.42	9	16.7	5.56	7	11.1	4.20
70-74	40	36.6	5.78	19	40.2	9.22	21	33.8	7.37
75-79	60	62.2	8.03	36	91.7	15.28	24	42.0	8.56
80-84	75	114.9	13.27	45	187.3	27.92	30	72.7	13.28
85+	131	230.8	20.17	61	398.3	51.00	70	168.9	20.19
Total deaths	363			193			170		
Connecticut									
. Crude rate		12.6	-		13.7	-		11.5	-
. AAMR (1940 Std. mil.)		4.5	0.29		6.1	0.49		3.5	0.36
. AAMR (1970 Std. mil.)		7.0	0.39		10.0	0.73		5.4	0.47
. AAMR (2000 Std. mil.)		10.4	0.55		15.5	1.13		7.7	0.61
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.5	-		6.8	-		4.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	16	8.7	2.18	10	10.6	3.34	6	6.7	2.75
5-9	5	2.5	1.13	5	4.9	2.20	0	0.0	0.00
10-14	15	8.1	2.09	6	6.3	2.59	9	9.9	3.30
15-19	51	30.1	4.21	28	31.9	6.04	23	28.1	5.85
20-24	49	31.9	4.56	35	44.2	7.47	14	18.8	5.02
25-29	54	29.0	3.95	41	43.6	6.81	13	14.1	3.92
30-34	71	30.5	3.62	57	48.8	6.46	14	12.1	3.23
35-39	73	28.2	3.30	61	47.1	6.04	12	9.3	2.68
40-44	65	27.0	3.34	51	42.9	6.00	14	11.5	3.06
45-49	59	28.7	3.73	45	44.3	6.61	14	13.4	3.59
50-54	34	18.7	3.20	23	25.7	5.36	11	11.9	3.58
55-59	23	17.3	3.61	11	17.0	5.12	12	17.6	5.09
60-64	30	27.0	4.93	19	35.4	8.13	11	19.2	5.78
65-69	47	40.2	5.86	34	63.0	10.81	13	20.6	5.72
70-74	52	47.5	6.59	29	61.4	11.39	23	37.0	7.72
75-79	76	78.8	9.04	41	104.4	16.31	35	61.2	10.34
80-84	101	154.7	15.40	50	208.1	29.43	51	123.6	17.31
85+	128	225.5	19.94	57	372.2	49.30	71	171.3	20.33
Total deaths	949			603			346		
Connecticut									
. Crude rate		32.9	-		42.9	-		23.4	-
. AAMR (1940 Std. mil.)		25.0	0.95		35.1	1.57		15.5	1.07
. AAMR (1970 Std. mil.)		26.3	0.93		36.6	1.56		17.2	1.08
. AAMR (2000 Std. mil.)		30.6	1.00		43.8	1.80		19.6	1.10
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		30.1	-		42.8	-		17.8	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Motor vehicle accidents (E810-E825)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	0	0.0	0.00	1	1.1	1.12
5-9	3	1.5	0.88	3	3.0	1.71	0	0.0	0.00
10-14	12	6.5	1.87	4	4.2	2.11	8	8.8	3.11
15-19	42	24.8	3.82	21	24.0	5.23	21	25.6	5.59
20-24	27	17.6	3.38	18	22.7	5.36	9	12.1	4.03
25-29	30	16.1	2.95	22	23.4	4.99	8	8.7	3.08
30-34	26	11.2	2.19	17	14.5	3.53	9	7.8	2.59
35-39	18	7.0	1.64	11	8.5	2.56	7	5.4	2.04
40-44	23	9.5	1.99	16	13.5	3.36	7	5.7	2.17
45-49	21	10.2	2.23	15	14.8	3.81	6	5.8	2.35
50-54	18	9.9	2.33	12	13.4	3.87	6	6.5	2.64
55-59	6	4.5	1.84	3	4.6	2.67	3	4.4	2.54
60-64	12	10.8	3.12	8	14.9	5.27	4	7.0	3.48
65-69	13	11.1	3.08	11	20.4	6.15	2	3.2	2.25
70-74	17	15.5	3.77	10	21.2	6.69	7	11.3	4.26
75-79	15	15.6	4.02	7	17.8	6.74	8	14.0	4.94
80-84	14	21.4	5.73	6	25.0	10.20	8	19.4	6.86
85+	13	22.9	6.35	5	32.6	14.60	8	19.3	6.83
Total deaths	311			189			122		
Connecticut									
. Crude rate		10.8	-		13.4	-		8.2	-
. AAMR (1940 Std. mil.)		10.7	0.67		13.3	1.03		8.3	0.85
. AAMR (1970 Std. mil.)		10.6	0.65		13.0	1.00		8.3	0.83
. AAMR (2000 Std. mil.)		10.8	0.62		13.6	1.00		8.2	0.76
. HP 2000 (1940 Std. mil.)		10.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		16.0	-		21.7	-		10.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.51	1	1.0	0.99	0	0.0	0.00
10-14	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
15-19	1	0.6	0.59	1	1.1	1.14	0	0.0	0.00
20-24	4	2.6	1.30	4	5.1	2.53	0	0.0	0.00
25-29	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
30-34	2	0.9	0.61	2	1.7	1.21	0	0.0	0.00
35-39	1	0.4	0.39	1	0.8	0.77	0	0.0	0.00
40-44	1	0.4	0.41	1	0.8	0.84	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	2	1.1	0.78	2	2.2	1.58	0	0.0	0.00
55-59	3	2.3	1.30	3	4.6	2.67	0	0.0	0.00
60-64	4	3.6	1.80	4	7.5	3.73	0	0.0	0.00
65-69	14	12.0	3.20	9	16.7	5.56	5	7.9	3.55
70-74	13	11.9	3.30	7	14.8	5.60	6	9.7	3.94
75-79	28	29.0	5.49	20	50.9	11.39	8	14.0	4.94
80-84	40	61.3	9.69	23	95.7	19.96	17	41.2	10.00
85+	67	118.1	14.42	31	202.4	36.36	36	86.9	14.48
Total deaths	183			111			72		
Connecticut									
. Crude rate		6.3	-		7.9	-		4.9	-
. AAMR (1940 Std. mil.)		2.4	0.24		4.1	0.47		1.1	0.16
. AAMR (1970 Std. mil.)		3.7	0.30		6.2	0.61		1.9	0.24
. AAMR (2000 Std. mil.)		5.3	0.39		8.8	0.85		3.0	0.36
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.6	-		3.6	-		1.7	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.2	1.09	3	3.2	1.83	1	1.1	1.12
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	1.2	0.83	2	2.3	1.61	0	0.0	0.00
20-24	2	1.3	0.92	2	2.5	1.79	0	0.0	0.00
25-29	2	1.1	0.76	2	2.1	1.50	0	0.0	0.00
30-34	3	1.3	0.74	3	2.6	1.48	0	0.0	0.00
35-39	4	1.5	0.77	4	3.1	1.55	0	0.0	0.00
40-44	1	0.4	0.41	1	0.8	0.84	0	0.0	0.00
45-49	1	0.5	0.49	1	1.0	0.98	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	1	0.8	0.75	1	1.5	1.54	0	0.0	0.00
60-64	1	0.9	0.90	0	0.0	0.00	1	1.7	1.74
65-69	1	0.9	0.86	1	1.9	1.85	0	0.0	0.00
70-74	1	0.9	0.91	1	2.1	2.12	0	0.0	0.00
75-79	2	2.1	1.47	2	5.1	3.60	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	1	1.8	1.76	1	6.5	6.53	0	0.0	0.00
Total deaths	26			24			2		
Connecticut									
. Crude rate		0.9	-		1.7	-		0.1	-
. AAMR (1940 Std. mil.)		0.9	0.19		1.6	0.36		-	-
. AAMR (1970 Std. mil.)		0.9	0.18		1.6	0.34		-	-
. AAMR (2000 Std. mil.)		0.9	0.18		1.7	0.35		-	-
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.6	-		2.5	-		0.4	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.1	0.77	1	1.1	1.06	1	1.1	1.12
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.59	0	0.0	0.00	1	1.2	1.22
20-24	1	0.7	0.65	1	1.3	1.26	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.43	0	0.0	0.00	1	0.9	0.86
35-39	1	0.4	0.39	1	0.8	0.77	0	0.0	0.00
40-44	2	0.8	0.59	1	0.8	0.84	1	0.8	0.82
45-49	2	1.0	0.69	1	1.0	0.98	1	1.0	0.96
50-54	2	1.1	0.78	1	1.1	1.12	1	1.1	1.08
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	3	2.7	1.56	2	3.7	2.64	1	1.7	1.74
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	1	0.9	0.91	0	0.0	0.00	1	1.6	1.61
75-79	4	4.1	2.07	1	2.5	2.55	3	5.2	3.03
80-84	2	3.1	2.17	2	8.3	5.89	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	22			11			11		
Connecticut									
. Crude rate		0.8	-		0.8	-		0.7	-
. AAMR (1940 Std. mil.)		0.6	0.15		-	-		-	-
. AAMR (1970 Std. mil.)		0.7	0.15		-	-		-	-
. AAMR (2000 Std. mil.)		0.7	0.16		-	-		-	-
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.9	-		1.4	-		0.9	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	1.6	0.93	3	3.2	1.83	0	0.0	0.00
15-19	8	4.7	1.67	6	6.8	2.79	2	2.4	1.73
20-24	12	7.8	2.25	10	12.6	4.00	2	2.7	1.90
25-29	16	8.6	2.15	12	12.8	3.68	4	4.4	2.18
30-34	33	14.2	2.47	29	24.8	4.61	4	3.4	1.72
35-39	30	11.6	2.12	21	16.2	3.54	9	7.0	2.32
40-44	36	14.9	2.49	24	20.2	4.12	12	9.8	2.83
45-49	23	11.2	2.33	18	17.7	4.18	5	4.8	2.15
50-54	21	11.5	2.52	15	16.8	4.33	6	6.5	2.64
55-59	9	6.8	2.26	7	10.8	4.08	2	2.9	2.08
60-64	14	12.6	3.37	11	20.5	6.18	3	5.2	3.02
65-69	10	8.6	2.70	8	14.8	5.24	2	3.2	2.25
70-74	15	13.7	3.54	10	21.2	6.69	5	8.0	3.60
75-79	9	9.3	3.11	9	22.9	7.64	0	0.0	0.00
80-84	12	18.4	5.31	8	33.3	11.77	4	9.7	4.85
85+	6	10.6	4.32	6	39.2	16.00	0	0.0	0.00
Total deaths	257			197			60		
Connecticut									
. Crude rate		8.9	-		14.0	-		4.1	-
. AAMR (1940 Std. mil.)		7.8	0.52		12.2	0.92		3.5	0.49
. AAMR (1970 Std. mil.)		7.4	0.49		11.9	0.88		3.3	0.45
. AAMR (2000 Std. mil.)		8.5	0.53		13.8	0.99		3.9	0.51
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.5	-		17.4	-		4.0	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.2	1.09	3	3.2	1.83	1	1.1	1.12
5-9	2	1.0	0.72	1	1.0	0.99	1	1.0	1.04
10-14	2	1.1	0.76	2	2.1	1.49	0	0.0	0.00
15-19	7	4.1	1.56	5	5.7	2.55	2	2.4	1.73
20-24	10	6.5	2.06	10	12.6	4.00	0	0.0	0.00
25-29	2	1.1	0.76	2	2.1	1.50	0	0.0	0.00
30-34	3	1.3	0.74	2	1.7	1.21	1	0.9	0.86
35-39	6	2.3	0.95	3	2.3	1.34	3	2.3	1.34
40-44	5	2.1	0.93	3	2.5	1.46	2	1.6	1.16
45-49	2	1.0	0.69	1	1.0	0.98	1	1.0	0.96
50-54	4	2.2	1.10	4	4.5	2.24	0	0.0	0.00
55-59	3	2.3	1.30	1	1.5	1.54	2	2.9	2.08
60-64	1	0.9	0.90	1	1.9	1.86	0	0.0	0.00
65-69	3	2.6	1.48	2	3.7	2.62	1	1.6	1.59
70-74	1	0.9	0.91	1	2.1	2.12	0	0.0	0.00
75-79	1	1.0	1.04	1	2.5	2.55	0	0.0	0.00
80-84	1	1.5	1.53	0	0.0	0.00	1	2.4	2.42
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	57			42			15		
Connecticut									
. Crude rate		2.0	-		3.0	-		1.0	-
. AAMR (1940 Std. mil.)		2.2	0.31		3.4	0.55		1.0	0.27
. AAMR (1970 Std. mil.)		2.2	0.31		3.3	0.54		1.0	0.27
. AAMR (2000 Std. mil.)		2.1	0.28		3.1	0.48		1.0	0.27
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.1	-		12.5	-		3.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All injury (E800-E999)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	20	10.9	2.43	13	13.7	3.81	7	7.8	2.97
5-9	7	3.5	1.34	6	5.9	2.41	1	1.0	1.04
10-14	20	10.8	2.41	11	11.6	3.50	9	9.9	3.30
15-19	66	38.9	4.79	39	44.5	7.12	27	33.0	6.34
20-24	72	46.9	5.52	56	70.7	9.45	16	21.5	5.37
25-29	79	42.5	4.78	59	62.7	8.16	20	21.8	4.87
30-34	112	48.1	4.54	90	77.0	8.12	22	19.0	4.04
35-39	117	45.2	4.18	91	70.3	7.37	26	20.1	3.94
40-44	116	48.1	4.47	85	71.5	7.75	31	25.4	4.56
45-49	91	44.2	4.64	68	67.0	8.12	23	22.1	4.60
50-54	62	34.0	4.32	45	50.3	7.50	17	18.3	4.44
55-59	35	26.3	4.45	19	29.3	6.72	16	23.5	5.88
60-64	45	40.5	6.04	31	57.8	10.38	14	24.4	6.52
65-69	61	52.2	6.68	45	83.4	12.43	16	25.4	6.35
70-74	68	62.2	7.54	40	84.6	13.38	28	45.1	8.52
75-79	86	89.2	9.61	51	129.9	18.19	35	61.2	10.34
80-84	115	176.2	16.43	59	245.6	31.97	56	135.8	18.14
85+	134	236.1	20.40	63	411.4	51.83	71	171.3	20.33
Total deaths	1,306			871			435		
Connecticut									
. Crude rate		45.3	-		61.9	-		29.4	-
. AAMR (1940 Std. mil.)		36.4	1.15		52.6	1.94		20.9	1.23
. AAMR (1970 Std. mil.)		37.0	1.11		53.4	1.90		22.2	1.22
. AAMR (2000 Std. mil.)		42.6	1.19		62.7	2.14		25.5	1.26
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		49.8	-		74.7	-		25.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	0	0.0	0.00	1	1.1	1.12
5-9	2	1.0	0.72	2	2.0	1.39	0	0.0	0.00
10-14	12	6.5	1.87	4	4.2	2.11	8	8.8	3.11
15-19	41	24.2	3.78	20	22.8	5.10	21	25.6	5.59
20-24	25	16.3	3.25	16	20.2	5.05	9	12.1	4.03
25-29	29	15.6	2.90	22	23.4	4.99	7	7.6	2.88
30-34	26	11.2	2.19	16	13.7	3.42	10	8.6	2.73
35-39	17	6.6	1.59	11	8.5	2.56	6	4.6	1.89
40-44	22	9.1	1.95	15	12.6	3.26	7	5.7	2.17
45-49	21	10.2	2.23	15	14.8	3.81	6	5.8	2.35
50-54	18	9.9	2.33	12	13.4	3.87	6	6.5	2.64
55-59	6	4.5	1.84	3	4.6	2.67	3	4.4	2.54
60-64	12	10.8	3.12	8	14.9	5.27	4	7.0	3.48
65-69	13	11.1	3.08	11	20.4	6.15	2	3.2	2.25
70-74	18	16.5	3.88	11	23.3	7.02	7	11.3	4.26
75-79	15	15.6	4.02	7	17.8	6.74	8	14.0	4.94
80-84	14	21.4	5.73	6	25.0	10.20	8	19.4	6.86
85+	13	22.9	6.35	5	32.6	14.60	8	19.3	6.83
Total deaths	305			184			121		
Connecticut									
. Crude rate		10.6	-		13.1	-		8.2	-
. AAMR (1940 Std. mil.)		10.4	0.66		12.8	1.01		8.2	0.85
. AAMR (1970 Std. mil.)		10.3	0.64		12.6	0.97		8.2	0.83
. AAMR (2000 Std. mil.)		10.5	0.61		13.2	0.98		8.1	0.76
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.6	-		21.3	-		10.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.51	1	1.0	0.99	0	0.0	0.00
10-14	1	0.5	0.54	1	1.1	1.06	0	0.0	0.00
15-19	1	0.6	0.59	1	1.1	1.14	0	0.0	0.00
20-24	3	2.0	1.13	3	3.8	2.19	0	0.0	0.00
25-29	2	1.1	0.76	1	1.1	1.06	1	1.1	1.09
30-34	3	1.3	0.74	3	2.6	1.48	0	0.0	0.00
35-39	1	0.4	0.39	1	0.8	0.77	0	0.0	0.00
40-44	1	0.4	0.41	1	0.8	0.84	0	0.0	0.00
45-49	1	0.5	0.49	1	1.0	0.98	0	0.0	0.00
50-54	2	1.1	0.78	2	2.2	1.58	0	0.0	0.00
55-59	3	2.3	1.30	3	4.6	2.67	0	0.0	0.00
60-64	6	5.4	2.21	6	11.2	4.57	0	0.0	0.00
65-69	12	10.3	2.96	8	14.8	5.24	4	6.4	3.18
70-74	9	8.2	2.74	6	12.7	5.18	3	4.8	2.79
75-79	23	23.8	4.97	17	43.3	10.50	6	10.5	4.28
80-84	25	38.3	7.66	17	70.8	17.16	8	19.4	6.86
85+	38	67.0	10.86	17	111.0	26.92	21	50.7	11.06
Total deaths	132			89			43		
Connecticut									
. Crude rate		4.6	-		6.3	-		2.9	-
. AAMR (1940 Std. mil.)		2.1	0.23		3.7	0.45		0.7	0.16
. AAMR (1970 Std. mil.)		2.9	0.28		5.1	0.56		1.2	0.20
. AAMR (2000 Std. mil.)		3.9	0.34		6.8	0.73		1.9	0.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.5	-		3.6	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	7	4.1	1.56	7	8.0	3.02	0	0.0	0.00
20-24	10	6.5	2.06	10	12.6	4.00	0	0.0	0.00
25-29	10	5.4	1.70	10	10.6	3.36	0	0.0	0.00
30-34	17	7.3	1.77	17	14.5	3.53	0	0.0	0.00
35-39	19	7.3	1.68	15	11.6	2.99	4	3.1	1.54
40-44	15	6.2	1.61	11	9.2	2.79	4	3.3	1.64
45-49	7	3.4	1.29	5	4.9	2.20	2	1.9	1.36
50-54	11	6.0	1.82	10	11.2	3.54	1	1.1	1.08
55-59	4	3.0	1.50	3	4.6	2.67	1	1.5	1.47
60-64	8	7.2	2.55	8	14.9	5.27	0	0.0	0.00
65-69	5	4.3	1.91	4	7.4	3.71	1	1.6	1.59
70-74	6	5.5	2.24	6	12.7	5.18	0	0.0	0.00
75-79	1	1.0	1.04	1	2.5	2.55	0	0.0	0.00
80-84	6	9.2	3.75	6	25.0	10.20	0	0.0	0.00
85+	4	7.0	3.52	4	26.1	13.06	0	0.0	0.00
Total deaths	130			117			13		
Connecticut									
. Crude rate		4.5	-		8.3	-		0.9	-
. AAMR (1940 Std. mil.)		4.2	0.39		7.7	0.76		-	-
. AAMR (1970 Std. mil.)		3.9	0.36		7.4	0.71		-	-
. AAMR (2000 Std. mil.)		4.4	0.39		8.3	0.77		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.2	-		21.1	-		3.3	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	3	1.8	1.02	3	3.4	1.98	0	0.0	0.00
20-24	2	1.3	0.92	2	2.5	1.79	0	0.0	0.00
25-29	7	3.8	1.42	7	7.4	2.81	0	0.0	0.00
30-34	16	6.9	1.72	16	13.7	3.42	0	0.0	0.00
35-39	15	5.8	1.50	12	9.3	2.68	3	2.3	1.34
40-44	12	5.0	1.44	9	7.6	2.52	3	2.5	1.42
45-49	7	3.4	1.29	5	4.9	2.20	2	1.9	1.36
50-54	8	4.4	1.55	7	7.8	2.96	1	1.1	1.08
55-59	3	2.3	1.30	3	4.6	2.67	0	0.0	0.00
60-64	7	6.3	2.38	7	13.1	4.93	0	0.0	0.00
65-69	4	3.4	1.71	4	7.4	3.71	0	0.0	0.00
70-74	6	5.5	2.24	6	12.7	5.18	0	0.0	0.00
75-79	1	1.0	1.04	1	2.5	2.55	0	0.0	0.00
80-84	6	9.2	3.75	6	25.0	10.20	0	0.0	0.00
85+	4	7.0	3.52	4	26.1	13.06	0	0.0	0.00
Total deaths	101			92			9		
Connecticut									
. Crude rate		3.5	-		6.5	-		0.6	-
. AAMR (1940 Std. mil.)		3.0	0.31		5.5	0.61		-	-
. AAMR (1970 Std. mil.)		2.8	0.29		5.4	0.58		-	-
. AAMR (2000 Std. mil.)		3.3	0.33		6.4	0.67		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.9	-		10.7	-		1.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	4	2.4	1.18	4	4.6	2.28	0	0.0	0.00
20-24	8	5.2	1.84	8	10.1	3.57	0	0.0	0.00
25-29	2	1.1	0.76	2	2.1	1.50	0	0.0	0.00
30-34	1	0.4	0.43	1	0.9	0.86	0	0.0	0.00
35-39	4	1.5	0.77	3	2.3	1.34	1	0.8	0.77
40-44	3	1.2	0.72	2	1.7	1.19	1	0.8	0.82
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	3	1.6	0.95	3	3.4	1.94	0	0.0	0.00
55-59	1	0.8	0.75	0	0.0	0.00	1	1.5	1.47
60-64	1	0.9	0.90	1	1.9	1.86	0	0.0	0.00
65-69	1	0.9	0.86	0	0.0	0.00	1	1.6	1.59
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	28			24			4		
Connecticut									
. Crude rate		1.0	-		1.7	-		0.3	-
. AAMR (1940 Std. mil.)		1.2	0.23		2.1	0.44		-	-
. AAMR (1970 Std. mil.)		1.1	0.22		1.9	0.41		-	-
. AAMR (2000 Std. mil.)		1.0	0.20		1.8	0.37		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.5	-		9.3	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	7	3.0	1.14	7	6.0	2.26	0	0.0	0.00
35-39	9	3.5	1.16	5	3.9	1.73	4	3.1	1.54
40-44	11	4.6	1.38	8	6.7	2.38	3	2.5	1.42
45-49	19	9.2	2.12	17	16.7	4.06	2	1.9	1.36
50-54	18	9.9	2.33	15	16.8	4.33	3	3.2	1.87
55-59	8	6.0	2.13	6	9.3	3.78	2	2.9	2.08
60-64	18	16.2	3.82	15	28.0	7.22	3	5.2	3.02
65-69	13	11.1	3.08	10	18.5	5.86	3	4.8	2.75
70-74	6	5.5	2.24	3	6.3	3.66	3	4.8	2.79
75-79	9	9.3	3.11	5	12.7	5.70	4	7.0	3.50
80-84	4	6.1	3.06	3	12.5	7.21	1	2.4	2.42
85+	2	3.5	2.49	2	13.1	9.23	0	0.0	0.00
Total deaths	124			96			28		
Connecticut									
. Crude rate		4.3	-		6.8	-		1.9	-
. AAMR (1940 Std. mil.)		3.3	0.31		5.5	0.57		1.3	0.27
. AAMR (1970 Std. mil.)		3.5	0.32		5.8	0.60		1.4	0.28
. AAMR (2000 Std. mil.)		4.0	0.36		6.7	0.69		1.7	0.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.3	-		9.9	-		3.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1997

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	1.2	0.83	1	1.1	1.14	1	1.2	1.22
20-24	13	8.5	2.35	11	13.9	4.19	2	2.7	1.90
25-29	21	11.3	2.46	15	15.9	4.12	6	6.5	2.67
30-34	43	18.5	2.82	33	28.2	4.92	10	8.6	2.73
35-39	54	20.9	2.84	48	37.1	5.35	6	4.6	1.89
40-44	44	18.2	2.75	31	26.1	4.68	13	10.6	2.95
45-49	30	14.6	2.66	22	21.7	4.62	8	7.7	2.71
50-54	8	4.4	1.55	6	6.7	2.74	2	2.2	1.52
55-59	9	6.8	2.26	3	4.6	2.67	6	8.8	3.60
60-64	1	0.9	0.90	0	0.0	0.00	1	1.7	1.74
65-69	2	1.7	1.21	0	0.0	0.00	2	3.2	2.25
70-74	5	4.6	2.04	3	6.3	3.66	2	3.2	2.28
75-79	4	4.1	2.07	2	5.1	3.60	2	3.5	2.47
80-84	3	4.6	2.65	0	0.0	0.00	3	7.3	4.20
85+	2	3.5	2.49	1	6.5	6.53	1	2.4	2.41
Total deaths	241			176			65		
Connecticut									
. Crude rate		8.4	-		12.5	-		4.4	-
. AAMR (1940 Std. mil.)		7.7	0.52		11.4	0.89		3.9	0.52
. AAMR (1970 Std. mil.)		6.6	0.44		9.6	0.76		3.6	0.47
. AAMR (2000 Std. mil.)		8.0	0.52		11.8	0.90		4.2	0.53
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.6	-		7.8	-		3.3	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is higher
. THE U.S. HP 2000 target: CT rate is higher